Variceal hemorrhage of the colon secondary to pancreatic cancer

Fig. 1 Endoscopic view of colonic lumen in a 71-year-old woman with previous segmental colonic resection for adenocarcinoma, who presented with hematochezia. Colonoscopy showed colon varices extending proximal to the anastomosis, with superficial venules (red wale markings).

A 71-year-old woman was admitted due to recurrent lower gastrointestinal bleeding. She had undergone segmental colonic resection and adjuvant chemotherapy for colon adenocarcinoma 10 years earlier, without recurrence on follow-up. She reported experiencing several self-limited episodes of hematochezia over the past 6 months, without other symptoms.

Blood tests showed iron deficiency anemia (hemoglobin 9.9 g/dL). Colonoscopy revealed dilated, tortuous, bluish vessels protruding into the lumen and extending proximal to the anastomosis, numerous superficial venules, and fresh blood and clots in the lumen (● Fig. 1). Hemostasis was achieved by adrenaline injection and hemostatic clips. Computed tomography angiography demonstrated prominent collateral vessels near the colonic anastomosis, without signs of thrombosis (● Fig. 2). In addition, a nodular density adjacent to the pancreatic uncinate process was noted, with superior mesenteric vessels involvement. Subsequent endoscopic ultrasound (EUS) showed a 31-mm hypoechoic pancreatic head mass (● Fig. 3), with invasion of the spleno-portal confluence, generating a “stop” image on Doppler ultrasound (● Fig. 4). Transbulbar EUS-guided fine-needle aspiration (22-gauge needle) was performed. Pathologic analysis revealed pan-

Fig. 2 Computed tomography image obtained after the administration of intravenous contrast showing prominent collateral vessels adjacent to the colonic anastomosis (white arrow), without signs of thrombosis. In addition, a nodular density adjacent to the pancreatic uncinate process was observed (black arrow).

Fig. 3 Endoscopic ultrasound image showing a hypoechoic pancreatic head mass, with 31 mm dimension and irregular margins.

Fig. 4 Endoscopic Doppler ultrasound image demonstrating invasion of the spleno-portal confluence by the pancreatic mass, which generates a “stop” image on Doppler sign.

creatic ductal adenocarcinoma (● Fig. 5). Due to recurrent colonic bleeding, a vascular stent was placed through percutaneous transhepatic selective portography (● Fig. 6).

Colonic varices are a very rare cause of lower gastrointestinal bleeding, with a reported incidence of 0.07% [1]. Portal hy-

pertension is the most common etiology. Uncommon causes are congestive heart failure, mesenteric vein thrombosis, pancreatitis with splenic vein thrombosis, adhesions and, rarely, mesenteric vein obstruction [2]. It should prompt thorough evaluation, but can be idiopathic [1–3]. The present case is a peculiar con-

Fig. 5 Photomicrograph of the endoscopic ultrasound-guided fine-needle puncture specimen, demonstrating epithelial cells in cohesive papillary aggregates, with altered nucleocytoplasmic ratio, consistent with the diagnosis of pancreatic ductal adenocarcinoma. (Hematoxylin and eosin, $\times 40$).

Fig. 6 Selective transhepatic portography showing superior mesenteric vein obstruction over a length of 3 cm and retrograde filling of varicose vein knot. A vascular stent was placed.

dition – a patient with previous segmental colectomy due to carcinoma, presenting with recurrent hematochezia as a result of variceal hemorrhage due to a second primary (pancreatic) carcinoma and

mesenteric obstruction. This case highlights the importance of considering colonic varices in the differential diagnosis of lower gastrointestinal bleeding and the importance of thorough investigation.

Endoscopy_UCTN_Code_CCL_1AD_2AF

Competing interests: None

Teresa Pinto-Pais¹, Sónia Sousa Fernandes¹, Luísa Proença¹, Pedro Roquete², Tiago Pereira³, João Carvalho¹, José Fraga¹

¹ Department of Gastroenterology and Hepatology, Centro Hospitalar de Gaia/Espinho, Vila Nova de Gaia, Portugal

² Department of Histopathology, Laboratório Rodrigues Pereira, Porto, Portugal

³ Department of Radiology, Centro Hospitalar de Gaia/Espinho, Vila Nova de Gaia, Portugal

References

- 1 Han JH, Jeon WJ, Chae HB et al. A case of idiopathic colonic varices: a rare cause of hematochezia misconceived as tumor. *World J Gastroenterol* 2006; 12: 2629–2632
- 2 Sohn W, Lee HL, Lee KN. Variceal hemorrhage of ascending colon. *Clin Gastroenterol Hepatol* 2012; 10: A24
- 3 Francois F, Tadros C, Diehl D. Pan-colonic varices and idiopathic portal hypertension. *J Gastrointest Liver Dis* 2007; 16: 325–328

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377763>
Endoscopy 2014; 46: E500–E501
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Teresa Pinto-Pais, MD

Department of Gastroenterology and Hepatology
Centro Hospitalar de Gaia/Espinho
Rua Conceição Fernandes
Vila Nova de Gaia 4434-502
Portugal
Fax: +351-227865100
teresapintopais@gmail.com