Endoscopic placement of a fully covered self-expandable metallic stent to treat an arteriobiliary fistula

Fig. 1 Angiogram showing an arteriobiliary fistula after contrast injection via the catheter-port system for intra-arterial chemotherapy.

Fig. 2 Endoscopic image showing no hemobilia in the ampulla of Vater.

Fig. 3 Radiograph showing the fully covered self-expandable metallic stent placed over a guidewire lying across the arteriobiliary fistula.

Arteriobiliary fistula is an uncommon complication resulting from hepatic intra-arterial chemotherapy. We present a case of successful endoscopic intervention using a fully covered self-expandable

metallic stent (SEMS) to treat an iatrogenic arteriobiliary fistula.

A 71-year-old man with multiple liver metastases from pancreatic cancer was admitted. He had undergone distal pan-

createctomy. He had also undergone chemotherapy with multiple regimens during the past 43 months and percutaneous transfemoral implantation of the catheter-port system for intra-arterial chemotherapy to treat his liver metastases. Three months after catheter placement, he developed sepsis, with positive results on bacterial blood cultures. The reason for the sepsis could not be determined using computed tomography. However, angiography via the catheter revealed a fistula between the hepatic artery and the extra-hepatic bile duct (● **Fig. 1**), indicating the fistula as the cause of the sepsis. Because of catheter-related thrombosis, the implanted catheter-port system could not be removed, so we decided to seal the arteriobiliary fistula endoscopically. Duodenoscopy revealed no hemobilia in the ampulla of Vater (● **Fig. 2**). A fully covered SEMS (ComVi, 10×80 mm; Taewoong-Medical Co., Ltd., Gyeonggi-do, Korea) (● **Fig. 3**) was inserted, and the fistula was sealed successfully without any complications (● **Fig. 4**). The patient did not require any treatment for the arteriobiliary fistula thereafter.

Arteriobiliary fistula after intra-arterial chemotherapy has been reported in only a few cases [1–4]. Embolization of the catheter track or the branch of the hepatic artery is a reported treatment [4]. More recently, percutaneous transhepatic placement of a covered SEMS to treat an intrahepatic arteriobiliary fistula has been described [5]. To our knowledge, however, ours is the first report of an arteriobiliary fistula treated endoscopically using a fully covered SEMS. Endoscopic placement of a fully covered SEMS should be recognized as a treatment option for arteriobiliary fistulas, as it could obviate the need for an arteriogram.

Endoscopy_UCTN_Code_TTT_1AR_2AF

Competing interests: None

Hiroshi Kawakami¹, Munenori Okamoto², Masaki Kuwatani¹, Yoshimasa Kubota¹, Kazumichi Kawakubo¹, Yoko Abe¹, Shuhei Kawahata¹, Kimitoshi Kubo¹, Naoya Sakamoto¹

¹ Department of Gastroenterology and Hepatology, Hokkaido University Graduate School of Medicine, Sapporo, Japan

² Department of Internal Medicine, Sapporo Century Hospital, Sapporo, Japan

Fig. 4 Radiograph showing the arteriobiliary fistula successfully sealed using the fully covered self-expandable metallic stent (inset: endoscopic view of the stent passing through the ampulla of Vater to the duodenum).

References

- 1 Rosenberg HD, Wile AG, Aufrichtig D et al. Hepatic artery–biliary fistula. An unusual complication of infusion therapy. *Gastrointest Radiol* 1983; 8: 37–40
- 2 Noda M, Kusunoki M, Yanagai H et al. Hepatic artery–biliary fistula during infusion chemotherapy. *Hepatogastroenterology* 1996; 43: 1387–1389
- 3 Venturini M, Angeli E, Salvioni M et al. Complications after percutaneous transaxillary implantation of a catheter for intraarterial chemotherapy of liver tumors: Clinical relevance and management in 204 patients. *Am J Roentgenol* 2004; 182: 1417–1426
- 4 Takao H, Doi I, Makita K et al. Coil embolization of an arteriobiliary fistula caused by hepatic intra-arterial chemotherapy. *Cardiovasc Intervent Radiol* 2005; 28: 825–827
- 5 Tan KC, Kapoor BS. Hepatic arteriobiliary fistula successfully treated with an endobiliary covered stent. *J Vasc Interv Radiol* 2008; 19: 1521–1522

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377939>
 Endoscopy 2014; 46: E566–E567
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Hiroshi Kawakami, MD, PhD

Department of Gastroenterology and Hepatology
 Hokkaido University Graduate School of Medicine
 Kita 15, Nishi 7, Kita-ku
 Sapporo 060-8638
 Japan
 Fax: +81-11-7067867
 hiropon@med.hokudai.ac.jp