SYNLETT Spotlight

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Applications of Allenylsilanes in Organic Synthesis

Compiled by Sharada Prasanna Swain

Sharada Prasanna Swain was born in Bhirang, Orissa, India. In 2004, he received his M. Pharm from Manipal University, India. He then joined Torrent Pharmaceutical Ltd., India and attended in 2007 the doctoral program at the National Tsing Hua University, Hsinchu, Taiwan under the supervision of Prof. Dr. Jih Ru Hwu. Currently, he is working as a postdoctoral fellow at the National Central University, Jhongli, Taiwan. His current research involves the development of new synthetic methods and synthesis of novel anticancer agents.

Department of Chemistry, National Tsing Hua University, Hsinchu 30013, Taiwan
E-mail: d9623889@oz.nthu.edu.tw

Introduction

Allenylsilanes **2** are versatile reagents widely used in organic synthesis.¹ Generally, allenylsilanes react as propargyl anion equivalents in Lewis acid mediated² or thermal³ nucleophilic addition to electrophiles such as carbonyls, imines, Selectfluor, and *N*-bromosuccinimide, etc.^{1,4,5} The regioselective addition of allenylsilanes provides a β -vinyl cation, which is stabilized by a C–Si bond, which is called β effect. The C–Si bond in allenylsilanes is oriented cis-coplanar to the p-orbital of the carbocation and provides direct stabilization.⁵ Allenylsilanes undergo [3+2] annulations with α,β -unsaturated carbonyls, carbonyl compounds, imines, and nitrosyl cations to form cyclo-

pentenones, dihydrofurans, dihydropyrroles, and isoxazoles, respectively.^{1,4,5}

The efficient methods for the synthesis of allenylsilanes **2** are copper-mediated 1,3-substitution reactions of carbon nucleophiles to propargylic substrates **1** having a leaving group at the propargylic position.^{6,7}

Scheme 1 Synthesis of allenylsilanes

Abstracts

(A) *Reactions with Aldehydes:* Allenylsilanes react with aldehydes and ketones in the presence of titanium tetrachloride to provide homopropargylic alcohols in a regioselective manner. The reaction of chiral allenylsilanes with chiral aldehydes leads to the formation of mainly *syn* homopropargylic alcohols.³ The reaction of chiral 3,3-disubstituted allenylsilane **3** with paraformaldehyde in the presence of TiCl₄·2THF generates chiral homopropargylic alcohol **4**, which is the key intermediate in the total synthesis of (–)-histrionicotxin **5**.⁸

(B) *Reactions with Aldehydes:* The reaction of chiral 2-silyl-substituted α -allenyl alcohol **7** with aldehyde **6** in the presence of InBr₃ give rise to chiral homopropargylic alcohol **11**. The reaction proceeds via formation of oxocarbenium ion **8**, which undergoes a [3,3]-sigmatropic rearrangement to form the alcohol **11**. The alcohol **11** is the key intermediate in the total synthesis of the natural product (+)-neopeltolide **12**.⁹

(C) *Reactions with Aldehydes*: The addition of γ -trimethylsilyl allene esters **13** to aldehydes **14** in the presence of *i*-PrOLi leads to the formation of regiospecific γ -carbinols **18**. The addition of anionic catalyst *i*-PrOLi leads to the intermediate **15**, which possess enolate-like reactivity. The nucleophilic addition of intermediate **15** to aldehyde **14** generates intermediate **16**. Then, the silyl group undergoes a 1,3-shift and the nucleophile eliminates to form the intermediate **17**. This reaction is the key step in the total synthesis of the [3.2.1] bicyclic natural product vitisinol D.¹⁰

(D) *Reactions with Imines*: The enantioenriched allenylsilane **20** reacts with the in situ generated iminium ion generated from *t*-butyl carbamate **22** and aldehydes **21** in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ to form substituted 4,5-dihydropyrroles **24**. Similarly, the reaction of allenylsilane **20** with an iminium ion, generated in situ from methyl carbamate **23** and aldehydes **21** in the presence of TMSOTf, forms substituted 4,5-dihydrooxazines **25**.¹¹

(E) *Gold-Catalyzed Cycloisomerization*: In the presence of AuCl_3 , the γ -silyl-substituted allenyl ketones **29** undergoes cycloisomerization to 3-silyl furans **32**. The cyclization of the allenyl ketone **29** give rise to the intermediate gold-carbene **31**, and upon the 1,2-Si shift, the 3-silyl furan **32** is produced.¹²

(F) *The Pauson–Khand Reaction of 1,1-Disubstituted Allenylsilanes*: The Pauson–Khand reaction of 1,1-disubstituted allenylsilanes with terminal alkynes leads to 4-alkylidene-2-cyclopenten-1-ones in good yields. The reaction proceeds through a [2+2+1] pathway. A three-membered iron metacycle is generated by reaction of allenylsilane **33** with diiron nonacarbonyl. The iron metacycle undergoes complexation with alkyne **34**, and finally, a reductive elimination takes place to provide the 4-alkylidene-2-cyclopenten-1-one **35**.¹³

(G) *[2+2+2] Cycloaddition with Benzyne*: Benzyne possess a strained triple bond and are highly electrophilic. Allenylsilanes **37** react with two equivalents of benzyne **36** to generate (α -phenanthrenyl)vinylsilanes **38** in excellent yields. The reaction proceeds through a [2+2+2] pathway.¹⁴

References

- (1) (a) Marshall, J. A. *J. Org. Chem.* **2007**, *72*, 8153. (b) Bower, J. F.; Kim, I. S.; Patman, R. L.; Krische, M. J. *Angew. Chem. Int. Ed.* **2009**, *48*, 34. (c) Yu, S.; Ma, S. *Angew. Chem. Int. Ed.* **2012**, *51*, 3074.
- (2) (a) Felzmann, W.; Castagnolo, D.; Rosenbeiger, D.; Mulzer, J. *J. Org. Chem.* **2007**, *72*, 2182. (b) Brawn, R. A.; Panek, J. S. *Org. Lett.* **2009**, *11*, 4362.
- (3) Sabbasani, V. R.; Lee, D. *Org. Lett.* **2013**, *15*, 3954.
- (4) Fuchs, P. L. *Handbook of Reagents for Organic Synthesis: Reagents for Silicon-Mediated Organic Synthesis*; John Wiley & Sons: Chichester, **2011**.
- (5) Curtis-Long, M. J.; Aye, Y. *Chem. Eur. J.* **2009**, *15*, 5402.
- (6) Li, H.; Müller, D.; Guéneé, L.; Alexakis, A. *Org. Lett.* **2012**, *14*, 5880.
- (7) (a) Brummond, K. M.; DeForrest, J. E. *Synthesis* **2007**, 795. (b) Ding, C.-H.; Hou, X.-L. *Chem. Rev.* **2011**, *111*, 1914. (c) Yu, S.; Ma, S. *Chem. Commun.* **2011**, *47*, 5384.
- (8) Adachi, Y.; Kamei, N.; Yokoshima, S.; Fukuyama, T. *Org. Lett.* **2011**, *13*, 4446.
- (9) Guinchard, X.; Roulland, E. *Org. Lett.* **2009**, *11*, 4700.
- (10) Maity, P.; Lepore, S. D. *J. Am. Chem. Soc.* **2009**, *131*, 4196.
- (11) Brawn, R. A.; Panek, J. S. *Org. Lett.* **2009**, *11*, 473.
- (12) Dudnik, A. S.; Xia, Y.; Li, Y.; Gevorgyan, V. *J. Am. Chem. Soc.* **2010**, *132*, 7645.
- (13) Williams, D. R.; Shah, A. A.; Mazumder, S.; Baik, M.-H. *Chem. Sci.* **2013**, *4*, 238.
- (14) Hwu, J. R.; Swain, S. P. *Chem. Eur. J.* **2013**, *19*, 6556.