Z. JIN, S. CHEN, Y. WANG, P. ZHENG, S. YANG,* Y. R. CHI* (NANYANG TECHNOLOGICAL UNIVERSITY, SINGAPORE, SINGAPORE AND GUIZHOU UNIVERSITY, GUIYANG, P. R. OF CHINA)

β -Functionalization of Carboxylic Anhydrides with β -Alkyl Substituents through Carbene Organocatalysis
Angew. Chem. Int. Ed. **2014**, 53, 13506–13509.

β -Functionalization of Carboxylic Anhydrides with N-Heterocyclic Carbenes

Significance: Chi, Yang, and co-workers report the asymmetric β -functionalization of symmetrical aliphatic anhydrides **1**. Nucleophilic attack of the N-heterocyclic carbene catalyst to the anhydride generates the NHC-bound ester intermediate, which upon deprotonation forms nucleophile **2**. This adds to various electrophiles, such as alkylidene diketones, chalcones, and isatins, in a highly selective manner. Decarboxylation of the β -lactone intermediates **3** yields the final products. For almost all substrates tested, consistently very high enantioselectivities accompanied with good diastereoselectivities were achieved.

Comment: In continuation of the work by the Chi group on the activation of esters with NHC catalysts for the functionalization of the β -position (*Nature Chem.* **2013**, 5, 835), overcoming the limitation of β -aryl substrates is the main objective of the current work. Under the previous reaction conditions, only low yields (8–40%) were obtained. The presented solution for these challenging substrates utilizes anhydride substrates instead, affording the desired products in moderate to very good yields under the optimized conditions.

SYNFACTS Contributors: Benjamin List, Tim Katzenmeier
 Synfacts 2015, 11(1), 0096 Published online: 15.12.2014
 DOI: 10.1055/s-0034-1379626; Reg-No.: B13014SF

2015 © THIEME STUTTGART • NEW YORK