H. WANG, F. R. FRONCZEK, M. G. H. VINCENTE, K. M. SMITH* (LOUISIANA STATE UNIVERSITY, BATON ROUGE, USA)
Functionalization of 3,5,8-Trichlorinated BODIPY Dyes
J. Org. Chem. **2014**, *79*, 10342–10352.

Phosgene as Key Reagent for Trifunctionalization of BODIPY Dyes

Significance: The functionalization of BODIPY fluorescent dyes allows the tuning of their photo-physical, hydrophobic, and charge-transfer properties. Smith and co-workers utilize phosgene to access a 3,5,8-trichlorinated BODIPY. Exploiting the higher reactivity of the vinylic chloride in Stille couplings, regioselectively functionalized BODIPY dyes are synthesized.

Comment: Starting from a trichlorinated BODIPY and performing mixed Stille couplings with only four different organotin reagents, 14 new BODIPY dyes are reported in this paper. Photophysical properties are tunable via substitution at the 3-, 5-, and 8-positions. A plethora of regioselectively functionalized BODIPY dyes should be accessible through this strategy.

SYNFACTS Contributors: Timothy M. Swager, Sibo Lin
Synfacts 2015, 11(1), 0034 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379663; Reg-No.: S12214SF

2015 © THIEME STUTTGART • NEW YORK