H. W. PAULS* ET AL. (UNIVERSITY HEALTH NETWORK, TORONTO, CANADA AND CELTIC CATALYSTS, DUBLIN, IRELAND)

The Discovery of Polo-Like Kinase 4 Inhibitors: Identification of (1*R*,2*S*)-2-(3-((*E*)-4-(((*cis*)-2,6-Dimethylmorpholino)methyl)styryl)-1*H*-indazol-6-yl)-5'-methoxyspiro[cyclopropane-1,3'-indolin]-2'-one (CFI-400945) as a Potent, Orally Active Antitumor Agent

J. Med. Chem. **2015**, *58*, 147–169.

Synthesis of PLK4 Inhibitor CFI-400945

Significance: CFI-400945 is an inhibitor of Polo-like kinase 4 (PLK4) that is a lead for the treatment of various cancers. The synthesis depicted features a diastereoselective one-pot double $\text{S}_{\text{N}}2$ displacement reaction (**E** $\rightarrow$ **H**) for the creation of the cyclopropane ring. The authors propose that the stereoselectivity of the cyclopropanation is a consequence of π – π interactions that stabilize conformer **G**.

SYNFACTS Contributors: Philip Kocienski
 Synfacts 2015, 11(4), 0345 Published online: 18.03.2015
 DOI: 10.1055/s-0034-1380330; Reg-No.: K00915SF

Comment: Attempted hydrogenolysis of the benzyl protecting groups from a close relative of **H** was accompanied by partial ring opening of the cyclopropane. However, the benzyl groups were removed cleanly using potassium *tert*-butoxide in an oxygen-saturated solution in THF and DMSO (R. M. Williams, E. Kwast *Tetrahedron Lett.* **1989**, 30, 451).

Category

Synthesis of Natural Products and Potential Drugs

Key words

CFI-400946

Polo-like kinase 4 inhibitors

cyclopropanation

Sharpless asymmetric dihydroxylation

Suzuki–Miyaura coupling

Synfact
of the month