Over-the-scope clip (OTSC) closure of a gastro-bronchial fistula after esophagectomy

Fig. 1 Gastrobronchial fistula in a 60-year-old man. **a** Bronchoscopy showing the bronchial fistula opening (arrow). **b** Esophagogastroduodenoscopy shows the fistula opening in the fundus hemiplication of the gastric conduit.

Fig. 2 Over-the-scope clip (OTSC) closure of the gastrobronchial fistula. **a** Endoscopic view through the 11/6t OTSC just before fistula closure. **b** Simultaneous bronchoscopy: No narrowing of the bronchial lumen was seen during clip application. The diaphanoscopy of the gastroscope is clearly visible.

Fig. 3 Follow-up of the gastrobronchial fistula after over-the-scope clip (OTSC) closure. The clip is still in place surrounded by some granulation tissue 7 months after OTSC closure.

A 60-year-old man with Barrett carcinoma underwent thoracoabdominal esophagectomy with reconstruction by gastric pull-up and tubularization of the gastric conduit. The ventral part of the esophagogastrostomy was covered with the remnant gastric fundus as an anterior hemiplication. One year later, the patient developed a persistent cough, especially in a recumbent position. Assessments by endoscopy (gastroscopy, bronchoscopy) and computed tomography (CT) scan did not reveal any evidence of an intestino-bronchial fistula. Finally, the patient was treated for psychogenic coughing. During an operation for an incisional hernia 2 years after the esophageal resection, the anesthesiologist presumed the exist-

ence of an intestino-bronchial fistula because of respiratory deterioration with simultaneous ascension of gastric juice through the tracheal tube.

We therefore performed simultaneous gastroscopy and bronchoscopy immediately after the operation in the ICU. The initial gastroscopy did not reveal any signs of fistula but bronchoscopy clearly showed a fistula opening in the right main bronchus approximately 2 cm distal to the tracheal bifurcation (● **Fig. 1 a**). Detailed gastroscopic examination finally showed the fistula in a position that was very difficult to visualize, originating from the hemiplication formed by the gastric fundus (● **Fig. 1 b**). The thickness of the fistula wall was not more than 1–2 mm.

After intense discussion, we decided to attempt endoscopic fistula closure with an over-the-scope clip (OTSC), although there were no positive literature reports of such an approach with a gastrobronchial fistula.

The procedure was performed under general anesthesia. An 11/6t OTSC (Ovesco Endoscopy, Tübingen, Germany) was chosen. No grasper or anchor could be used owing to the narrow conditions in the small gastric plication. The fistula opening was only aspirated in the OTSC cap (● **Fig. 2 a**). Simultaneous bronchoscopy confirmed no narrowing of the bronchial lumen during clip application (● **Fig. 2 b**).

After the procedure, the patient reported being symptom free. Endoscopic control 3 weeks later showed the OTSC in the gastric plication surrounded by granulation tissue. On bronchoscopy, the fistula had completely healed. The clip was still in situ 7 months later and the patient free of symptoms (● **Fig. 3**).

To our knowledge, this is the first report of successful gastrobronchial fistula closure with an OTSC. There are only two cases of successful OTSC fistula closures (one esophagobronchial and one esophago-tracheal) [1,2] and one describing a combined approach (OTSC and self-expandable covered metal stent) [3]. In our case, the clip may have grasped not only the thin fistula wall but also well-perfused tissue of the gastric plication. This might have promoted healing of the fistula.

In conclusion, the use of an OTSC is justified when attempting gastrobronchial fistula closure.

Endoscopy_UCTN_Code_TTT_1AO_2AI

Competing interests: None

**Andreas Fischer¹, Jens Höppner²,
Stefan Utzolino², Hans-Jürgen
Richter-Schrag¹**

¹ University Hospital Freiburg, Germany,
Interdisciplinary Endoscopy, Department
of General and Visceral Surgery, Depart-
ment of Medecin II, Freiburg, Germany

² University Hospital Freiburg, Germany,
Department of General and Visceral
Surgery, Freiburg, Germany

References

- 1 Zolotarevsky E, Kwon Y, Bains M et al. Esophagobronchial fistula closure using a novel endoscopic over-the-scope-clip. *Ann Thorac Surg* 2012; 3: E69–70
- 2 Traina M, Curcio G, Tarantino I et al. New endoscopic over-the-scope clip system for closure of a chronic tracheoesophageal fistula. *Endoscopy* 2010; 42 (Suppl. 02): E54–55
- 3 Rebelo A, Moutinho-Ribeiro P, Cotter J. Complex endoscopic resolution of a large bronchoesophageal fistula. *Gastrointest Endosc* 2011; 4: 833–834

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1390729>
Endoscopy 2014; 46: E638–E639
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Andreas Fischer

University Hospital Freiburg, Germany
Interdisciplinary Endoscopy
Department of General and Visceral Surgery
Department of Medecin II
Sir Hans A. Krebs Str.
79106 Freiburg
Fax: +49-761-27025411
andreas.fischer@uniklinik-freiburg.de