Endoscopic treatment of a completely obstructed ileorectal anastomosis

Fig. 1 Endoscopic treatment of a completely obstructed ileorectal anastomosis. A guidewire is passed through a 19-gauge needle.

Fig. 2 Endoscopic view of the second endoscope after balloon dilation.

Fig. 3 Endoscopic aspect of the anastomosis after stent removal.

Complete anastomotic obstruction secondary to a benign disease is rare, with only a few reports of endoscopic management in the literature [1–3]. We describe a case of the successful endoscopic management of a completely obstructed ileorectal anastomosis with the use of an endoscopic rendezvous technique. A 55-year-old man was admitted in September of 2009 for severe alcoholic pancreatitis. Organ necrosis developed, so that subtotal colectomy and lateral ileostomy were required. In May 2010, an ileorectal anastomosis was established, and ileostomy closure was postponed. After 3 months, the patient underwent a rectoscopy, which revealed complete obstruction of the anastomosis. At the same time, a second operator conducted an endoscopic procedure through the ileostomy. Transillumination was obtained from both sides. Fluoroscopy showed that the two endoscope tips were almost in contact with each other. We punctured the upper side of the anastomosis with a 19-gauge needle (Cook Endoscopy, Winston-Salem, North Carolina, USA) and passed a Metro II guidewire (Cook Endoscopy) through the needle (► **Fig. 1**). A 10-Fr cystotome (Cook Endoscopy) was then passed from below upward over the guidewire, and dilation with a 10-mm Quantum Maverick balloon (Boston Scientific, Natick, Massachusetts, USA) was carried out (► **Fig. 2**), followed by the placement of a fully covered self-expandable Wallstent (Boston Scientific) measuring 80×10 cm (► **Video 1**). After 1 week, the stent was removed (► **Fig. 3**), and the new orifice was further dilated to 15 mm. At 6 months after the procedure, the patient's bowel habits had returned to normal. Complete obstruction of an anastomosis presents a particular challenge because it requires a real recanalization. The technique of rendezvous was first proposed in the management of complex esophageal strictures [4]. The use of this therapeutic approach to treat complete obstruction of a colorectal anastomosis under fluoroscopic control [5], endoscopic ultrasound [6], or computed tomography [7] has been reported. Our case is, to our knowledge, the first one in which an ileorectal anastomosis was treated.

Video 1

Video showing transillumination of the obstructed anastomosis, puncture with a 19-gauge needle, passage of the guidewire, passage of the cystotome, balloon dilation, and stent deployment. The final result is a functional anastomosis.

Endoscopy_UCTN_Code_TTT_1AQ_2AF

Competing interests: None

Najib Al Ghossaini¹, Vincent Maunoury², Richard Delcenserie³, Philippe Bulois⁴

¹ Saint Omer Hospital, Gastroenterology Department, Saint Omer, France

² Lille University Hospital, Gastroenterology Department, Lille, France

³ Amiens University Hospital, Gastroenterology Department, Amiens, France

⁴ La Louvière Private Hospital, Générale de Santé, Lille, France

References

- 1 Mukai M, Kishima K, Iizuka S et al. Endoscopic hook knife cutting before balloon dilatation of a severe anastomotic stricture after rectal cancer resection. *Endoscopy* 2009; 41: E193–E194
- 2 Repici A, Pagano N, Romeo F et al. Endoscopic treatment of a complete colonic anastomotic stricture by the rendez-vous technique. *Dig Liv Dis* 2010; 42 (Suppl. 02): 98–S99
- 3 Curcio G, Spada M, di Francesco F et al. Completely obstructed colorectal anastomosis: a new nonelectrosurgical endoscopic approach before balloon dilatation. *World J Gastroenterol* 2010; 16: 4751–4754
- 4 Bueno R, Swanson SJ, Jaklitsch MT et al. Combined antegrade and retrograde dilation: a new endoscopic technique in the management of complex esophageal stricture. *Gastrointest Endosc* 2001; 54: 368–372
- 5 Kaushik N, Rubin J, McGrath K. Treatment of benign complete colonic anastomotic obstruction by using an endoscopic rendez-vous technique. *Gastrointest Endosc* 2006; 4: 727–730
- 6 De Lusong MA, Shah JN, Soetikno R et al. Treatment of a completely obstructed co-

lonic anastomotic stricture by using a prototype forward-array echoendoscope and facilitated by SpyGlass (with videos). *Gastrointest Endosc* 2008; 68: 988–992

- 7 Albertsmeier M, Rittler P, Hoffmann RT et al. Treatment of a completely obstructed colonic anastomosis using a CT-guided endoscopic rendezvous technique. *Endoscopy* 2011; 43 (Suppl. 02): E5–E6

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1390753>
Endoscopy 2014; 46: E609–E610
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Philippe Bulois, MD
 20 rue du Ballon
 F-59000 Lille
 France
 Fax: +33320576947
philippebulois@yahoo.fr