Primary malignant melanoma involving the whole esophagus: a rare case with rarer presentation

Fig. 1 Endoscopy image showing growth with irregular projections, black hue and luminal narrowing in a 50-year-old man with a 2-month history of dysphagia and weight loss.

Fig. 2 Endoscopy image showing subepithelial growth with blue-black hue.

Fig. 3 Endoscopic ultrasound image showing a solid lesion with intraluminal and extraluminal growth.

Fig. 5 Biopsies (H&E stain, magnification: ×40) image showing malignant cells arranged in sheets.

Fig. 4 Computed tomography (CT) image showing the extent of the tumor along the whole length of the esophagus with loss of planes to the posterior trachea and aorta.

A 50-year-old man presented with a 2-month history of dysphagia and weight loss. A gastroscopy was done and showed an irregular submucosal bulge with a blue-black hue along the whole length of the esophagus (● Fig. 1 and ● Fig. 2). An

endoscopic ultrasound was performed to look for vascularity, and showed a solid lesion with intraluminal and extraluminal growth limited to the serosa (● Fig. 3). A computed tomography (CT) chest scan was done and this showed a solid lesion

involving the whole length of the esophagus, loss of planes to the posterior trachea and aorta, and multiple bony lesions (● Fig. 4). Endoscopic biopsies showed malignant cells arranged in sheets with moderate cytoplasm and moderately pleomorphic nuclei (● Fig. 5), and a coarse brown granular pigment with Masson-Fontana stain. Immunohistochemistry results were as follows: CK negative, HMB-45, S-100, and Vimentin positive, suggestive of malignant melanoma. Primary malignant melanoma of the esophagus arises from melanin cells of the mucosal epithelial basal layer. It is a rare disease with extremely poor prognosis owing to its high metastatic potential. It represents 0.1% to 0.2% of all esophageal malignant tumors and generally presents with dysphagia (80% cases), retrosternal or epigastric discomfort or pain [1,2]. The lower third of the esophagus is the most common site followed by the middle and upper esophagus [1]. Our patient had involvement of the whole esophagus which makes it a rarer presentation. Diagnosis can be suspected during endoscopy if the mass has a black or dark brown pigment [2]. Positive immunohistochemistry for S-100, and HMB-45 helps in diagnosis [2]. Surgery is the mainstay of treatment when possible [1,3].

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AB

Competing interests: None

**Narendra Singh Choudhary¹,
Rajesh Puri¹, Ruchika Goel²,
Randhir Sud¹**

¹ Institute of Digestive and Hepatobiliary
Sciences, Medanta, The Medicity,
Gurgaon, India

² Department of Histopathology, Medanta,
The Medicity, Gurgaon, India

References

- 1 *Sabanathan S, Eng J, Pradhan GN.* Primary malignant melanoma of the esophagus. *Am J Gastroenterol* 1989; 84: 1475–1481
- 2 *Li YH, Li X, Zou XP.* Primary malignant melanoma of the esophagus: a case report. *World J Gastroenterol* 2014; 20: 2731–2734
- 3 *Li B, Lei W, Shao K et al.* Characteristics and prognosis of primary malignant melanoma of the esophagus. *Melanoma Res* 2007; 17: 239–242

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1390783>
Endoscopy 2014; 46: E621–E622
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Rajesh Puri

Institute of Digestive and Hepatobiliary Sciences
Medanta
The Medicity
Sector 38
Gurgaon
Haryana 122001
India
Fax: +91-124-4834111
purirajesh69@gmail.com