Gastric tuberculosis resembling depressed type, early gastric cancer

Fig. 1 Esophagogastroduodenoscopy (EGD) showing a depressed lesion with converging folds on the posterior wall of the upper gastric corpus in a 60-year-old asymptomatic man.

Fig. 2 EGD with chromoendoscopy revealed an irregularly shaped, depressed lesion of which the anal portion appeared to resemble a whitish nodule (arrow).

Fig. 3 Image enhanced endoscopy with magnifying narrow-band imaging demonstrating the spiral-shaped, dilated small vessels or microvessels with an amorphous surface.

Fig. 4 Endoscopic ultrasonography (EUS) demonstrating a hypoechoic lesion localized in the deep portion of the mucosa and the superficial submucosa.

A 60-year-old asymptomatic man was referred to our hospital for evaluation of a gastric lesion detected by esophagogastroduodenoscopy (EGD) in a medical check-up. EGD showed an irregularly shaped, depressed lesion with converging folds on the posterior wall of the upper gastric corpus (● Fig. 1, ● Fig. 2). Magnifying endoscopy with narrow-band imaging at the anal portion of the lesion revealed spiral-shaped, dilated small vessels/microvessels with an amorphous surface (● Fig. 3). Endoscopic ultrasonography (EUS) demonstrated a hypoechoic lesion localized in the deep portion of the mucosa and the superficial submucosa (● Fig. 4). A biopsy from the lesion showed granulomatous inflammation with caseous necrosis and Langerhans giant cells. Acid-fast bacilli were detected by both Ziehl-Neelsen staining (● Fig. 5) and mycobacterium culture. A polymerase chain reaction test for tuberculosis was also positive. Fluorine-18 fluorodeoxyglucose positron emission tomography (PET) showed markedly increased accumulation in the lymph nodes of the mediastinum, pulmonary hilum, and upper abdomen (● Fig. 6). Chest computed tomography (CT) revealed no evidence of pulmonary tuberculosis. Colonoscopy, small-bowel capsule endoscopy, and bronchoscopy showed normal findings. Biopsy from the inguinal lymph node demonstrated non-specific inflammation without any neoplastic cells or granulomas. Thus, the patient was diagnosed as having gastric tuberculosis with systemic lymphadenopathy, and subsequently underwent anti-tuberculous treatment. Both the gastric lesion and lymphadenopathy had resolved 6 months later.

Gastric tuberculosis is rare, and its endoscopic appearance can vary [1–4]. To confirm a definitive diagnosis of tuberculosis, EUS-guided, fine-needle aspiration or surgery is sometimes required [1,4,5]. The gastric lesion in our case resembled depressed-type, early gastric cancer endoscopically; however when the amorphous area (probably composed of inflammation or granulation tissue covered with thin epithelium) was viewed using magnifying endoscopy with narrow-band imaging, it was clearly different from that of gastric cancer. We thus consider that magnifying endoscopy with narrow-band imaging is useful in the differential diagnosis between gastric tuberculosis and gastric cancer.

Fig. 5 Histologic image of a biopsy specimen from the gastric lesion showing acid-fast bacilli (Ziehl-Neelsen stain, magnification: × 1000).

Fig. 6 Fluorine-18 fluorodeoxyglucose positron emission tomography (PET) showing markedly increased accumulation in the lymph nodes of the mediastinum, pulmonary hilum, and upper abdomen.

Endoscopy_UCTN_Code_CCL_1AB_2AD_3AF

Competing interests: None

Hiroki Yaita¹, Shotaro Nakamura², Koichi Kurahara¹, Toshifumi Morishita¹, Shuji Kochi¹, Yumi Oshiro³, Naohiko Hamaguchi⁴, Tadahiko Fuchigami¹

¹ Division of Gastroenterology, Matsuyama Red Cross Hospital, Ehime, Japan

² Department of R/D for Surgical Support System, Center for Advanced Medical Innovation, Kyushu University, Fukuoka, Japan

³ Department of Pathology, Matsuyama Red Cross Hospital, Ehime, Japan

⁴ Department of Respiratory Medicine, Matsuyama Red Cross Hospital, Ehime, Japan

References

- 1 Liu PF, Chang CS, Wang J et al. Primary gastric tuberculosis. *Endoscopy* 2009; 41: E327–E328
- 2 Lim JU, Kim YH, Choi CW et al. Gastric tuberculosis presenting with a huge abdominal mass. *Singapore Med J* 2013; 54: e244–e246
- 3 Kim DY, Bang S, Park BK et al. Tuberculous mesenteric lymphadenitis involving the gastric wall: case report. *Gastrointest Endosc* 2005; 62: 799–802
- 4 Kim SH, Park JH, Kang KH et al. Gastric tuberculosis presenting as a submucosal tumor. *Gastrointest Endosc* 2005; 61: 319–322
- 5 Rana SS, Bhasin DK, Srinivasan R et al. Gastric outlet obstruction caused by tuberculosis and diagnosed by endoscopic ultrasound-guided fine needle aspiration. *Endoscopy* 2011; 43: E117–E118

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1390865>
Endoscopy 2014; 46: E669–E670
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Hiroki Yaita, MD

Division of Gastroenterology
 Matsuyama Red Cross Hospital
 1 Bunkyo-cho
 Matsuyama-shi
 Ehime 790-8524
 Japan
 Fax: +81-89-9269916
hyaita@matsuyama.jrc.or.jp