Argon plasma coagulation of a bleeding angioectasia in a jejunal diverticulum by single-balloon enteroscopy

Fig. 1 A large diverticulum in the proximal jejunum with a bulky clot.

Fig. 2 Bleeding angioectasia located in the fundus of the diverticulum.

An 86-year-old woman with a history of cardiac and chronic renal failure was referred to our gastroenterology outpatient clinic because of an iron deficiency anemia and intermittent, self-limited hematochezia episodes. Colonoscopy and upper endoscopy revealed no alterations. A capsule endoscopy showed a nonbleeding angioectasia in the proximal ileum. Single-balloon enteroscopy was proposed to treat the angioectasia. During the procedure, a large jejunal diverticulum with a bulky clot was seen in the proximal jejunum (Fig. 1). The clot was removed after water irrigation, and an oozing bleeding was found from an angioectasia located in the fundus of the diverticulum (Fig. 2). Distal tattooing was performed. Bleeding was then controlled with combination therapy: first a submucosal epi-

nephrene injection (Fig. 3) followed by argon plasma coagulation (APC) of the angioectasia (Fig. 4). No immediate or delayed complications were recorded, and there was no recurrence of bleeding during 1 year of follow-up.

Small-bowel diverticula are a rare condition usually characterized by mucosal and submucosal herniation through the muscular layer on the mesenteric border [1]. Although they are usually asymptomatic, malabsorption, abdominal discomfort, obstruction, bloating, bleeding, or perforation may occur [2]. Currently, capsule enteroscopy and device-assisted enteroscopy are the mainstays for diagnosis [3]. Small-bowel angioectasias are a common cause of obscure gastrointestinal bleeding, mostly in the elderly. APC by device-assisted enteroscopy has been shown

to be effective and safe in the management of small-bowel angioectasias [4]. Identification of a bleeding angioectasia in a jejunal diverticulum is an exceptional finding. The risk of diverticular perforation after APC may be increased. To the best of our knowledge, this is the first case report of APC by single-balloon enteroscopy in a bleeding angioectasia located in a jejunal diverticulum.

Endoscopy_UCTN_Code_CCL_1AC_2AF

Competing interests: None

Carlos Fernandes¹, Rolando Pinho¹, Adélia Rodrigues¹, Germano Vilas-Boas², Iolanda Ribeiro¹, Joana Silva¹, Ana Ponte¹, João Carvalho¹

¹ Department of Gastroenterology and Hepatology, Centro Hospitalar Vila Nova Gaia/Espinho, Vila Nova de Gaia, Portugal

² Department of Gastroenterology, Hospital Pedro Hispano, Matosinhos, Portugal

References

- 1 Williams RA, Davidson DD, Serota AI et al. Surgical problems of diverticula of the small intestine. *Surg Gynecol Obstet* 1981; 152: 621–626
- 2 Johnson KN, Fankhauser GT, Chapital AB et al. Emergency management of complicated jejunal diverticulosis. *Am Surg* 2014; 80: 600–603
- 3 Ell C, May A, Nachbar L et al. Push-and-pull enteroscopy in the small bowel using the double-balloon technique: results of a prospective European multicenter study. *Endoscopy* 2005; 37: 613–616
- 4 Samaha E, Rahmi G, Landi B et al. Long-term outcome of patients treated with double balloon enteroscopy for small bowel vascular lesions. *Am J Gastroenterol* 2012; 107: 240–246

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1390918>
Endoscopy 2015; 47: E62
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Carlos Fernandes
Centro Hospitalar Vila Nova Gaia
Gastroenterology and Hepatology
Rua Conceição Fernandes
Vila Nova de Gaia 4434
Portugal
Fax: +351-22-7830209
carlosdpfernandes@gmail.com

Fig. 3 Appearance after submucosal epinephrine injection to control bleeding.

Fig. 4 Appearance after argon plasma coagulation of the angioectasia.