Endoscopic ultrasonography-guided liver abscess drainage using a dedicated, wide, fully covered self-expandable metallic stent with flared-ends



Fig. 1 An abscess in the left lobe of the liver identified at computed tomography (CT) in an 84-year-old man with a 12-day history of high fever and epigastric pain.



Endoscopic ultrasonography (EUS)-guided drainage of liver abscesses has recently become available. We present here a case of successful drainage of a liver abscess using a dedicated wide fully covered selfexpandable metallic stent (FCSEMS) with flared ends.

An 84-year-old man was admitted to another hospital with a history of high fever and epigastric pain for 12 days. Computed tomography (CT) revealed a  $10.3 \times$ 6.1-cm abscess in the left lobe of the liver (**•** Fig. 1). He was referred to our hospital because of failure of a 1-week antibiotic regimen. Esophagogastroduodenoscopy revealed a bulging mass in the stomach (**•** Fig.2).

We attempted EUS-guided drainage through a transgastric approach. Using a 19-gauge needle, we punctured the abscess and placed a 0.025-inch guidewire (**•** Fig.3). A 6-Fr wire-guided diathermic dilator (Cysto-Gastro-Set; Endo-Flex, Voerde, Germany) was used to dilate the needle tract using a blended cut mode. Finally, a dedicated wide FCSEMS with flared Fig.2 Esophaqoqastroduodenoscopy (EGD)

Fig. 2 Esophagogastroduodenoscopy (EGD) shows a bulging mass in the upper body of the stomach

ends (NAGI stent, 16×3 cm; Taewoong-Medical, Seoul, Korea) was placed, without any complications (**> Fig. 4**, **> Fig. 5**). After stone clearance from the common bile duct, the patient was discharged on postoperative day 10 without removal of the SEMS (**> Fig. 6**).

Seven cases of EUS-guided drainage of liver abscess, including one case of multiple abscesses, have been reported to date [1,2]. The left lobe of the liver, the caudate lobe, and the gastrohepatic space usually lie in close proximity to the stomach or duodenum [1]. Therefore, EUS-guided liver abscess drainage might be safe and effective in the management of these areas. Single or double plastic stents were used in most of the reported cases [1]; the newly designed dedicated anchoring FCSEMS with a "yo-yo" shape was placed in only one patient [2]. It has been suggested that a dedicated FCSEMS is the ideal stent for treating liver abscesses and pancreatic fluid collection because of its antimigration feature and because it allows direct insertion of an endoscope through it [3].

Endoscopy\_UCTN\_Code\_TTT\_1AS\_2AC

### Competing interests: None



Fig. 4 The dedicated, wide, fully covered selfexpandable metallic stent (FCSEMS) with flared ends (NAGI stent; Taewoong-Medical, Seoul, Korea)



Fig. 6 Endoscopic view through the NAGI stent on postoperative day 8, showing only necrotic tissues on the surface of the liver.



Fig. 5 Radiograph showing placement of the NAGI stent into the cavity of the liver abscess. Inset: endoscopic view of the purpose-designed stent.



## Kawakubo<sup>2</sup>, Masaki Kuwatani<sup>3</sup>, Yoshimasa Kubota<sup>1</sup>, Yoko Abe<sup>1</sup>, Shuhei Kawahata<sup>1</sup>, Kimitoshi Kubo<sup>1</sup>, Naoya Sakamoto<sup>2</sup>

- <sup>1</sup> Department of Gastroenterology and Hepatology, Hokkaido University Hospital
- <sup>2</sup> Department of Gastroenterology and Hepatology, Hokkaido University Graduate School of Medicine, Sapporo, Japan
- <sup>3</sup> Division of Endoscopy, Hokkaido University Hospital

### References

- 1 Singhal S, Changela K, Lane D et al. Endoscopic ultrasound-guided hepatic and perihepatic abscess drainage: an evolving technique. Therap Adv Gastroenterol 2014; 7: 93 - 98
- 2 Alcaide N, Vargas-Garcia AL, de la Serna-Higuera C et al. EUS-guided drainage of liver abscess by using a lumen-apposing metal stent (with video). Gastrointest Endosc 2013; 78: 941-942
- 3 Kawakami H, Itoi T, Sakamoto N. Endoscopic ultrasound-guided transluminal drainage for peripancreatic fluid collections: Where we are now? Gut Liver 2014; 8: 341-355

#### Bibliography

**DOI** http://dx.doi.org/ 10.1055/s-0034-1391125 Endoscopy 2014; 46: E982-E983 © Georg Thieme Verlag KG Stuttgart · New York ISSN 0013-726X

# **Corresponding author**

Hiroshi Kawakami, MD, PhD Department of Gastroenterology and Hepatology Hokkaido University Hospital Kita 14, Nishi 5, Kita-ku Sapporo 060-8648 Japan Fax: +81-11-7067867 hiropon@med.hokudai.ac.jp