

An alternative method of endoscopic retrieval of a proximally migrated biliary plastic stent using a “sphincterotome hooking the stent” technique

Stent migration occurs in about 5%–10% of patients undergoing biliary stenting. The risk of proximal stent migration is higher with benign biliary strictures, stenosis of the lower bile duct, bile duct diameter >10mm, duration of stent placement >1 month, and the use of straight and 10-Fr stents [1]. Devices that have been used for the endoscopic removal of stents are biopsy forceps, grasping forceps, basket catheters, balloon catheters, snares, ultra slim cholangioscopes, and the Soehendra stent retriever [2–4]. We present the case of a 38-year-old woman who was treated with sphincterotomy and insertion of a 6-cm, 10-Fr, straight, plastic, biliary stent for a post-cholecystectomy bile duct leakage. Gastroduodenoscopy 30 days later revealed that the stent was not “in situ,” and removal of the proximally migrated stent was attempted.

Initially, a guidewire was advanced through the lumen of the stent (▶ Fig. 1) and then a stone retrieval balloon was used, over the wire, to remove the stent. However, this was unsuccessful because the tip of the balloon catheter could not advance past the proximal end of the stent. A sphincterotome was then used, which was advanced successfully over the wire, to 2–3 cm beyond the proximal end of the stent, probably because of its stiffness compared with the balloon. Then, the proximal end of the sphincterotome was bent to form a “hook” on which to catch the stent (▶ Fig. 2), and the sphincterotome was pulled back slowly. When the hooked stent appeared through the papilla (▶ Fig. 3), the endoscope was pulled back in the stomach, from where the stent was finally retrieved using a grasping forceps (▶ Video 1).

We propose the use of this previously unreported, simple, and technically easy method for the endoscopic retrieval of a proximally migrated biliary stent.

Endoscopy_UCTN_Code_CPL_1AK_2AD

Competing interests: None



Fig. 1 Radiological image showing a guidewire being advanced through the lumen of the stent.



Fig. 2 Radiological image showing the sphincterotome, which was advanced through the stent and then bent in order to “hook” the stent.

Video 1

Endoscopic retrieval of a proximally migrated, plastic, biliary stent.




Fig. 3 Endoscopic image showing sphincterotome and “hooked” stent appearing through the papilla.

**Christos Liatsos¹, Nikolaos Kyriakos¹,
Grigorios Anthopoulos¹, Michalis
Galanopoulos¹, Evangelos Kalafatis¹,
Christos Mavrogiannis²**

¹ Gastroenterology Department, 401 Army General Hospital of Athens, Athens, Greece

² Academic Department of Gastroenterology, Faculty of Nursing, Kifissia General and Oncology Hospital, Athens University, Athens, Greece

References

- 1 Kawaguchi Y, Ogawa M, Kawashima Y et al. Risk factors for proximal migration of biliary tube stents. *World J Gastroenterol* 2014; 20: 1318–1324
- 2 Katsinelos P, Kountouras J, Paroutoglou G et al. Migration of plastic biliary stents and endoscopic retrieval: an experience of three referral centers. *Surg Laparosc Endosc Percutan Tech* 2009; 19: 217–221
- 3 Tarnasky PR, Cotton PB, Baillie J et al. Proximal migration of biliary stents: attempted endoscopic retrieval in forty-one patients. *Gastrointest Endosc* 1995; 42: 513–520
- 4 Arhan M, Odemiş B, Parlak E et al. Migration of biliary plastic stents: experience of a tertiary center. *Surg Endosc* 2009; 23: 769–775

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1391242>
Endoscopy 2015; 47: E64–E65
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Nikolaos Kyriakos, MD, PhD
 Gastroenterology Department
 401 Army General Hospital
 N.Vitali 38
 Cholargos
 Athens 15562
 Greece
 Fax: +30-210-7494095
 nikos_kiriakos@yahoo.gr