Aortic wall rupture from a mediastinal tumor invasion diagnosed by endoscopic ultrasound

Rupture of the aorta from tumor infiltration is a rare event, described in only a few case reports to date [1–4]. It represents a life-threatening condition with the prognosis strongly dependent on its early recognition, which allows life-saving emergency interventional therapeutic procedures to be performed [5].

We present the case of a 57-year-old man with a posterior mediastinal mass who was referred for endoscopic ultrasound-guided fine-needle aspiration (EUS-FNA). During the procedure, a large parietal break in the posterior wall of the aorta was observed, which appeared to communicate with the inner part of the

mediastinal tumor. The vascular nature of the tumor was confirmed by color Doppler examination (● Fig. 1, ● Video 1). This finding had not been present on a thoracic computed tomography (CT) scan 2 weeks before the procedure.

The EUS-FNA procedure was aborted and another chest CT scan was immediately performed, confirming the aortic pseudoaneurysm formation due to tumor invasion. Aortography with placement of two endoprostheses was immediately carried out, with success. A repeat EUS 2 days later, with FNA of the tissue surrounding the now obliterated pseudoaneurysm (● Fig. 2), was performed using

a 25-gauge needle (● Video 1), and a diagnosis of high degree spindle cell sarcoma was made.

Vascular structures represent important landmarks during EUS examinations. Thus, endosonographers can incidentally detect abnormalities of the aorta or any other big vessel, which in some cases might be mistaken for a lesion of a different nature [6–8]. In the presented case, the use of EUS made it possible to diagnose aortic pseudoaneurysm formation, which constituted a “ticking time bomb” in this patient, making it essential to abort the risky endoscopic procedure (EUS-FNA), which could have become lethal [9]. The procedure was rescheduled only

Video 1

Case study of a 57-year-old man with mediastinal tumor invasion into the aorta.

Fig. 1 Endoscopic ultrasound view of the aortic wall defect (arrows) communicating with the inner part of the tumor. **a** B-mode. **b** Color Doppler examination.

Fig. 2 Computed tomography of the chest showing pseudoaneurysm (yellow arrow) before and after endograft placement into the thoracic aorta (empty arrow). **a** Before endograft placement. **b** After endograft placement, which resulted in complete resolution of the pseudoaneurysm.

after this procedural risk had been reduced, as saving a life is undoubtedly infinitely more important than a tissue diagnosis.

Endoscopy_UCTN_Code_CCL_1AF_2AC

Competing interests: Dr. Larghi is a consultant for Cook Medical and Boston Scientific, and has received research support from Olympus.

**Mihai Rimbaş^{1,2}, Fabia Attili¹,
Alberto Larghi¹**

¹ Digestive Endoscopy Unit, Catholic University, Rome, Italy

² Gastroenterology Department, Colentina Clinical Hospital, Bucharest, Romania

References

- 1 Tsuchida R, Kasahara N, Inobe M et al. Aortic intramural hematoma associated with metastatic carcinoma. *Pathol Res Pract* 2010; 206: 839–845
- 2 Falcone JL, Go MR, Baril DT et al. Vascular wall invasion in neurofibromatosis-induced aortic rupture. *Vasc Endovascular Surg* 2010; 44: 52–55
- 3 Stambo G, Valentin M, Kerr TM et al. Endovascular treatment of an acutely ruptured abdominal aorta from tumor invasion by an unresectable retroperitoneal leiomyosarcoma. *Ann Vasc Surg* 2008; 22: 568–570
- 4 Cantwell CP, Stack J. Abdominal aortic invasion by leiomyosarcoma. *Abdom Imaging* 2006; 31: 120–122
- 5 Collaud S, Waddell TK, Yasufuku K et al. Thoracic aortic endografting facilitates the resection of tumors infiltrating the aorta. *J Thorac Cardiovasc Surg* 2014; 147: 1178–1182
- 6 Muraki S, Inguu A, Baba M et al. [A case report of a pseudoaneurysm of the thoracic aorta (thrombotic closure type) showing like a mediastinal tumor – value of endoscopic ultrasonography in differential diagnosis]. *Nihon Kyobu Geka Gakkai Zasshi* 1996; 44: 1917–1920
- 7 Varadarajulu S, Eloubeidi MA. Diagnosis of an aneurysm masquerading as a pancreatic-cyst lesion at EUS. *Gastrointest Endosc* 2007; 65: 721–725
- 8 Nagamatsu H, Takahashi K, Ueo T et al. A case of splenic artery aneurysm simulating a pancreas tumor. *Nihon Shokakibyo Gakkai Zasshi* 2011; 108: 1420–1427
- 9 Lachter J. Fatal complications of endoscopic ultrasonography: a look at 18 cases. *Endoscopy* 2007; 39: 747–750

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1391243>
Endoscopy 2015; 47: E66–E67
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Alberto Larghi, MD, PhD
Digestive Endoscopy Unit
Catholic University
Largo A. Gemelli 8
00168, Rome
Italy
Fax: +39-06-30156581
albertolarghi@yahoo.it