Chronic diarrhea because of villous atrophy unrelated to celiac disease

A 71-year-old woman was admitted with a 7-month history of watery diarrhea, which had led to an unintentional 30 kg (27%) of weight loss and admittance to the intensive care unit (ICU) on two separate occasions as a result of dehydration. Endoscopy and video capsule evaluation revealed villous atrophy of the entire small bowel, with fissuring, nodularity, and loss of folds, as shown for both duodenum (● Fig. 1 a) and ileum (● Fig. 1 b). The mucosa appeared fragile, with ulcers after biopsies (● Fig. 1 c). Histology confirmed complete villous atrophy and showed lengthened regenerative crypts, only a few intra-epithelial lymphocytes, and thickening of the basal membrane in both proximal (● Fig. 2) and distal small-bowel biopsies. Colonoscopy revealed a pale and edematous mucosa with superficial ulcerations, more pronounced distally (● Fig. 3). Microscopic evaluation showed subtle

inflammation in colon biopsies with focal erosion, a focally thickened basal membrane, and some apoptotic cells in the epithelium (● Fig. 4 a, b). Infectious, ischemic, and malignant disorders were excluded. Serum anti-tTG IgA and anti-gliadin IgG were negative during and after gluten exposure, ruling out celiac disease. The clinical presentation and diagnostic findings were most compatible with adult-onset autoimmune enteropathy [1], affecting an extensive part of the digestive tract (stomach to rectum). Immunosuppressive therapy was started; however high dose prednisolone, increasing doses of azathioprine, and immunoglobulins failed to induce any clinical response. The patient continued to produce voluminous diarrhea; however 3 weeks after starting therapy with oral budesonide (3×3 mg daily, pulverized in the morning, granules at noon, capsule at night) [1], the patient recovered, with

formed stools, clinical improvement, and weight gain. Duodenal biopsies revealed completely restored villous architecture (● Fig. 5). The patient has remained well for 20 months of follow-up.

Thus, autoimmune enteropathy should be considered after exclusion of celiac disease when severe diarrhea is associated with villous atrophy. Topical immunosuppressive treatment should be applied.

Endoscopy_UCTN_Code_CCL_1AC_2AD

Competing interests: None

Anja Ursula van Lent¹, R. Bart Takkenberg¹, Susanne van Eeden², Chris Mulder³, Ulrich Beuers¹

¹ Department of Gastroenterology and Hepatology, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

² Department of Pathology, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

³ Department of Gastroenterology and Hepatology, VU University Medical Center, University of Amsterdam, Amsterdam, The Netherlands

Fig. 1 A 71-year-old woman was admitted with a 7-month history of watery diarrhea and weight loss. Endoscopic images showing: **a** villous atrophy in the duodenum; **b** villous atrophy in the ileum; **c** ulcers after biopsies in the duodenum.

Fig. 2 Proximal small-bowel biopsy (hematoxylin and eosin (H&E) stain) of duodenal mucosa demonstrating severe villous atrophy, some inflammation, and thickening of the basal membrane.

Fig. 3 Colonoscopy of the distal colon showing pale and edematous mucosa with superficial ulcerations.

Fig. 4 Colonic mucosa (H&E staining) showing: **a** focal erosion with homogenization of the lamina propria; **b** some apoptotic cells (arrows) in the epithelium.

Fig. 5 Restored villous architecture after topical budesonide treatment: **a** endoscopic view; **b** microscopic view.

Reference

- 1 Akram S, Murray JA, Pardi DS et al. Adult autoimmune enteropathy: Mayo Clinic Rochester experience. *Clin Gastroenterol Hepatol* 2007; 5: 1282–1290

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1391247>
Endoscopy 2015; 47: E71–E72
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Anja Ursula van Lent, MD, PhD
 Department of Gastroenterology and Hepatology
 Academic Medical Center
 University of Amsterdam
 P.O. Box 22700
 1100 DE Amsterdam
 The Netherlands
 Fax: +31-20-5669608
a.u.vanlent@amc.uva.nl