Esophageal fully covered self-expanding metal stent for combined treatment of an aorto-esophageal fistula

Fig. 1 Computed tomography (CT) scan showing a thoracic aortic pseudoaneurysm causing a mass effect on the esophagus.

Fig. 2 Gastrografin esophagogram showing discontinuity in the medial aspect of the esophagus (arrow) at the level of the aortic arch.

We report on a 63-year-old man with a history of coronary artery bypass in 1995 who presented with massive hematemesis and hemodynamic instability. A computed tomography (CT) scan revealed a thoracic aortic pseudoaneurysm causing a mass effect on the esophagus (► Fig. 1) and rupture of the aorta immediately after the origin of the left subclavian artery, with the suspicion of an aorto-esophageal fistula.

Endovascular aortic treatment was undertaken with a carotid–subclavian bypass using a polytetrafluoroethylene stent. A Gastrografin esophagogram 4 days after admission revealed a discontinuity in the medial aspect of the esophagus at the level of the aortic arch (► Fig. 2). An upper gastrointestinal endoscopy performed the following day confirmed a 4-cm long esophageal rupture 20 cm from the incisors (► Fig. 3). A fully covered, 110-mm × 20-mm, Nitinol self-expanding metal stent (Hanarostent M.I.Tech Co. Inc., Seoul, South Korea) was placed under direct visualization to isolate the esophageal perforation and a nasogastric tube was subsequently placed.

The patient was initially kept on enteral feeding, later progressing to an oral diet. Twenty weeks after the placement of the esophageal stent, the patient remains alive, with no evidence of mediastinitis, aortic stent-graft infection or new episodes of gastrointestinal bleeding, with the esophageal stent being correctly placed (► Fig. 4).

Aorto-esophageal fistulas are a rare cause of massive upper gastrointestinal bleeding; one of the main causes is erosion of the esophageal wall by thoracic aortic aneurysms [1]. Endovascular aortic repair enables aortic bleeding to be rapidly controlled [2]; however, it does not repair the esophagus, so leaves the risk of mediastinitis and aortic graft contamination.

Previous reports have considered surgical repair of the esophageal lesion once the patient's condition has been stabilized [3], but secondary esophageal reconstruc-

Fig. 3 Upper gastrointestinal endoscopy showing a 4-cm long esophageal rupture 20 cm from the incisors, with no evidence of spontaneous purulent drainage.

Fig. 4 Computed tomography (CT) scan showing the correctly placed esophageal stent.

tion can cause significant surgical trauma. Esophageal stents are universally used to treat esophageal perforations [4]. In selected cases, they may be a less invasive and equally effective alternative to surgical repair, bypassing the defect to allow it to heal, while allowing oral feeding to be continued.

Endoscopy_UCTN_Code_TTT_1AO_2AI

Competing interests: None

**Eduardo Rodrigues-Pinto,
Pedro Pereira, Guilherme Macedo**

Gastroenterology Department,
Centro Hospitalar São João, Porto

References

- 1 Hollander JE, Quick G. Aortoesophageal fistula: a comprehensive review of the literature. *Am J Med* 1991; 91: 279–287
- 2 Prokakis C, Koletsis E, Apostolakis E et al. Aortoesophageal fistulas due to thoracic aorta aneurysm: surgical versus endovascular repair. Is there a role for combined aortic management? *Med Sci Monit* 2008; 14: RA48–RA54
- 3 Topel I, Stehr A, Steinbauer MG et al. Surgical strategy in aortoesophageal fistulae: endovascular stentgrafts and in situ repair of the aorta with cryopreserved homografts. *Ann Surg* 2007; 246: 853–859
- 4 Sharma P, Kozarek R. Practice Parameters Committee of American College of Gastroenterology. Role of esophageal stents in benign and malignant diseases. *Am J Gastroenterol* 2010; 105: 258–273

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1391256>
Endoscopy 2015; 47: E73–E74
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Eduardo Rodrigues-Pinto, MD
 Gastroenterology Department
 Centro Hospitalar São João
 Al. Prof. Hernâni Monteiro
 4200-319 Porto
 Portugal
 Fax: +351-22-5513601
edu.gil.pinto@gmail.com