Endoscopic ultrasound-guided pancreatic duct drainage using antegrade stenting

Fig. 1 Endoscopic ultrasound (EUS)-guided pancreatic duct antegrade stenting was performed on a 66-year-old woman. At endoscopic retrograde cholangiopancreatography (ERCP) the guidewire was successfully advanced into the pancreatic duct through the stenosis, but the ERCP cannula and other devices, including the dilation catheter, could not be advanced.

Fig. 2 The pancreatic duct has been punctured using a 19-G needle (endosonographic image).

Fig. 3 The contrast medium was injected through the 19-G needle.

Endoscopic ultrasound (EUS)-guided pancreatic duct access in patients with an inaccessible papilla or failed endoscopic retrograde cholangiopancreatography (ERCP) has been reported previously [1, 2]. This technique consists of the EUS-guided rendezvous technique, and EUS-guided pancreatic duct stenting via the

stomach (pancreaticogastrostomy), or the duodenum (pancreaticoduodenostomy) [2]. However, the technical success rate of the EUS rendezvous technique is not very high [3], and it is not indicated in patients with altered anatomy, for example following a Roux-en-Y procedure. Stent occlusion is a risk of EUS-guided pancreatico-

duodenostomy, and it may easily occur because of food impaction. Novel technical tips for EUS-guided pancreatic duct antegrade stenting are presented here (▶ **Video 1**).

A 66-year-old woman was admitted to our hospital because of abdominal pain. She had previously had chronic pancreatitis. On magnetic resonance cholangiopancreatography (MRCP) and EUS imaging, the pancreatic duct was observed to be dilated from the head of the pancreas, around which a pancreatic stone was also seen. This pancreatic stenosis was treated using a pancreatic stent. First, we performed ERCP, through which we observed the stenosis of the pancreatic duct. However, we were unable to pass the ERCP cannula through the stenosis site (▶ **Fig. 1**). Next, the scope was changed from a duodenoscope to a convex echoendoscope. To avoid any intervening vessels, the pancreatic duct was first punctured using a 19-G needle under Doppler imaging (▶ **Fig. 2**). After pancreatic juice had been aspirated, contrast medium was injected, and images of the pancreatic duct were obtained (▶ **Fig. 3**). Then, a 0.025-inch stiff guidewire was inserted, and was easily advanced into the duodenum through the stenosis. Next, the 19-G needle was exchanged for an ERCP catheter to dilate the fistula (▶ **Fig. 4**). Although an EUS-guided rendezvous approach was considered, EUS-guided antegrade stenting of the pancreatic duct was attempted because no leakage of pancreatic juice was seen on endosonographic imaging. The 7-Fr straight plastic stent was inserted antegradely, from the duodenum to the pancreatic duct (▶ **Fig. 5**). No adverse events were seen.

Although validation is required in a prospective clinical trial, this novel technique appears to be safe and effective as an option for EUS-guided pancreatic duct drainage.

Video 1

During endoscopic ultrasound (EUS)-guided pancreatic duct antegrade stenting, the main pancreatic duct is punctured using a 19-G needle, and contrast medium is injected. A 0.025-inch guidewire is inserted through the 19-G needle and advanced into the duodenum. To dilate the fistula, a cannula used for endoscopic retrograde cholangiopancreatography (ERCP) is inserted. Finally, a 7-Fr plastic stent is placed antegradely from the duodenum to the pancreatic duct.

Fig. 4 A 0.025-inch guidewire and ERCP cannula have been easily advanced into the duodenum through the stenosis.

Fig. 5 A 7-Fr plastic stent has been placed antegradely.

Endoscopy_UCTN_Code_TTT_1AS_2AD

Competing interests: None

**Takeshi Ogura¹, Saori Onda¹,
Tatsushi Sano¹, Kazuhiro Yamamoto²,
Daisuke Masuda¹, Akira Imoto¹,
Kazuhide Higuchi¹**

¹ Second Department of Internal Medicine,
Osaka Medical College, Osaka, Japan

² Department of Radiology, Osaka Medical
College, Osaka, Japan

References

- 1 Kahaleh M, Hernandez AJ, Tokar J et al. EUS-guided pancreaticogastrostomy: analysis of its efficacy to drain inaccessible pancreatic ducts. *Gastrointest Endosc* 2007; 65: 224–230
- 2 Itoi T, Yasuda I, Kurihara T et al. Technique of endoscopic ultrasonography-guided pancreatic duct intervention (with video). *J Hepatobiliary Pancreat Sci* 2014; 21: E4–E9
- 3 Giovannini M. Endoscopic ultrasonography-guided pancreatic drainage. *Gastrointest Endosc Clin N Am* 2012; 22: 221–230

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1391422>
Endoscopy 2015; 47: E161–E162
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Takeshi Ogura, PhD, MD
Second Department of Internal Medicine
Osaka Medical College
1-1 Daigakuchou, Takatsukishi
Osaka 464-8681
Japan
Fax: +81-52-7635233
oguratakeshi0411@yahoo.co.jp