Intra-abdominal organ strangulation during intraoperative endoscopic retrograde cholangiopancreatography with rendezvous cannulation: case report

Intraoperative rendezvous endoscopic retrograde cholangiopancreatography (ERCP) is one of the single-stage options available for managing common bile duct stones during laparoscopic cholecystectomy [1]. The rendezvous procedure allows immediate biliary cannulation and reduces the risk of post-ERCP pancreatitis [1,2]. We report a case of omentum and stomach strangulation caused by the trans-cystic guidewire during a routine rendezvous ERCP, an unusual complication.

A 41-year-old woman was admitted to the emergency department with a 3-month history of recurrent episodes of upper right abdominal pain. Examination with abdominal ultrasound demonstrated cholelithiasis. Because of the patient's intense abdominal pain, laparoscopic cholecystectomy was performed acutely. During perioperative cholangiography, stones were detected in the common bile duct, and it was decided to perform ERCP directly after the cholecystectomy.

At the beginning of the ERCP procedure, the guidewire was easily identified emerging from the papilla of Vater. During attempts to withdraw the guidewire, a high degree of resistance was encountered. Perioperative X-ray imaging revealed a twisted guidewire in the abdomen (● Fig. 1). Laparoscopy was performed immediately and showed the guidewire twisted around a structure that seemed to be part of the intestine, dark in color and with uncertain viability. Conversion of the procedure to laparotomy revealed strangulation of the omentum and anterior wall of the gastric antrum, with uncertain viability. The guidewire was cut into pieces and removed with the assistance of orthopedic scissors. The omentum and gastric antrum were fully recovered 20 minutes later (● Fig. 2), and the abdomen was closed routinely. The patient was able to mobilize the same day and the day after was discharged from the hospital in good clinical condition.

The possibility of the strangulation of intra-abdominal organs should be kept in mind when resistance that is hard to explain is encountered during extraction of a guidewire. Laparotomy remains the safest way to determine both viability of the abdominal organs and extraction of the guidewire [3–5].

Endoscopy_UCTN_Code_CPL_1AM_2AI

Competing interests: None

Ioannis Gkekas, Jonathan Palmgren, Pia Näsval, Michael Dahlberg

Department of Surgery, Sunderby Medical Center, Luleå, Sweden

References

- 1 Swahn F, Regnér S, Enochsson L et al. Endoscopic retrograde cholangiopancreatography with rendezvous cannulation reduces pancreatic injury. *World J Gastroenterol* 2013; 19: 6026–6034
- 2 Swahn F, Nilsson M, Arnelo U et al. Rendezvous cannulation technique reduces post-ERCP pancreatitis: a prospective nationwide study of 12,718 ERCP procedures. *Am J Gastroenterol* 2013; 108: 552–559
- 3 Bailey A, Bourke M, Williams S et al. A prospective randomized trial of cannulation technique in ERCP: effects on technical success and post-ERCP pancreatitis. *Endoscopy* 2008; 40: 296–301
- 4 Kwon C, Song SH, Hahm KB et al. Unusual complications related to endoscopic retrograde cholangiopancreatography and its endoscopic treatment. *Clin Endosc* 2013; 46: 251–259
- 5 Samei H, Baky RA, Dowais A. A rare post-ERCP complication: a case report of pneumoperitoneum, pneumoretroperitoneum, and surgical emphysema following unsuccessful ERCP. *Internet J Surg* 2008; 21

Fig. 1 Perioperative X-ray image shows a twisted guidewire in the abdomen of a patient undergoing endoscopic retrograde cholangiopancreatography during surgery.

Fig. 2 Fully recovered gastric antrum and the sign of strangulation by the guidewire.

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1391865>
Endoscopy 2015; 47: E231
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Ioannis Gkekas, MD
 Department of Surgery
 Sunderby Medical Center
 971 80 Luleå
 Sweden
 Fax: +46-920-283320
gkekasioannis@windowslive.com