Endoscopic ultrasound-guided duodenojejunal anastomosis to treat postsurgical Roux-en-Y hepaticojejunostomy stricture: a dream or a reality?

Fig. 1 Radiologic view of endoscopic ultrasound-guided puncture of the afferent loop in a 67-year-old man with recurrent cholangitis due to benign stricture of a bilioenteric anastomosis. The patient had a history of sclerosing cholangitis and left hepatectomy with Roux-en-Y bilioenteric reconstruction.

Fig. 2 Radiologic view of the stent between the two loops.

Roux-en-Y hepaticojejunostomy stricture develops in 8% to 40% of patients, and re-intervention is frequently required [1,2]. Surgery remains the gold standard to treat this problem. We present the case of a 67-year-old man with recurrent cholangitis due to benign stricture of a bilioenteric anastomosis. The patient had a history of sclerosing cholangitis and left hepatectomy with Roux-en-Y bilioenteric reconstruction.

In another hospital, the patient had been treated via a percutaneous trans-hepatic approach with an uncovered self-expandable metal stent (u-SEMS). Biliary lithiasis and several episodes of cholangitis recurred soon thereafter. Many attempts were made to remove the stones radiologically, but they all failed. Percutaneous internal-external drainage was then placed temporarily, and the patient was referred to our tertiary endoscopic center.

A peroral endoscopic procedure with a rendezvous technique was attempted by passing a guidewire through the internal-external drainage, but it did not succeed because the anastomotic loop was too long and tortuous. Therefore, a new endoscopic approach with an endoscopic ultrasound (EUS)-guided transenteric anastomosis was used. After the injection of contrast medium into the jejunal loop, an EUS-guided puncture was performed from the duodenal portion with a 19-gauge needle (ECHO-19, Cook Medical) (Fig. 1); a 0.035-guidewire was then placed into the jejunal loop, and a duodenojejunal fistula was created by pushing an 8.5-Fr cystoenterostome (XS 1341, Endoflex) on the guidewire. Finally, a 20-mm-long and 16-mm-diameter uSEMS (Nagi stent; Taewoong Medical) was left in place to allow consolidation of the endoscopic fistula (Fig. 2). After 3 days, an operative gastroscope (Pentax) was advanced through the stent to perform endoscopic retrograde cholangiopancreatography (ERC) and remove the previous uSEMS (Fig. 3). The patient was discharged after 2 days. No complications or procedure-related symptoms have been reported during 12 months of clinical follow-up.

EUS-guided duodenojejunal anastomosis is a feasible endoscopic approach in selected patients [3,4]. Expert and skilled endoscopists are needed to perform the described procedure successfully.

Endoscopy_UCTN_Code_TTT_1AS_2AB

Competing interests: None

Fig. 3 Passage of the scope through the created anastomosis.

**Massimiliano Mutignani¹,
Raffaele Manta¹, Francesco
Pugliese¹, Antonio Rampoldi²,
Lorenzo Dioscoridi¹, Edoardo Forti¹**

¹ Diagnostic and Interventional Digestive Endoscopy, Niguarda Ca' Granda Hospital, Milan, Italy

² Interventional Radiology Unit, Niguarda Ca' Granda Hospital, Milan, Italy

References

- 1 Chan SM, Teoh AY. Endoscopic ultrasound-guided biliary drainage: a review. *Curr Treat Options Gastroenterol* 2015; 13: 171–184
- 2 Siripun A, Sripongpun P, Ovarlarnporn B. Endoscopic ultrasound-guided biliary intervention in patients with surgically altered anatomy. *World J Gastrointest Endosc* 2015; 7: 283–289
- 3 Ogura T, Higuchi K. Technical tips of endoscopic ultrasound-guided choledochoduodenostomy. *World J Gastroenterol* 2015; 21: 820–828
- 4 Jürgensen C, Wentrup R, Zeitz M. Endoscopic ultrasound (EUS)-guided transduodenal drainage of an obstructed jejunal loop after hepaticojejunostomy as treatment for recurrent biliary sepsis. *Endoscopy* 2013; 45 (Suppl. 02): E40–E41

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1392424>
Endoscopy 2015; 47: E350–E351
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Massimiliano Mutignani, MD
Endoscopy Unit
Niguarda Ca' Granda Hospital
Piazza dell'Ospedale Maggiore 3
20162 Milan
Italy
Fax: +39-02-64442911
massimiliano.mutignani@ospedaleniguarda.it