A novel method for closure of a choledochoduodenostomy in sump syndrome

Fig. 1 Fluoroscopic image showing drainage of the contrast dye via the choledochoduodenostomy anastomosis (arrow).

A 57-year-old woman with a history of a complicated open cholecystectomy and side-to-side choledochoduodenostomy (CDD) 25 years previously was admitted to our hospital with cholangitis. She had undergone five endoscopic retrograde cholangiopancreatography (ERCP) procedures with stone and debris extractions in another hospital during the preceding 2 years.

ERCP revealed a patent biliary sphincterotomy and CDD orifice that was proximal to the papilla of Vater. Injection of contrast showed that dye drained via the CDD anastomosis (● **Fig. 1**). Several small pieces of debris were extracted using a balloon-tipped catheter (Eliminator Biliary Balloon Dilators; ConMed, Utica, New York, USA), and an 8-cm long, fully covered self-expanding metal stent (SEMS; WallFlex; Boston Scientific, Natick, Massachusetts, USA) was placed through the papilla of Vater (● **Fig. 2**). After a period of 3 months, the fully covered SEMS was retrieved uneventfully, and complete closure of the CDD orifice was observed during the ERCP procedure (● **Fig. 3a**). A repeat injection of contrast confirmed the closure of the CDD orifice with no evidence of any drainage of the contrast dye (● **Fig. 3b**).

Limited data exist regarding endoscopic closure of CDDs in sump syndrome. Ell et al. [1] used an amplatzer septal occluder for successful CDD closure; however, this technique seems to be difficult and may not be available in many endoscopy centers. Recently, Ellison et al. [2] used an over-the-scope clip; however, the success

rate of this method depends on the diameter of the CDD orifice. In addition, we believe that this method could result in formation of a biliary stricture. The new method presented, using a fully covered SEMS, is technically easy, safe, and effective for closure of a CDD in sump syndrome. We suggest that a minimum stent indwell time of 3 months is needed for successful closure of a CDD.

Endoscopy_UCTN_Code_TTT_1AR_2AG

Competing interests: None

Galip Ersoz, Fatih Tekin, Ilker Turan, Oktay Tekesin, Omer Ozutemiz

Department of Gastroenterology, Ege University Medical School, Izmir, Turkey

References

- 1 Ell C, Boosfeld C, Henrich R et al. Endoscopic treatment of the "sump syndrome" after choledochoduodenostomy: a new technique using an amplatzer septal occluder. *Z Gastroenterol* 2006; 44: 1231–1235
- 2 Ellison JC, Tarnasky PR. Endoscopic management of sump syndrome: closure of choledochoduodenostomy. *Gastrointest Endosc* 2014; 80: 725–726

Fig. 2 The fully covered self-expanding metal stent seen on: **a** fluoroscopic view; **b** endoscopic view showing it passing through the papilla of Vater; **c** endoscopic view through the choledochoduodenostomy anastomosis.

Fig. 3 **a** Endoscopic view 3 months later after retrieval of the self-expanding metal stent showing complete closure of the choledochoduodenostomy orifice. **b** Fluoroscopic image during endoscopic retrograde cholangiopancreatography (ERCP) 3 months later showing no evidence of drainage after injection of contrast dye, confirming the complete closure of the choledochoduodenostomy.

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1392504>
Endoscopy 2015; 47: E365–E366
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Fatih Tekin, MD
Ege Universitesi Tip Fakultesi
Gastroenteroloji Bilim Dalı
Bornova 35100
Izmir
Turkey
Fax: +90-232-3427764
drtekinfatih@gmail.com