Clip and snare lifting technique to assist cannulation of a papilla hidden behind a mucosal fold

Endoscopic retrograde cholangiopancreatography (ERCP) is a technically complex endoscopic procedure with significant rates of severe complications (0.8%) and mortality (0.02%) [1]. Complication rates increase in those patients where cannulation is difficult, defined as a situation in which the operator using a standard technique fails to achieve cannulation within 5 minutes, makes five unsuccessful attempts, or two passages of the guidewire into the pancreas [2,3]. Features that are known to be associated with difficult, time-consuming, or impossible cannulations are failure to identify the papilla within the diverticulum, a floppy papilla on the edge of diverticula, and a papilla that is hidden by a mucosal fold (● Fig. 1 a) [4]. The latter situation

can be particularly challenging and has few possible solutions [5].

We hereby describe the first two patients in which we used an endoclip and a snare to lift the mucosal fold and allow access to a hidden papilla. The first patient was a 70-year-old woman with a bile leak after hemihepatectomy. She had a flat papilla, underneath a floppy mucosal fold, that could not be cannulated. The second patient was a 55-year-old woman with a post-cholecystectomy bile leak. She had a papilla that was hidden behind a fold and had undergone two ERCPs and a precut sphincterotomy because of unsuccessful bile duct cannulation.

In both patients an endoclip (Resolution Clip; Boston Scientific Corporation, Natick, Massachusetts, USA) was placed on

the fold. The duodenoscope was withdrawn and was then re-introduced with a biopsy forceps inside the working channel that was used to grasp a polypectomy snare (Sensation Short Throw, 13 mm; Boston Scientific) so that this was introduced beside the scope. The snare loop was then tightened slightly around the shaft of the endoclip (● Fig. 1 b) and the wire was pulled up through the mouth (● Fig. 1 c), which slightly altered the alignment of the papillary region, so allowing the opening of the papilla to be visualized (● Fig. 1 d) and successfully cannulated.

These are the first successful cases of clip and snare-assisted cannulation, a way of accessing the bile duct in otherwise apparently impossible conditions, by either pulling or pushing (● Fig. 2) a snared clip that had been previously positioned on the mucosal fold that was hiding the papilla.

Endoscopy_UCTN_Code_TTT_1AR_2AB

Fig. 1 Endoscopic views showing: **a** the mucosal fold with the papilla hidden behind it; **b, c** the snare being tightened around the previously positioned endoscopic clip, and being pulled back; **d** the papilla, which is successfully visualized after use of a push-and-pull technique, subsequently allowing the common bile duct to be cannulated.

Competing interests: None

Roberto Valente^{1,2}, Francisco Baldaque-Silva¹, Antti Siiki³, Jeanne Lübke⁴, Lars Enochson¹, Matthias Lohr¹, Urban Arnelo¹

¹ Pancreatic Surgery Unit, Department of Clinical Science, Intervention and Technology (CLINTEC), Karolinska Institute, Stockholm, Sweden

² Digestive and Liver Disease Unit, Sant'Andrea Hospital, Sapienza University of Rome, Rome, Italy

³ Department of Gastroenterology and Alimentary Tract Surgery, Tampere University Hospital, Tampere, Finland

⁴ Division of Surgery, Tygerberg Hospital and the University of Stellenbosch, Tygerberg, South Africa

References

- 1 Salminen P, Laine S, Gullichsen R. Severe and fatal complications after ERCP: analysis of 2555 procedures in a single experienced center. *Surg Endosc* 2008; 22: 1965–1970
- 2 Udd M, Kylänpää L, Halttunen J. Management of difficult bile duct cannulation in ERCP. *World J Gastrointest Endosc* 2010; 2: 97–103
- 3 Halttunen J, Meisner S, Aabakken L et al. Difficult cannulation as defined by a prospective study of the Scandinavian Association for Digestive Endoscopy (SADE) in 907 ERCPs. *Scand J Gastroenterol* 2014; 49: 752–758
- 4 Tham TC, Kelly M. Association of periampullary duodenal diverticula with bile duct stones and with technical success of endoscopic retrograde cholangiopancreatography. *Endoscopy* 2004; 36: 1050–1053
- 5 Fujita N, Noda Y, Kobayashi G et al. ERCP for intradiverticular papilla: two-devices-in-one-channel method. *Endoscopic Retrograde Cholangiopancreatography. Gastrointest Endosc* 1998; 48: 517–520

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1392923>
Endoscopy 2015; 47: E517–E518
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Urban Arnelo, MD, PhD

Department for Clinical Science, Intervention and Technology (CLINTEC) – Division of Surgery
 Karolinska Institute
 Huddinge, K53
 Stockholm S-14186
 Sweden
 Urban.Arnelo@ki.se