Overtube-guided endoscopic extraction of a rectal foreign body: lifting not only the embargo

Fig. 1 Plain radiograph of the abdomen of a 65-year-old man presenting to the emergency department shows a long, radio-paque, cylindrical foreign body in the rectum and sigmoid.

Fig. 2 Proctosigmoidoscopy revealing a cigar case: **a** in the rectum; **b** in the sigmoid.

A 65-year-old man presented to the emergency department with a foreign body in his rectum. He had inserted a Cuban cigar case into his anus approximately 9 hours before admission. The case had no sharp edges, and its cap was on.

The patient's abdomen was soft and not distended. Abdominal radiography showed a long, radiopaque, cylindrical foreign body in the rectum and sigmoid (● Fig. 1), with no signs of perforation. Proctosigmoidoscopy, performed without sedation, confirmed the presence of the object (● Fig. 2). Several attempts were made to remove it with a polypectomy snare, but the foreign body was very long

and inflexible, so that it could not pass through the acute angle of the rectosigmoid junction.

An esophageal overtube was then introduced up to the proximal edge of the cylindrical case, reducing the angle of the rectosigmoid junction so that the distal tip of the case could be disimpacted (● Video 1). With its distal tip fixed by a polypectomy snare, the foreign body was moved to the distal rectum and finally removed with manual assistance (● Fig. 3). Afterward, the endoscope was removed, and no complication followed the procedure. The patient was asymptomatic at discharge.

Video 1

Overtube-guided extraction of a rectal foreign body.

Retained rectal foreign bodies are not uncommon and often present a management dilemma [1,2]. Frequently, hospital admission is delayed until after several efforts have been made to remove the object at home [1]. Extraction can be difficult, and severe complications may occur [1].

Low-lying foreign bodies are usually palpable on digital rectal examination and can often be removed manually [3]. High-lying foreign bodies are located proximally to the rectosigmoid junction, and proctosigmoidoscopy is required for removal [3,4]. Surgery should be reserved for unstable patients, those with perforation, and those in whom transanal attempts at removal have failed [1].

To the best of our knowledge, no previous cases in which this technique was used have been described. The use of an overtube can straighten the colorectum, facilitating the removal of a rectal foreign body.

Endoscopy_UCTN_Code_TTT_1AQ_2AH

Competing interests: None

Marco Silva, Andreia Albuquerque, Armando Ribeiro, Hélder Cardoso, Guilherme Macedo

Gastroenterology Department, Centro Hospitalar São João, Faculty of Medicine of the University of Porto, Porto, Portugal

Fig. 3 The cigar case after removal and the esophageal overtube used during the extraction.

References

- 1 Coskun A, Erkan N, Yakan S et al. Management of rectal foreign bodies. *World J Emerg Surg* 2013; 8: 11
- 2 Cologne KG, Ault GT. Rectal foreign bodies: what is the current standard? *Clin Colon Rectal Surg* 2012; 25: 214–218
- 3 Anderson KL, Dean AJ. Foreign bodies in the gastrointestinal tract and anorectal emergencies. *Emerg Med Clin North Am* 2011; 29: 369–400
- 4 Lyons MF, Tsuchida AM. Foreign bodies of the gastrointestinal tract. *Med Clin North Am* 1993; 77: 1101–1114

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1393392>
Endoscopy 2015; 47: E563–E564
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Marco Silva, MD
 Gastroenterology Department
 Centro Hospitalar São João
 Alameda Professor Hernâni Monteiro
 4200-319 Porto
 Portugal
 Fax: +351-225-025-766
marcocostasilva87@gmail.com