Protective Effect of Tetrahydrocurcumin against Cisplatin-Induced Renal Damage: *In Vitro* and *In Vivo* Studies

Authors

Kyung Il Song^{1*}, Jun Yeon Park^{2*}, Seungyong Lee^{3,4}, Dahae Lee^{2,4}, Hyuk-Jai Jang⁵, Su-Nam Kim³, Hyeonseok Ko⁶, Hyun Young Kim⁴, Jae Wook Lee³, Gwi Seo Hwang², Ki Sung Kang², Noriko Yamabe²

Affiliations

The affiliations are listed at the end of the article

Key words

- cisplatin
- nephrotoxicity
- tetrahydrocurcumin
- oxidative renal damage
- COX-2

received July 23, 2014
revised January 15, 2015
accepted January 16, 2015

Bibliography

DOI <http://dx.doi.org/10.1055/s-0035-1545696>
 Published online February 26, 2015
 Planta Med 2015; 81: 286–291
 © Georg Thieme Verlag KG
 Stuttgart · New York ·
 ISSN 0032-0943

Correspondence

Noriko Yamabe
 College of Korean Medicine
 Gachon University
 1342 SeongnamDaero
 Seongnam 461–701
 Korea
 Phone: + 82 3 17 50 54 21
 Fax: + 82 3 17 50 54 16
 norikoy@gachon.ac.kr

Correspondence

Ki Sung Kang
 College of Korean Medicine
 Gachon University
 1342 SeongnamDaero
 Seongnam 461–701
 Korea
 Phone: + 82 3 17 50 54 02
 Fax: + 82 3 17 50 54 16
 kkang@gachon.ac.kr

Abstract

▼ The adverse effects of anticancer drugs can prompt patients to end their treatment despite the efficacy. Cisplatin is a platinum-based molecule widely used to treat various forms of cancer, but frequent and long-term use of cisplatin is limited due to severe nephrotoxicity. In the present study, we investigated the protective effect and mechanism of tetrahydrocurcumin on cisplatin-induced kidney damage, oxidative stress, and inflammation to evaluate its possible use in renal damage. Cisplatin-induced LLC-PK1 renal cell damage was significantly reduced by tetrahydrocurcumin treatment. Additionally, the protective effect of tetrahydrocurcumin on cisplatin-induced oxidative renal damage was investigated

in rats. Tetrahydrocurcumin was orally administered every day at a dose of 80 mg/kg body weight for ten days, and a single dose of cisplatin was administered intraperitoneally (7.5 mg/kg body weight) in 0.9% saline on day four. The creatinine clearance levels, which were markers of renal dysfunction, in cisplatin-treated rats were recovered nearly back to normal levels after administration of tetrahydrocurcumin. Moreover, tetrahydrocurcumin exhibited protective effects against cisplatin-induced oxidative renal damage in rats by inhibiting cyclooxygenase-2 and caspase-3 activation. These results collectively provide therapeutic evidence that tetrahydrocurcumin ameliorates renal damage by regulating inflammation and apoptosis.

Introduction

▼ The usual treatments for cancer are surgery, chemotherapy, radiation, or a combination of these methods [1,2]. Chemotherapy is the use of anticancer drugs to treat cancerous cells. It has been used for many years, and is one of the most common treatments for cancer [3,4]. Anticancer drugs interfere with the growth of tumor cells, eventually causing their death. However, they may also affect the growth of normal cells, causing many adverse effects, some of which may be serious [5]. Cisplatin is a platinum-based anticancer drug widely used to treat various forms of cancer in humans [6,7]. However, this treatment has some major limitations for use. Within an hour following injection, some patients suffer from side effects that include perception and hearing disorders, tinnitus, and inner ear dysfunctions that

cause dizziness [8,9]. In addition, frequent and long-term use of cisplatin is restricted because of severe nephrotoxicity [10,11]. These side effects can prompt patients to stop treatment despite its efficacy. Therefore, a better understanding of these side effects is crucial for the continued clinical use of cisplatin.

We have investigated naturally occurring antioxidants that can protect the kidney from cisplatin-induced damage [12–14]. Many researchers have shown the involvement of oxidative stress in cisplatin-induced nephrotoxicity [15,16]. Cisplatin induces the generation of various reactive oxygen species (ROS) by inactivating the cellular antioxidant system, disrupting the mitochondrial respiratory chain, or interacting with microsomal cytochrome P450 [17,18]. Natural antioxidants provide beneficial effects against cisplatin chemotherapy, not only by protecting the kidney from cisplatin-induced oxidative damages, but also by inducing cancer cell death. For example, we have reported that antioxidants from *Panax ginseng* exhibited both renoprotective and anticancer ef-

* These two authors contributed equally to the work described in this study.

Fig. 1 Effect of tetrahydrocurcumin on cisplatin-induced nephrotoxicity in LLC-PK1 cells. **A** Chemical structure of THC. **B** Representative microscopic images showing the protective effect of THC against cisplatin-induced nephrotoxicity in cells. **C** Dose-dependent protective effect of THC against cisplatin-induced nephrotoxicity in cells. **D** No cytotoxic effects of THC at the same doses, which was effectively protected from cisplatin-induced cell damage. $P < 0.05$ compared to the cisplatin-treated control value. (Color figure available online only.)

fects, providing evidence for its use as a chemotherapeutic adjuvant [19,20].

Curcumin is an active ingredient from the plant *Curcuma longa* L. (Zingiberaceae), which has various pharmacological effects, including anti-inflammatory, antioxidant, and anticarcinogenic activities [21–23]. Tetrahydrocurcumin (THC) is an active metabolite of curcumin found in the gastrointestinal tract and a reduced analog of curcumin with phenolic and β -diketo moieties (● **Fig. 1 A**). THC possesses stronger antioxidant activity than other curcuminoids, including curcumin, demethoxycurcumin, and bisdemethoxycurcumin [24,25]. However, little is known about the effects of THC on cisplatin-induced nephrotoxicity. Therefore, the *in vitro* and *in vivo* effects and molecular mechanisms of THC on cisplatin-induced nephrotoxicity were investigated in this study.

Results and Discussion

Recent literature indicates that ROS play critical roles in the development and progression of kidney damage [26–29]. Antioxidant treatment can prevent oxidative damage and might delay

the progression of kidney disease. Therefore, the use of antioxidant agents as inhibitors against oxidative stress may be considered an important therapeutic approach for kidney disease. In the present study, we have investigated the protective effects and mechanism of THC on cisplatin-induced kidney damage, oxidative stress, and inflammation to evaluate its possible use in renal damage.

The protective effect of THC against cisplatin-induced oxidative damage was tested using LLC-PK1 cells, which are renal tubular cells that are the most vulnerable renal tissue to oxidative stress [26]. ● **Fig. 1 B** shows the morphological changes of cisplatin-treated LLC-PK1 cells that were also treated with THC. As shown in ● **Fig. 1 C**, 25 μ M cisplatin treatment significantly decreased cell viability to about 50% of that of untreated control cells. In contrast, pretreatment with THC markedly restored cell viability in a dose-dependent manner. However, THC demonstrated no cytotoxic effects at the same doses, which was effectively protected from cisplatin-induced cell damage (● **Fig. 1 D**). Then, we further examined the effect of THC on cisplatin-induced oxidative renal damage in rats.

The body weight gains of rats after cisplatin and cisplatin + THC treatments were markedly reduced compared to normal rats

Fig. 2 Effect of tetrahydrocurcumin on body weight, kidney weight, food intake, and the renal function parameter in the cisplatin-induced renal damage rat model. **A** Changes in body weight. **B** Changes in food intake. **C** Changes in kidney weight. **D** Creatinine clearance. $P < 0.05$ compared to the cisplatin-treated control value.

(● Fig. 2A). Similarly, food intake amounts were slightly lowered after cisplatin treatments and gradually recovered in the vehicle-treated groups (● Fig. 2B). The sharp decrease in the food intake was observed on day four in cisplatin and cisplatin + THC treated groups. These results are in accordance with earlier reports that noted a decrease in body weight gain after cisplatin injection [30]. Cisplatin intoxication resulted in a significant body weight reduction and kidney weight increase. However, these changes were not significantly ameliorated by THC treatment (● Fig. 2C). Regarding renal functional parameters, cisplatin-injected rats exhibited decreased creatinine clearance levels compared to those of the vehicle-treated group. The reduced creatinine clearance level of cisplatin-treated rats was significantly recovered by co-treatment with THC (● Fig. 2D).

● Fig. 3A shows the effect of THC on organic cation transporter-2, cyclooxygenase-2, and procaspase-3 protein expression in cisplatin-treated rat kidneys. OCT-2 is a renal uptake transporter that plays a key role in disposition and the renal clearance of drugs and endogenous compounds [31]. OCT-2 protein expression in kidney tissue was slightly increased after cisplatin treatment and was decreased by THC treatment, but the differences were not statistically significant (● Fig. 3B). COX-2 is an inducible component of the prostaglandin synthesis cascade and is inducible in normal cells by many cytokines, mitogens, and proinflammatory factors [32]. Recent studies have demonstrated that COX-2 is highly expressed in response to inflammation in kidney tissue. Therefore, it may be intimately involved in prostaglandin-dependent renal inflammatory processes [33]. Thus, the development of COX-2 inhibitors serves as a paradigm for molecularly targeted renoprotective agents. Caspase-3 is potentially the most

important effector enzyme in apoptosis, providing a common pathway to both death receptor- and mitochondria-dependent apoptotic mechanisms [34, 35]. COX-2 and procaspase-3 protein expression levels were significantly increased after cisplatin-injection, but cisplatin co-treatment with THC resulted in almost complete renoprotection. These results imply that THC may alleviate oxidative stress by preventing caspase-3 activation and related inflammation in the kidney.

PAS staining was performed on renal sections to measure tubular damage. As shown in the representative pictures of renal sections, severe tubulointerstitial injuries, including tubular epithelial cell detachments, cystic dilatation of tubules, and inflammatory cell infiltration, occurred in the kidneys of cisplatin-treated animals (● Fig. 4). However, the increased tubular damage in cisplatin-treated rats was reduced by co-treatments with THC (● Fig. 4). Therefore, THC was effective in alleviating cisplatin-induced tubulointerstitial injuries.

In summary, kidney cell damage induced by cisplatin was significantly inhibited by treatment with THC. In addition, the renal dysfunction of cisplatin-treated rats was markedly ameliorated by THC extract administration. The renoprotective effect of THC was associated with the caspase-dependent anti-inflammatory pathway. Taken together, these results demonstrate that THC exerted a renoprotective effect in cisplatin-treated rats and, therefore, its use can be considered to prevent kidney damage during or after chemotherapy.

Fig. 3 Effect of tetrahydrocurcumin on protein expression levels of organic cation transporter-2, cyclooxygenase-2, and procaspase-3 of the rat kidney. **A** OCT-2 (60 kDa), COX-2 (72 kDa), and cleaved caspase-3 (17, 19 kDa) protein expressions evaluated by Western blot analysis. GAPDH (37 kDa) was used as an internal control. **B** Quantitative data for the OCT-2, COX-2, and cleaved caspase-3 Western blot analysis. Western blot assays were done in triplicate for each protein and were repeated at least three times. $P < 0.05$ compared to the cisplatin-treated control value.

Fig. 4 Effect of THC on the cisplatin-induced tubular damage in renal cortex tissues. **A** PAS staining of a representative renal section. **B** Quantitative data for the tubular damage of PAS staining. Original magnification of **A** is $\times 100$. $P < 0.05$ compared to the cisplatin-treated control value. (Color figure available online only.)

Materials and Methods

Chemicals and reagents

THC and cisplatin (purity 98–100%) were purchased from Sigma. A stock solution of chemicals for cell-based assays was prepared in 100% DMSO and stored at -20°C until use. When required, the stock solution was diluted with cell culture media to the appropriate concentration. The final concentrations of DMSO in the culture media were adjusted to less than 0.5% (v/v), which exhibits no toxic effect for cells. Dulbecco's modified Eagle's medium (DMEM) was purchased from Cellgro and FBS was from Invitrogen. OCT-2, COX-2, cleaved caspase-3, GAPDH, and horseradish peroxidase (HRP) conjugated anti-rabbit antibodies were purchased from Cell Signaling and EZ-Cytox reagent was from Daeil Lab Service.

Renoprotective effect against cisplatin-induced damage in kidney cells

The renoprotective effect against oxidative renal damage was evaluated using LLC-PK1 cells [26]. LLC-PK1 (pig kidney epithelium, CL-101) cells were purchased from ATCC and cultured in DMEM supplemented with 10% FBS, 1% penicillin/streptomycin, and 4 mM L-glutamine at 37°C with 5% CO_2 in the air. The cells were seeded in 96-well culture plates at 1×10^4 cells per well and allowed to adhere for 2 h. Thereafter, the test sample and/or radical donor, 25 μM cisplatin, were added to culture medium. Twenty-four hours later, the medium containing the test sample and/or radical donor was removed. The cells were incubated with serum free medium (90 μL /well) and Ez-Cytox reagent (10 μL /well) at 37°C for 2 h. Cell viability was measured by absorbance at 450 nm using a microplate reader.

Renoprotective effect against cisplatin-induced oxidative damage in rats

All procedures involving the use of live animals as described in this study were approved in May 2014 by the Institutional Animal Care and Use Committee of the Gachon University (approval number GIACUC-R2014002) and strictly followed the NIH guidelines for the humane treatment of animals. Male Wistar rats weighing 140–160 g were used for evaluating the protection of THC against cisplatin-induced nephrotoxicity. The rats were housed with a temperature of $23 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ humidity [conditions with a standard light (12 h light/dark)]. The rats were given free access to water and a normal diet containing 10 kcal% fat for a period of one week after their arrival and were then divided into four groups based on their body weight and assigned to the vehicle, THC, cisplatin, or cisplatin + THC.

Group 1: vehicle ($n = 4$), received water (no sample treatment); group 2: THC ($n = 4$), treated with THC (80 mg/kg) in aqueous solution orally for ten days; group 3: cisplatin ($n = 4$) received water (no sample treatment); group 4: cisplatin + THC ($n = 4$), treated with THC (80 mg/kg) in aqueous solution orally for ten days.

THC was orally administered every day at a dose of 80 mg/kg body weight, while vehicle-treated rats were orally given water. The 80 mg/kg of THC dosage was chosen according to the literature [25]. After four days, rats in two groups (cisplatin and cisplatin + THC) were administered a single dose of cisplatin intraperitoneally (7.5 mg/kg body weight) in 0.9% saline. Animals in the vehicle group received an equivalent amount of normal saline for ten days. The rats were sacrificed six days after cisplatin administration under light ether anesthesia. Twenty-four hour urine samples were collected using a metabolic cage. Blood sam-

ples were collected from the abdominal aorta, and the kidneys were removed. All the preparations and analyses of various parameters were performed simultaneously under similar experimental conditions to avoid any day-to-day variations. During the experimental period, their body weights were measured daily.

Plasma biomarker analyses

Blood samples were collected in tubes containing 0.18 M EDTA and centrifuged at 5000 rpm for 5 min at 4°C . After centrifugation, the plasma was separated for the estimation of creatinine. Creatinine levels were determined by a rate-blanked kinetic Jaffe method. Creatinine clearance was calculated on the basis of the urinary Cr, serum Cr, urine volume, and body weight using the following equation:

$$\text{Creatinine clearance (mL/min/g body weight)} = [\text{urinary Cr (mg/dL)} \times \text{urine volume (mL)/serum Cr (mg/dL)}] \times [1000/\text{body weight (g)}] \times [1/1440 (\text{min})]$$

Histological analysis of kidney

Kidney samples were fixed in 10% buffered formalin phosphate, dehydrated, embedded in paraffin, sectioned in 3 μm thickness, and stained with periodic acid-Schiff (PAS) reagent (Fisher Scientific) for histological examination. Tubular damage in PAS-stained sections was examined under a microscope and scored based on the percentage of cortical tubules showing epithelial necrosis: 0, normal; 1, 1–25%; 2, 26–50%; 3, 51–75%; 4, > 76%. Tubular necrosis was defined as the loss of the proximal tubular brush border, blebbing of apical membranes, tubular epithelial cell detachment from the basement membrane, or intraluminal aggregation of cells and proteins, as described previously [36].

Preparation of whole-cell extracts from tissue

Whole-cell extracts from the kidney tissue were prepared according to the manufacturer's (Cell Signaling) instructions using RIPA buffer supplemented with a 1 $\times$ protease inhibitor cocktail and 1 mM phenylmethylsulfonyl fluoride (PMSF, Sigma).

Western blot analysis

Proteins (whole-cell extracts, 30 μg /lane; nuclear extracts, 10 μg /lane; cytosolic extracts, 20 μg /lane) were separated by SDS-PAGE, transferred to polyvinylidene fluoride (PVDF) membranes (Millipore) for 1 h at semi-dry, and blocked in blocking buffer for 1 h at room temperature. The PVDF membranes were incubated with the primary antibody against OCT-2 (1:1000 dilution), COX-2 (1:1000 dilution), cleaved caspase-3 (1:1000 dilution), and GAPDH (1:1000 dilution) overnight at 4°C , washed three times for 5 min in wash buffer, incubated with the HRP-conjugated secondary antibody (1:2000 dilution, anti-rabbit) for 1 h at room temperature, washed three times, and then detected with enhanced chemiluminescence (ECL) solution (GE Healthcare).

Statistical analysis

Statistical significance was determined through one-way analysis of variance (ANOVA) followed by a multiple comparison test with a Bonferroni adjustment. P values less than 0.05 were considered statistically significant.

Acknowledgements

This research was funded by the Gangneung Asan Hospital Biomedical Research Center Promotion Fund and Gachon University Research Fund 2014 (GCU-2014-0116).

Conflict of Interest

The authors declare no conflict of interest.

Affiliations

- ¹ Department of Internal Medicine, University of Ulsan College of Medicine, Gangneung Asan Hospital, Gangneung, Korea
- ² College of Korean Medicine, Gachon University, Seongnam, Korea
- ³ Natural Products Research Institute, Korea Institute of Science and Technology, Gangneung, Korea
- ⁴ Department of Food Science, Gyeongnam National University of Science and Technology, Jinju, Korea
- ⁵ Department of Surgery, University of Ulsan College of Medicine, Gangneung Asan Hospital, Gangneung, Korea
- ⁶ Laboratory of Molecular Oncology, Cheil General Hospital & Women's Healthcare Center, Kwandong University College of Medicine, Seoul, Korea

References

- 1 Cai XZ, Huang WY, Qiao Y, Du SY, Chen Y, Chen D, Yu S, Che RC, Liu N, Jiang Y. Inhibitory effects of curcumin on gastric cancer cells: a proteomic study of molecular targets. *Phytomedicine* 2013; 20: 495–505
- 2 Wilken R, Veena MS, Wang MB, Srivatsan ES. Curcumin: a review of anti-cancer properties and therapeutic activity in head and neck squamous cell carcinoma. *Mol Cancer* 2011; 10: 12
- 3 Hurley LH. DNA and its associated processes as targets for cancer therapy. *Nat Rev Cancer* 2002; 2: 188–200
- 4 Miltenburg NC, Boogerd W. Chemotherapy-induced neuropathy: A comprehensive survey. *Cancer Treat Rev* 2014; 40: 872–882
- 5 Chorawala MR, Oza PM, Shah GB. Mechanisms of anticancer drugs resistance: an overview. *Int J Pharm Sci Drug Res* 2012; 4: 1–9
- 6 Arany I, Safirstein RL. Cisplatin nephrotoxicity. *Semin Nephrol* 2003; 23: 23460–23464
- 7 Zhu M, Chen J, Yin H, Jiang H, Wen M, Miao C. Propofol protects human umbilical vein endothelial cells from cisplatin-induced injury. *Vascul Pharmacol* 2014; 61: 72–79
- 8 Kohn S, Fradis M, Ben-David J, Zidan J, Robinson E. Nephrotoxicity of combined treatment with cisplatin and gentamicin in the guinea pig: glomerular injury findings. *Ultrastruct Pathol* 2002; 26: 371–382
- 9 Li Q, Tian Y, Li D, Sun J, Shi D, Lin F, Gao Y, Liu H. The effect of lipocisplatin on cisplatin efficacy and nephrotoxicity in malignant breast cancer treatment. *Biomaterials* 2014; 35: 6462–6472
- 10 Luke DR, Vadiei K, Lopez-Berestein G. Role of vascular congestion in cisplatin induced acute renal failure in the rat. *Nephrol Dial Transplant* 1992; 7: 1–7
- 11 Miller RP, Tadagavadi RK, Ramesh G, Reeves WB. Mechanisms of Cisplatin nephrotoxicity. *Toxins (Basel)* 2010; 2: 2490–2518
- 12 Kang KS, Ham J, Kim YJ, Park JH, Cho EJ, Yamabe N. Heat-processed *Panax ginseng* and diabetic renal damage: active components and action mechanism. *J Ginseng Res* 2013; 37: 379–388
- 13 Kim JH, Han IH, Yamabe N, Kim YJ, Lee W, Eom DW, Choi P, Cheon GJ, Jang HJ, Kim SN, Ham J, Kang KS. Renoprotective effects of Maillard reaction products generated during heat treatment of ginsenoside Re with leucine. *Food Chem* 2014; 15: 114–121
- 14 Lee JH, Lee W, Lee S, Jung Y, Park SH, Choi P, Kim SN, Ham J, Kang KS. Important role of Maillard reaction in the protective effect of heat-processed ginsenoside Re-serine mixture against cisplatin-induced nephrotoxicity in LLC-PK1 cells. *Bioorg Med Chem Lett* 2012; 22: 5475–5479
- 15 Nath KA, Norby SM. Reactive oxygen species and acute renal failure. *Am J Med* 2000; 109: 665–678
- 16 Taguchi T, Nazneen A, Abid MR, Razzaque MS. Cisplatin-associated nephrotoxicity and pathological events. *Contrib Nephrol* 2005; 148: 107–121
- 17 Jiang M, Dong Z. Regulation and pathological role of p53 in cisplatin nephrotoxicity. *Pharmacology* 2008; 327: 300–307
- 18 Pabla N, Dong Z. Cisplatin nephrotoxicity: mechanisms and renoprotective strategies. *Kidney Int* 2008; 73: 994–1007
- 19 Yamabe N, Song KI, Lee W, Han IH, Lee JH, Ham J, Kim SN, Park JH, Kang KS. Chemical and free radical-scavenging activity changes of ginsenoside Re by Maillard reaction and its possible use as a renoprotective agent. *J Ginseng Res* 2012; 36: 256–262
- 20 Yamabe N, Kim YJ, Lee SY, Cho EJ, Park SH, Ham JY, Kim HY, Kang KS. Increase in antioxidant and anticancer effects of ginsenoside Re-lysine mixture by Maillard reaction. *Food Chem* 2013; 138: 876–883
- 21 Huang MT, Lysz T, Ferraro T, Abidi TF, Laskin JD, Conney AH. Inhibitory effects of curcumin on *in vitro* lipoxygenase and cyclooxygenase activities in mouse epidermis. *Cancer Res* 1991; 51: 813–819
- 22 Huang MT, Ma W, Yen P, Xie JG, Han J, Frenkel K, Grunberger D, Conney AH. Inhibitory effects of topical application of low doses of curcumin on 12-O-tetradecanoylphorbol-13-acetate-induced tumor promotion and oxidized DNA bases in mouse epidermis. *Carcinogenesis* 1997; 18: 83–88
- 23 Sandur SK, Ichikawa H, Pandey MK, Kunnumakkara AB, Sung B, Sethi G, Aggarwal BB. Role of pro-oxidants and antioxidants in the anti-inflammatory and apoptotic effects of curcumin (diferuloylmethane). *Free Radic Biol Med* 2007; 43: 568–580
- 24 Yoysungnoen P, Wirachwong P, Changtam C, Suksamrarn A, Patumraj S. Anti-cancer and anti-angiogenic effects of curcumin and tetrahydrocurcumin on implanted hepatocellular carcinoma in nude mice. *World J Gastroenterol* 2008; 7: 2003–2009
- 25 Murugan P, Pari L. Effect of tetrahydrocurcumin on lipid peroxidation and lipids in streptozotocin-nicotinamide-induced diabetic rats. *Basic Clin Pharmacol Toxicol* 2006; 99: 122–127
- 26 Yokozawa T, Cho EJ, Hara Y, Kitani K. Antioxidative activity of green tea treated with radical initiator 2, 2'-azobis (2-amidinopropane) dihydrochloride. *J Agric Food Chem* 2000; 48: 5068–5073
- 27 Forbes JM, Coughlan MT, Cooper ME. Oxidative stress as a major culprit in kidney disease in diabetes. *Diabetes* 2008; 57: 1446–1454
- 28 Heyman SN, Rosen S, Rosenberger C. A role for oxidative stress. *Contrib Nephrol* 2011; 174: 138–148
- 29 Li JM, Shah AM. ROS generation by nonphagocytic NADPH oxidase: potential relevance in diabetic nephropathy. *J Am Soc Nephrol* 2003; 14: 221–226
- 30 Takeda H, Sadakane C, Hattori T, Katsurada T, Ohkawara T, Nagai K, Asaka M. Rikkunshito, an herbal medicine, suppresses cisplatin-induced anorexia in rats via 5-HT₂ receptor antagonism. *Gastroenterology* 2008; 134: 2004–2013
- 31 Jonker JW, Schinkel AH. Pharmacological and physiological functions of the polyspecific organic cation transporters: OCT1, 2, and 3 (SLC22A1–3). *J Pharmacol Exp Ther* 2004; 308: 2–9
- 32 Harris RC. Cyclooxygenase-2 in the kidney. *J Cardiovasc Pharmacol* 2006; 47: 37–42
- 33 Kim GH. Renal effects of prostaglandins and cyclooxygenase-2 inhibitors. *Electrolyte Blood Press* 2008; 6: 35–41
- 34 Cohen GM. Caspases: the executioners of apoptosis. *Biochem* 1997; 326: 1–16
- 35 Yang B, Johnson TS, Thomas GL, Watson PF, Wagner B, Nahas AM. Apoptosis and caspase-3 in experimental anti-glomerular basement membrane nephritis. *J Am Soc Nephrol* 2001; 12: 485–495
- 36 Mukhopadhyay P, Rajesh M, Pan H, Patel V, Mukhopadhyay B, Bátkai S, Gao B, Haskó G, Pacher P. Cannabinoid-2 receptor limits inflammation, oxidative/nitrosative stress, and cell death in nephropathy. *Free Radic Biol Med* 2010; 48: 457–467