S. ZHU, N. NILJANSKUL, S. L. BUCHWALD* (MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, USA)
 A Direct Approach to Amines with Remote Stereocentres by Enantioselective CuH-Catalysed Reductive Relay Hydroamination
Nat. Chem. **2016**, *8*, 144–150.

Aliphatic Amines Formed by Asymmetric Copper-Catalyzed Hydroamination

Selected examples:

Postulated mechanism:

Significance: Aliphatic amines are present in many pharmaceutical agents and they are often essential for activity. The potential of stereoisomers to have contrasting biological effects means that asymmetric approaches to construct amine-containing remote stereocenters has been a challenge. Buchwald and co-workers report an asymmetric copper-catalyzed hydroamination reaction that generates γ - and δ -chiral aliphatic amines.

Comment: The copper-catalyzed hydroamination reaction produced the desired aliphatic amines in good to excellent yields and with excellent enantio- and diastereoselectivities. The proposed mechanism starts with the formation of an alkylcopper species and subsequent β -alkoxide elimination to form the enantioenriched alkene. The hydroxylamine reagent regenerates the CuH catalyst, which undergoes the anti-Markovnikov hydrocupration to form the terminal alkylcopper species. Finally, interception with the hydroxylamine generates the product, and the hydroxylamine reagent regenerates the CuH catalyst.

SYNFACTS Contributors: Mark Lautens, Hyung Yoon
 Synfacts 2016, 12(4), 0367 Published online: 15.03.2016
 DOI: 10.1055/s-0035-1561818; Reg-No.: L01816SF