A. E. COTMAN, D. CAHARD, B. MOHAR* (UNIVERSITY OF LJUBLJANA AND NATIONAL INSTITUTE OF CHEMISTRY, LJUBLJANA, SLOVENIA; UNIVERSITÉ ET INSA DE ROUEN, MONT SAINT AIGNAN, FRANCE)

Stereoarrayed CF₃-Substituted 1,3-Diols by Dynamic Kinetic Resolution: Ruthenium(II)-Catalyzed Asymmetric Transfer Hydrogenation

Angew. Chem. Int. Ed. **2016**, *55*, 5294–5298.

Ruthenium(II)-Catalyzed Asymmetric Transfer Hydrogenation Reaction

Transfer hydrogenation catalysts:

Selected examples:

Significance: In this report, ruthenium-based transfer hydrogenation catalysts are described. The catalysts exhibit impressive turnover numbers with excellent enantioselectivities for the selective reduction of CF₃-substituted 1,3-dicarbonyl compounds. The substrates include benzo-fused cyclic ketones which undergo reduction to furnish one stereoisomer.

SYNFACTS Contributors: Mark Lautens, Alvin Jang
 Synfacts 2016, 12(06), 0589 Published online: 17.05.2016
 DOI: 10.1055/s-0035-1562190; Reg-No.: L04416SF

Comment: Transfer hydrogenation (TH) reactions are milder and safer alternatives to metal-on-carbon-based hydrogenation reactions. Although the scope is usually limited to the reduction of carbonyls, TH avoids the use of highly pressurized systems. These reported ruthenium-based transfer hydrogenation catalysts are notable achievements in this field.

Category

Metal-Catalyzed
Asymmetric
Synthesis and
Stereoselective
Reactions

Key words

transfer
hydrogenation

ruthenium catalysis

asymmetric
reduction

Synfact
of the month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.