

The Next 10 Years in Voice Evaluation and Treatment

Julie M. Barkmeier-Kraemer, Ph.D.¹ and Rita R. Patel, Ph.D.²

ABSTRACT

Voice disorders are thought to affect approximately one third of all individuals within the United States during their lifetime. Individuals who require the use of their voice as part of their occupations are at highest risk for developing voice problems. Unfortunately, efficient diagnosis and effective management of voice disorders can be challenged by difficulty accessing professionals with the necessary expertise to diagnose and treat voice problems efficiently. Within the next decade, technological advancements show promise for improving the efficiency and effectiveness of intervention for voice disorders. Exciting developments in laryngeal imaging, modeling of patient-specific vocal patterns, and implementation of smart mobile technology and telehealth will greatly improve the accuracy of diagnosing voice problems and enhance implementation and carryover of effective voice treatment methods to daily communication demands.

KEYWORDS: Voice, voice disorders, evaluation, treatment, technology

Learning Outcomes: As a result of this activity, the reader will be able to (1) identify and discuss current directions in laryngeal imaging technology; (2) identify and discuss current directions in developing patient-specific methods for diagnosis and treatment; and (3) identify and discuss future methods for implementing mobile smart technology and telehealth technology to address current challenges in voice treatment.

Voice disorders affect nearly one-third of individuals in the United States age 20 to 66 years of age during their lifetime, with 39% of older people aged 45 to 64 years of

age reporting voice problems.¹ Of those diagnosed with voice disorders, two thirds are female.² Twenty-eight million people in the United States are estimated to require their

¹Division of Otolaryngology, University of Utah, Salt Lake City, Utah; ²Department of Speech and Hearing Sciences, Indiana University, Bloomington, Indiana.

Address for correspondence: Julie M. Barkmeier-Kraemer, Ph.D., Division of Otolaryngology, University of Utah, 50 N. Medical Dr 3C120, Salt Lake City, UT 84132 (e-mail: JulieB.Kraemer@hsc.utah.edu).

Forecasting the Future: Expert Opinions on the Next Decade for Adult Speech and Language Disorders; Guest Editor, Audrey L. Holland, Ph.D.

Semin Speech Lang 2016;37:158-165. Copyright © 2016 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA. Tel: +1(212) 584-4662.

DOI: <http://dx.doi.org/10.1055/s-0036-1583547>.

ISSN 0734-0478.

voice for job performance.³ Teachers represent one such voice-intensive occupation with 20% of teachers missing work due to voice problems.³ Approximately \$2.5 billion dollars are spent annually within the United States on teacher sick days and voice treatment expenses.³ Epidemiology research estimates that individuals with voice disorders miss an average of 39.2 work days/year/person due to a voice problem.⁴ One of the larger barriers to all individuals with voice problems is delayed referral to an appropriate health care provider. Delayed referral costs are estimated to be \$711 per person when the delay is greater than 3 months compared with \$271 per person when appropriately referred within the first 3 months of onset of a voice disorder.⁵

Over one million, or 6 to 9%, of all children in the United States also suffer from dysphonia.^{6,7} The proportion of school-aged children exhibiting a voice disorder ranges from 6 to 23%, depending upon the cohort. Studies have shown that voice disorders have an adverse impact on children academically as well as psychologically.⁸⁻¹⁰ Hence, early identification and treatment of voice disorders is critical.

By 2026, individuals with voice problems will likely enjoy a more accessible, efficient, and convenient method for detection and treatment of voice disorders. Evaluation and treatment of voice disorders will likely involve machine-based diagnostics using technological advances that map abnormal voicing patterns onto underlying pathophysiology (Fig. 1). Furthermore, voice therapy will be available in the comfort of the patient's home via telehealth or be implemented using smart technology during the course of daily life activities. Voice therapy approaches will be recommended for each individual using artificial intelligence capable of identifying optimal custom-based treatment methods from recorded voice use patterns. Optimal voice therapy outcomes require accurate implementation of specified voicing techniques and exercises that can be applied within typical communication activities. To ensure accurate implementation of therapeutic methods and goals, smart devices and phone apps will utilize patient-specific principles of behavior modification to cue individuals to complete treatment exercises at des-

ignated times during the day or use targeted voice therapy techniques during conversation when voice sensors determine aberrant performance. Thus, implementation of new technology can be used outside the clinic setting to facilitate improved implementation of therapeutic voice methods within the context of daily life activities. Technological advances by 2026 will revolutionize the efficiency and accuracy with which various voice disorders are diagnosed and effective voice therapy or medical/surgical treatments are recommended.

HIGH-SPEED VIDEO IMAGING

Comprehensive evaluation of voice is multidimensional; however, direct imaging of vocal fold vibratory patterns is currently fundamental for appropriate evaluation of the cause of the dysphonia. High-speed video (HSV) imaging, though not the current gold standard in laryngeal imaging, has transformed voice evaluations in state-of-the-art voice centers. Compared with the current gold standard of laryngeal imaging, videostroboscopy, current HSV imaging systems have the advantage of evaluating severely dysphonic voice qualities and rapid transient fluctuations in voice.¹¹⁻¹⁷ HSV imaging offers high temporal resolution of up to 8,000 frames per second compared with stroboscopy, which can only capture images at an average rate of 30 frames per second. Thus, stroboscopy frequently cannot accurately track the signal used to trigger stroboscopic lighting due to severely impaired vocal fold vibratory patterns. HSV imaging does not have this limitation because the high rate of image capture does not require synchronization with the voice signal. That is, valid HSV images without movement artifact can be obtained irrespective of the degree and duration of the voice disturbance during image recordings. Clinical indications for the use of HSV imaging support its value in evaluating severely affected voices,^{11,12} differentiating adductor spasmodic dysphonia from muscle tension dysphonia,¹⁷ comparing voice onset patterns during different types of singing,¹⁸ age-related voice changes,^{19,20} diplophonia,¹⁶ tremor,¹³ and various structural abnormalities affecting the vocal folds (e.g., sulcus vocalis, nodules and polyps).²¹⁻²⁴ HSV



Figure 1 The future of voice evaluations as imagined in on the cover of the Winter 2015 publication of The PHAROS as illustrated by Jim M'Guinness.

imaging also provides new insights into evaluating the effects of superficial dehydration on vibratory motion,²⁵ evaluating the growth and development of the vibratory motion in chil-

dren,^{26,27} monitoring changes in voice physiology following voice therapy,²⁸ and quantifying outcomes of phonosurgical treatment of early glottic carcinoma.²⁹ Furthermore, HSV

imaging combined with multiple arrays of lasers provides valuable insights into the basic mechanisms of phonation in adults and children by aiding in quantifying not only the two-dimensional vibratory motion,^{30–33} but also the three-dimensional vocal fold vibration kinematics in vivo, which is unavailable on standard laryngeal imaging.³⁴

The current limitation to clinical use of HSV imaging is the significant time required to complete visual-perceptual analysis of the long and detailed recordings. For example, 4 seconds of HSV recorded at 4,000 frames per second requires ~27 minutes to watch the entire recording at 10 frames per second using the playback rate. Moreover, visual-perceptual analysis of HSV is subjective and often results in low inter- and intra-rater reliability. However, recent work addressing these limitation using methods of machine learning, feature extraction, and automated analysis of HSV recordings have recently been shown to distinguish laryngeal pathologies with great accuracy.^{35–37} Thus, it is likely that HSV imaging with automated multifeature objective analysis will become the standard method by which clinical laryngeal imaging and analysis will be conducted in the evaluation of voice disorders during the upcoming decade.

DIAGNOSIS USING PATIENT-SPECIFIC VOICE MODELING

In addition to advances in laryngeal imaging, current efforts are underway to implement the use of bioinformatics and artificial intelligence for diagnosing and treating voice disorders. Artificial intelligence can detect patterns that the humans cannot perceive or predict and has the capacity to synthesize large amounts of data within seconds. Current voice modeling efforts study tissue-based biomarkers or voice physiology parameters to determine patient-specific patterns associated with each voice disorder and predict optimal custom-based treatment approaches for optimal outcomes.

One approach to simulating patient-specific information for treatment purposes is the use of biomarker agent-based models. Li and colleagues studied laryngeal secretion composition associated with onset of inflammatory

responses to phonotrauma and mapped recovery patterns in a small group of individuals.³⁸ Using the agent-based models approach, they determined which biomarkers from laryngeal secretions best predicted the recovery trajectory of speakers across a baseline, vocal loading (i.e., phonotrauma), and recovery period using three different paradigms of voice use. An extension of the initial model showed promise for developing simulation approaches based on laryngeal tissue secretion biomarkers that could eventually help clinicians determine optimal timing and treatment approaches for patients with laryngeal tissue injuries (e.g., post-laryngeal surgery, or trauma).³⁹

Another method to develop future applications for improved diagnostic accuracy entails the acquisition of physiologic parameters associated with normal and abnormal voice use for simulation followed by empirical testing. Simulation-based modeling is useful for testing the accuracy of predictions about voice output patterns associated with parameters of voice physiology. Such methods are being utilized to develop future diagnostic tools that will be able to detect the presence and characteristics of such voice disorders as vocal tremor,^{40,41} vocal fold paralysis,^{42–45} and functional voice abnormalities.^{46–49}

SMART MOBILE TECHNOLOGY

Another promising approach for evaluation and treatment purposes is the development of mobile technology for recording daily voicing patterns or cueing and facilitating the use of voice therapy techniques during daily life activities. The first form of these tools utilized attachment of an accelerometer to the skin overlying the larynx to record the frequency and intensity of the voice across an 8-hour period. Recordings are stored on a small computer device worn in a waist pack or in a pocket.^{50–53} Early work using these devices was used to compare patterns of voice use in teachers between work and nonwork environments.^{50,54,55} Voicing patterns of teachers compared with nonteachers elucidated the concept of “vocal dose” levels associated with onset of voice problems.⁵⁵ More recent work extended the information gained from the voice sensors to predict subglottal pressures,⁵⁶ aerodynamic

patterns,⁵⁷ and muscle activation patterns associated with normal and abnormal voicing.^{52,58,59} In the near future, these devices could be purchased by consumers as disposable sensors that communicate with smartphones to record voice use patterns and eventually provide biofeedback to the user regarding their accuracy in implementing targeted therapeutic voice use patterns in daily life.⁵³

Recent use of smartphones to facilitate accuracy and adherence to voice therapy methods also shows promise for future implementation of media that can be used by clinicians to ensure consistent and accurate patient practice of voice therapy methods.^{60,61} The use of voice therapy smartphone applications should lead to custom-based treatment recommendations such that patients are cued to practice at selected times during the day by their smartphone or mobile device based on patient-specific principles of behavior change to optimize acquisition and carryover of the target therapeutic skills. Partnered with physiologic feedback from the dosimeter or ambulatory phonation monitor, future treatment plans may enable direct cueing to patients to improve voice production techniques within their typical daily environment when aberrant physiology is detected by the sensor placed on their throat.

TELEHEALTH

Another advancement changing the face of voice therapy is the increased utilization of telemedicine/telehealth. Although literature on the efficacy and effectiveness of telehealth for voice treatment is not extensive, telehealth appears to be utilized to deliver voice therapy with comparable levels of treatment outcomes.⁶²⁻⁶⁵ The use of telehealth to deliver voice therapy will be particularly important for individuals with voice disorders living in rural regions without voice specialists or those with transportation barriers. Thus, implementation of telehealth and improvements in telehealth technology will enable improved access to voice treatment in the comfort of an individual's home or local site with necessary reception specifications (e.g., local library, medical center).

Current common barriers to use of telehealth for voice treatment include the lack of insurance

coverage by Medicare and other third-party payers as well as the need for the clinician to be licensed in the state where the patient resides. Although the latter is not a problem for in-state patients, regions where multiple state regions are served by one voice center make it necessary to maintain multiple state licenses and associated requirements. Hopefully, the ability of patients to access qualified providers for voice evaluation and treatment through telehealth will be addressed in upcoming efforts by the American Speech-Language-Hearing Association to initiate changes to current legislative and licensure barriers. We predict that implementation of telehealth for voice evaluation and treatment will improve accessibility to state-of-the-art clinical methods for improving diagnostic precision and treatment planning for those with voice disorders.

In summary, the area of voice and voice disorders faces an exciting time as technological advancements may significantly improve diagnostic precision and treatment planning through the development of improved imaging and computer diagnostic systems that will soon be available to communication science and disorder specialists.⁶⁶ With early detection of various conditions and information regarding risk factors at their fingertips, clinicians will be more involved in preventive care. With the aid of technological advances in artificial intelligence, communication science and disorder clinicians will be able to use computers to quickly determine evidenced-based diagnoses and highlight the risk factors, thereby facilitating early detection and treatment of various voice disorders. Finally, mobile technology methods will eventually be available to improve the efficiency with which voice treatment methods can be trained so that patients can quickly carryover improved voicing techniques and exercises into their daily lives. The increased availability and use of technology by voice patients to facilitate improved voicing methods in their daily lives will help reduce overall medical costs associated with inaccurate diagnoses and extended use of ineffective treatment methods. We predict that technological advances within the upcoming decade will have an impact on evaluating and treating voice disorders. This should result in significantly more efficient and effective approaches to voice

disorder management that is more accessible to the elderly and parents of children with voice disorders and to those living in rural regions than currently.

REFERENCES

1. Roy N, Merrill RM, Thibeault S, Gray SD, Smith EM. Voice disorders in teachers and the general population: effects on work performance, attendance, and future career choices. *J Speech Lang Hear Res* 2004;47(3):542–551
2. Cohen SM, Kim J, Roy N, Asche C, Courey M. Direct health care costs of laryngeal diseases and disorders. *Laryngoscope* 2012;122(7):1582–1588
3. Verdolini K, Ramig LO. Review: occupational risks for voice problems. *Logoped Phoniatr Vocol* 2001;26(1):37–46
4. Cohen SM, Kim J, Roy N, Asche C, Courey M. The impact of laryngeal disorders on work-related dysfunction. *Laryngoscope* 2012;122(7):1589–1594
5. Cohen SM, Kim J, Roy N, Courey M. Delayed otolaryngology referral for voice disorders increases health care costs. *Am J Med* 2015;128(4):426.e11–426.e18
6. Wilson DK. *Voice Problems in Children*. 3rd ed. Baltimore, MD/London, UK: Williams and Wilkins; 1987
7. McMurray JS. Disorders of phonation in children. *Pediatr Clin North Am* 2003;50(2):363–380
8. Ruddy BH, Sapienza CM. Treating voice disorders in the school-based setting: working within the framework of IDEA. *Lang Speech Hear Serv Sch* 2004;35(4):327–332
9. Connor NP, Cohen SB, Theis SM, Thibeault SL, Heatley DG, Bless DM. Attitudes of children with dysphonia. *J Voice* 2008;22(2):197–209
10. Lass NJ, Ruscello DM, Bradshaw KH, Blankenship BL. Adolescents' perceptions of normal and voice-disordered children. *J Commun Disord* 1991;24(4):267–274
11. Olthoff A, Woywod C, Kruse E. Stroboscopy versus high-speed laryngography: a comparative study. *Laryngoscope* 2007;117(6):1123–1126
12. Patel R, Dailey S, Bless D. Comparison of high-speed digital imaging with stroboscopy for laryngeal imaging of glottal disorders. *Ann Otol Rhinol Laryngol* 2008;117(6):413–424
13. Hertegård S, Larsson H, Wittenberg T. High-speed imaging: applications and development. *Logoped Phoniatr Vocol* 2003;28(3):133–139
14. Kunduk M, Yan Y, McWhorter AJ, Bless D. Investigation of voice initiation and voice offset characteristics with high-speed digital imaging. *Logoped Phoniatr Vocol* 2006;31(3):139–144
15. Hoppe U, Rosanowski F, Döllinger M, Lohscheller J, Schuster M, Eysholdt U. Glissando: laryngeal motorics and acoustics. *J Voice* 2003;17(3):370–376
16. Kiritani S, Hirose H, Imagawa H. High-speed digital image-analysis of vocal cord vibration in diplophonia. *Speech Commun* 1993;13(1–2):23–32
17. Patel RR, Liu L, Galatsanos N, Bless DM. Differential vibratory characteristics of adductor spasmodic dysphonia and muscle tension dysphonia on high-speed digital imaging. *Ann Otol Rhinol Laryngol* 2011;120(1):21–32
18. Freeman E, Woo P, Saxman JH, Murry T. A comparison of sung and spoken phonation onset gestures using high-speed digital imaging. *J Voice* 2012;26(2):226–238
19. Yamauchi A, Imagawa H, Yokonishi H, et al. Evaluation of vocal fold vibration with an assessment form for high-speed digital imaging: comparative study between healthy young and elderly subjects. *J Voice* 2012;26(6):742–750
20. Ahmad K, Yan Y, Bless D. Vocal fold vibratory characteristics of healthy geriatric females—analysis of high-speed digital images. *J Voice* 2012;26(6):751–759
21. Chodara AM, Krausert CR, Jiang JJ. Kymographic characterization of vibration in human vocal folds with nodules and polyps. *Laryngoscope* 2012;122(1):58–65
22. Yamauchi A, Yokonishi H, Imagawa H, et al. Visualization and estimation of vibratory disturbance in vocal fold scar using high-speed digital imaging. *J Voice* 2015. pii: S0892-1997(15)00151-4. doi: 10.1016/j.jvoice.2015.07.003. [Epub ahead of print]
23. Bohr C, Kräck A, Dubrovskiy D, et al. Spatiotemporal analysis of high-speed videolaryngoscopic imaging of organic pathologies in males. *J Speech Lang Hear Res* 2014;57(4):1148–1161
24. Bohr C, Kraeck A, Eysholdt U, Ziethe A, Döllinger M. Quantitative analysis of organic vocal fold pathologies in females by high-speed endoscopy. *Laryngoscope* 2013;123(7):1686–1693
25. Patel RR, Walker R, Sivasankar PM. Spatiotemporal quantification of vocal fold vibration following exposure to superficial laryngeal dehydration: a preliminary study. *J Voice* 2015. pii: S0892-1997(15)00157-5. doi: 10.1016/j.jvoice.2015.07.009. [Epub ahead of print]
26. Patel R, Donohue KD, Unnikrishnan H, Kryscio RJ. Kinematic measurements of the vocal-fold displacement waveform in typical children and adult populations: quantification of high-speed endoscopic videos. *J Speech Lang Hear Res* 2015;58(2):227–240
27. Patel RR, Dubrovskiy D, Döllinger M. Measurement of glottal cycle characteristics between children and adults: physiological variations. *J Voice* 2014;28(4):476–486

28. Patel RR, Pickering J, Stemple J, Donohue KD. A case report in changes in phonatory physiology following voice therapy: application of high-speed imaging. *J Voice* 2012;26(6):734–741
29. Mehta DD, Deliyiski DD, Zeitels SM, Quatieri TF, Hillman RE. Voice production mechanisms following phonosurgical treatment of early glottic cancer. *Ann Otol Rhinol Laryngol* 2010;119(1):1–9
30. Jiang JJ, Chang CI, Raviv JR, Gupta S, Banzali FM Jr, Hanson DG. Quantitative study of mucosal wave via videokymography in canine larynges. *Laryngoscope* 2000;110(9):1567–1573
31. Schubert S, Hoppe U, Döllinger M, Lohscheller J, Eysholdt U. High-precision measurement of the vocal fold length and vibratory amplitudes. *Laryngoscope* 2002;112(6):1043–1049
32. Patel RR, Donohue KD, Johnson WC, Archer SM. Laser projection imaging for measurement of pediatric voice. *Laryngoscope* 2011;121(11):2411–2417
33. Patel RR, Donohue KD, Lau D, Unnikrishnan H. In vivo measurement of pediatric vocal fold motion using structured light laser projection. *J Voice* 2013;27(4):463–472
34. Semmler M, Kniesburges S, Birk V, Ziethe A, Patel R, Doellinger M. 3D reconstruction of human laryngeal dynamics based on endoscopic high-speed recordings. *IEEE Trans Med Imaging* 2016 [Epub ahead of print]
35. Panek D, Skalski A, Zielinski T, Deliyiski DD. Voice pathology classification based on high-speed videoendoscopy. *Conf Proc IEEE Eng Med Biol Soc* 2015;2015:735–738
36. Voigt D, Döllinger M, Yang A, Eysholdt U, Lohscheller J. Automatic diagnosis of vocal fold paresis by employing phonovibrograms features and machine learning methods. *Comput Methods Programs Biomed* 2010;99(3):275–288
37. Voigt D, Döllinger M, Braunschweig T, Yang A, Eysholdt U, Lohscheller J. Classification of functional voice disorders based on phonovibrograms. *Artif Intell Med* 2010;49(1):51–59
38. Li NY, Verdolini K, Clermont G, et al. A patient-specific in silico model of inflammation and healing tested in acute vocal fold injury. *PLoS ONE* 2008;3(7):e2789
39. Li NY, Vodovotz Y, Kim KH, Mi Q, Hebda PA, Abbott KV. Biosimulation of acute phonotrauma: an extended model. *Laryngoscope* 2011;121(11):2418–2428
40. Lester RA, Barkmeier-Kraemer J, Story BH. Physiologic and acoustic patterns of essential vocal tremor. *J Voice* 2013;27(4):422–432
41. Lester RA, Story BH. The effects of physiological adjustments on the perceptual and acoustical characteristics of simulated laryngeal vocal tremor. *J Acoust Soc Am* 2015;138(2):953–963
42. Chhetri DK, Neubauer J, Bergeron JL, Sofer E, Peng KA, Jamal N. Effects of asymmetric superior laryngeal nerve stimulation on glottic posture, acoustics, vibration. *Laryngoscope* 2013;123(12):3110–3116
43. Chhetri DK, Neubauer J, Berry DA. Neuromuscular control of fundamental frequency and glottal posture at phonation onset. *J Acoust Soc Am* 2012;131(2):1401–1412
44. Erath BD, Peterson SD, Zañartu M, Wodicka GR, Plesniak MW. A theoretical model of the pressure field arising from asymmetric intraglottal flows applied to a two-mass model of the vocal folds. *J Acoust Soc Am* 2011;130(1):389–403
45. Khosla S, Oren L, Gutmark E. An example of the role of basic science research to inform the treatment of unilateral vocal fold paralysis. *SIG 3 Perspectives on Voice and Voice Disorders. ASHA* 2014;24:37–50
46. Yang A, Berry DA, Kaltenbacher M, Döllinger M. Three-dimensional biomechanical properties of human vocal folds: parameter optimization of a numerical model to match in vitro dynamics. *J Acoust Soc Am* 2012;131(2):1378–1390
47. Yang A, Stingl M, Berry DA, et al. Computation of physiological human vocal fold parameters by mathematical optimization of a biomechanical model. *J Acoust Soc Am* 2011;130(2):948–964
48. Chhetri DK, Neubauer J, Sofer E, Berry DA. Influence and interactions of laryngeal adductors and cricothyroid muscles on fundamental frequency and glottal posture control. *J Acoust Soc Am* 2014;135(4):2052–2064
49. Stepp CE, Hillman RE, Heaton JT. A virtual trajectory model predicts differences in vocal fold kinematics in individuals with vocal hyperfunction. *J Acoust Soc Am* 2010;127(5):3166–3176
50. Popolo PS, Svec JG, Titze IR. Adaptation of a Pocket PC for use as a wearable voice dosimeter. *J Speech Lang Hear Res* 2005;48(4):780–791
51. Mehta DD, Van Stan JH, Zañartu M, et al. Using ambulatory voice monitoring to investigate common voice disorders: research update. *Front Bioeng Biotechnol* 2015;3(3):155
52. Ghassemi M, Van Stan JH, Mehta DD, et al. Learning to detect vocal hyperfunction from ambulatory neck-surface acceleration features: initial results for vocal fold nodules. *IEEE Trans Biomed Eng* 2014;61(6):1668–1675
53. Mehta DD, Zañartu M, Feng SW, Cheyne HA II, Hillman RE. Mobile voice health monitoring using a wearable accelerometer sensor and a smartphone platform. *IEEE Trans Biomed Eng* 2012;59(11):3090–3096
54. Halpern AE, Spielman JL, Hunter EJ, Titze IR. The inability to produce soft voice (IPSV): a tool to detect vocal change in school-teachers. *Logoped Phoniatr Vocol* 2009;34(3):117–127

55. Titze IR, Hunter EJ. Comparison of vocal vibration-dose measures for potential-damage risk criteria. *J Speech Lang Hear Res* 2015;58(5):1425–1439
56. Zañartu M, Ho JC, Mehta DD, Hillman RE, Wodicka GR. Subglottal impedance-based inverse filtering of voiced sounds using neck surface acceleration. *IEEE Trans Audio Speech Lang Process* 2013;21(9):1929–1939
57. Llico AF, Zañartu M, González AJ, et al. Real-time estimation of aerodynamic features for ambulatory voice biofeedback. *J Acoust Soc Am* 2015;138(1):EL14–EL19
58. Smith N, Rivera L, Dietrich M, Shyu CR, Page M, DeSouza G. Detection of simulated vocal dysfunctions using complex sEMG patterns. *IEEE J Biomed Health Inform* 2015
59. Stepp CE, Heaton JT, Jetté ME, Burns JA, Hillman RE. Neck surface electromyography as a measure of vocal hyperfunction before and after injection laryngoplasty. *Ann Otol Rhinol Laryngol* 2010;119(9):594–601
60. van Leer E, Connor NP. Use of portable digital media players increases patient motivation and practice in voice therapy. *J Voice* 2012;26(4):447–453
61. van Leer E, Connor NP. Predicting and influencing voice therapy adherence using social-cognitive factors and mobile video. *Am J Speech Lang Pathol* 2015;24(2):164–176
62. Mashima PA, Brown JE. Remote management of voice and swallowing disorders. *Otolaryngol Clin North Am* 2011;44(6):1305–1316, viii
63. Mashima PA, Birkmire-Peters DP, Syms MJ, Holtel MR, Burgess LP, Peters LJ. Telehealth: voice therapy using telecommunications technology. *Am J Speech Lang Pathol* 2003;12(4):432–439
64. Kelchner L. Telehealth and the treatment of voice disorders: a discussion regarding evidence. *SIG 3 Perspectives on Voice and Voice Disorders* 2013;23:88–94
65. Howell S, Tripoliti E, Pring T. Delivering the Lee Silverman Voice Treatment (LSVT) by web camera: a feasibility study. *Int J Lang Commun Disord* 2009;44(3):287–300
66. Bond WF, Schwartz LM, Weaver KR, Levick D, Giuliano M, Graber ML. Differential diagnosis generators: an evaluation of currently available computer programs. *J Gen Intern Med* 2012;27(2):213–219