Editorial

Long-Acting Reversible Contraceptives: An Important Approach to Reduce Unintended Pregnancies

Contraceptivos reversíveis de longa duração: uma importante medida para reduzir as gestações não planejadas

Carolina Sales Vieira¹

¹ Department of Gynecology and Obstetrics, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo – USP, Ribeirão Preto, SP, Brazil

Rev Bras Ginecol Obstet 2016;38:207–209.

Unintended pregnancies are a major public health concern worldwide. In Brazil, as much as 55% of pregnancies are unplanned.¹ Unplanned pregnancy is associated with an increased risk of maternal, neonatal and infant morbidity and mortality.² It also reduces educational and career opportunities for mothers, and it may contribute to socioeconomic deprivation and widening income disparities.³ The total cost attributed to unplanned pregnancy in Brazil is estimated to be R\$ 4.1 billion (roughly US\$ 1.85 billion) annually.⁴

Part of the high rate of unplanned pregnancies may be due to the relatively low use of long-acting reversible contraceptives (LARCs), specifically contraceptive implants and intrauterine devices (IUDs).⁵ Less than 2% of Brazilian women who take contraceptives use LARCs.⁶ By comparison, in the UK, LARCs are used by 31% of women using contraceptives, and the rate of unplanned pregnancies there is estimated to be 16.2%.⁷ There are many reasons for the low prevalence rate of women using LARCs in Brazil, some of which are: only one type of LARC is available for free in the public health system (copper IUD); a lack of training in LARC methods in the majority of the obstetrics and gynecology residence programs; and biased information and inadequate counseling on LARCs being offered by some healthcare providers.

The CHOICE project was responsible for bringing LARCs and unplanned pregnancies to the spotlight of the reproductive planning discussion. The CHOICE project was an observational cohort study developed to promote the use of LARC methods in the St. Louis region (USA). It was designed to investigate if high and stagnant rates of unintended pregnancy could be reduced by increasing the uptake of LARC

methods. In order to achieve this objective, the project removed two major barriers in the use of LARC methods: the lack of access to free LARCs and the lack of adequate information on the safety and efficacy of these methods.⁸ When the barriers of cost, access and knowledge were removed, 75% of the CHOICE cohort chose a LARC method at baseline enrollment.⁹ The continuation rates of LARC methods were higher than those of non-LARC methods at 12 and 24 months (86 against 55% at 12 months; 77 against 41% at 24 months).^{9,10} Overall, 84% of LARC users were satisfied with the method at 12 months, while only 53% of participants using short-acting methods were satisfied at the same period.⁹ Although there is a concern that an increased uptake of LARC methods could increase risk-taking sexual behavior, the CHOICE project showed that the provision of no-cost contraception was not associated with increased risk-taking sexual behaviors.¹¹ Additionally, the superiority of LARC methods was confirmed over short-acting methods; implants and IUDs were 22 times more effective than oral contraceptive pills, patches, or rings.¹² In order to evaluate the population impact of this huge increase of LARCs use in the St. Louis region, the average annual rates of teen pregnancy, birth and induced abortion among the CHOICE participants were compared with the national rates of these outcomes. When compared with the national data, the CHOICE project showed over 75% reduction in all three outcomes.¹³

In Brazil, studies using LARCs also showed low rates of premature discontinuations, and high continuation and satisfaction rates when adequate counseling was provided.^{14,15} Therefore, counseling and evidence-based information are crucial to facilitate the decisions of women regarding a

Address for correspondence
Carolina Sales Vieira, Avenida
Bandeirantes, 3900 - Campus
Universitário - 14049-900,
Ribeirão Preto, SP, Brazil
(e-mail: carol.sales@uol.com.br).

DOI <http://dx.doi.org/10.1055/s-0036-1583761>.
ISSN 0100-7203.

Copyright © 2016 by Thieme Publicações
Ltda, Rio de Janeiro, Brazil

License terms

contraceptive method.¹⁴ In the family planning clinic of the University of Campinas (Brazil), where there is free access to LARCs, in the past 15 years the rates of women who opted to continue using LARC methods until menopause were higher than of those who opted for female or male sterilization. The annual number of sterilizations dropped markedly in the same period, as we can observe in this issue of RBGO.¹⁶

In addition to a positive impact on women's health outcomes, increasing LARC uptake has also a favorable economic impact.¹⁷ For example, it was estimated that if 10% of women aged 20–29 years in the US switched from oral contraception to LARCs, the total cost of unplanned pregnancies would be reduced by US\$ 288 million per year.¹⁸ With savings of more than US\$7 for each US\$1 spent, LARC methods were shown to be more cost-effective than short-acting methods or no method.¹⁹

In vulnerable populations, such as adolescents or drug users, the use of LARCs can have even a more prominent effect on women's health outcomes and on cost-effectiveness than short-acting methods. Women aged less than 21 years using short-acting methods were twice more likely to experience an unintended pregnancy than older women using the same method. Among LARC users, there was no difference in the risk of unplanned pregnancy across age groups.¹² In Brazil, unintended pregnancy is a major problem among crack cocaine users. Of 45,600 Brazilian children who lived in public foster care in 2013, 81% were from parents who are addicted to drugs, and while the majority of these children have a family, as many as 77% of them do not receive any visits.²⁰ The number of pregnancies per women (3.4) among Brazilian crack cocaine users is almost double the national rate.²¹ These women also have a higher rate of syphilis (20.4 against 1.6%), HIV (8.2 against 0.4%), and hepatitis C (2.2 against 1.38%) when compared with reported national rates.^{21,22} The increased rate of sexual infectious diseases and obstetrics morbidity (prematurity and fetal demise) in this population increases the social and economic costs of unplanned pregnancies.²¹ A Brazilian study estimated that with the use of an etonogestrel-releasing implant by 101 female crack cocaine users, the public health system could potentially save R\$ 341,643.50 (roughly US\$ 94,980), considering only the costs of the pregnancies.²¹

The use of LARCs is also important during the postpartum period,^{23–25} which offers a window of opportunity for contraceptive counseling and initiation. This approach reduces rapid repeat pregnancy by over 80%, especially in vulnerable populations like adolescents.^{24,25}

Considering all advantages of LARCs, many health organizations, non-governmental organizations and medical societies are recommending adequate counseling on LARC methods and improvement of access to LARCs to all candidates, including nulliparous women and adolescents.^{5,26,27} The World Health Organization (WHO) also included all LARC methods in the list of essential medicines for a basic healthcare system. In this list is included the most effective, safe and cost-effective medicines for priority conditions.²⁸ Despite these recommendations and a request from the Brazilian Federation of Gynecology and Obstetrics Associations (Febrasgo), the Brazilian government refuses to include the etonogestrel-releasing implant and the levonorgestrel-releasing intrauterine system in the national list of essential medicines. The Brazilian government said that these methods do not present advantages over the contraceptive methods currently offered by the public health system, and that goes against current scientific evidence.^{5,9,10,12,13,17–19,26,27,29}

In order to empower women when it comes to family planning, we must offer counseling and access to all methods of contraception, enabling women to make informed decisions about whether and when to have children. LARCs are the most cost-effective contraceptive methods; they have the highest efficacy and continuation rates among all contraceptives, and show the most prominent effect on reducing unplanned pregnancy and abortion rates. Increasing the use of highly effective contraceptive methods may be part of the solution to decrease the persistent high rate of unintended pregnancies. In order to increase LARC uptake, it is important to promote free access and information on these methods for women and adolescents, and to educate their partners, providers, and policymakers about the potential usefulness of LARCs. Finally, it is essential to implement family planning policies based on cost-effectiveness and the best evidence available.

References

- Viellas EF, Domingues RM, Dias MA, et al. Prenatal care in Brazil. *Cad Saude Publica* 2014;30(Suppl 1):S1–S15
- Singh A, Singh A, Mahapatra B. The consequences of unintended pregnancy for maternal and child health in rural India: evidence from prospective data. *Matern Child Health J* 2013;17(3):493–500
- Parks C, Peipert JF. Eliminating health disparities in unintended pregnancy with long-acting reversible contraception (LARC). *Am J Obstet Gynecol* 2016 Feb 12. Ahead of print. doi: 10.1016/j.ajog.2016.02.017
- Le HH, Connolly MP, Bahamondes L, Cecatti JG, Yu J, Hu HX. The burden of unintended pregnancies in Brazil: a social and public health system cost analysis. *Int J Womens Health* 2014;6:663–670
- Committee on Gynecologic Practice Long-Acting Reversible Contraception Working Group. Committee Opinion No. 642: increasing access to contraceptive implants and intrauterine devices to reduce unintended pregnancy. *Obstet Gynecol* 2015;126(4):e44–e48
- Brasil. Ministério da Saúde [Internet]. Pesquisa Nacional de Demografia e Saúde da Criança e da Mulher - PNDS 2006. Brasília (DF)Ministério da Saúde2009 [citado 2016 Fev 10]. Disponível em: http://bvsms.saude.gov.br/bvs/publicacoes/pnds_crianca_mulher.pdf
- Wellings K, Jones KG, Mercer CH, et al. The prevalence of unplanned pregnancy and associated factors in Britain: findings from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3). *Lancet* 2013;382(9907):1807–1816
- Secura GM, Allsworth JE, Madden T, Mullersman JL, Peipert JF. The Contraceptive CHOICE Project: reducing barriers to long-acting reversible contraception. *Am J Obstet Gynecol* 2010;203(2):115.e1–115.e7
- Peipert JF, Zhao Q, Allsworth JE, et al. Continuation and satisfaction of reversible contraception. *Obstet Gynecol* 2011;117(5):1105–1113

- 10 O'Neil-Callahan M, Peipert JF, Zhao Q, Madden T, Secura G. Twenty-four-month continuation of reversible contraception. *Obstet Gynecol* 2013;122(5):1083-1091
- 11 Secura GM, Adams T, Buckel CM, Zhao Q, Peipert JF. Change in sexual behavior with provision of no-cost contraception. *Obstet Gynecol* 2014;123(4):771-776
- 12 Winner B, Peipert JF, Zhao Q, et al. Effectiveness of long-acting reversible contraception. *N Engl J Med* 2012;366(21):1998-2007
- 13 Secura GM, Madden T, McNicholas C, et al. Provision of no-cost, long-acting contraception and teenage pregnancy. *N Engl J Med* 2014;371(14):1316-1323
- 14 Modesto W, Bahamondes MV, Bahamondes L. A randomized clinical trial of the effect of intensive versus non-intensive counselling on discontinuation rates due to bleeding disturbances of three long-acting reversible contraceptives. *Hum Reprod* 2014;29(7):1393-1399
- 15 Guazzelli CA, de Queiroz FT, Barbieri M, Torloni MR, de Araujo FF. Etonogestrel implant in postpartum adolescents: bleeding pattern, efficacy and discontinuation rate. *Contraception* 2010;82(3):256-259
- 16 Ferreira JM, Monteiro I, Castro S, Villarroel M, Silveira C, Bahamondes L. The use of long acting reversible contraceptives and the relationship between discontinuation rates due to menopause and to female and male sterilizations. *Rev Bras Ginecol Obstet* 2016;38(5):210-217
- 17 Mavranezouli I; LARC Guideline Development Group. The cost-effectiveness of long-acting reversible contraceptive methods in the UK: analysis based on a decision-analytic model developed for a National Institute for Health and Clinical Excellence (NICE) clinical practice guideline. *Hum Reprod* 2008;23(6):1338-1345
- 18 Trussell J, Henry N, Hassan F, Prezioso A, Law A, Filonenko A. Burden of unintended pregnancy in the United States: potential savings with increased use of long-acting reversible contraception. *Contraception* 2013;87(2):154-161
- 19 Foster DG, Rostovtseva DP, Brindis CD, Biggs MA, Hulett D, Darney PD. Cost savings from the provision of specific methods of contraception in a publicly funded program. *Am J Public Health* 2009;99(3):446-451
- 20 Conselho Nacional do Ministério Público. (CNMP) [Internet]. CNMP divulga dados sobre acolhimento de crianças e adolescentes. 2013 [citado 2015 Dez 12]. Disponível em: http://www.cnmp.mp.br/portal_2015/todas-as-noticias/3702-cnmp-divulga-dados-sobre-acolhimento-de-criancas-e-adolescentes
- 21 Sakamoto LC, Malavasia AL, Karasin AL, Frajzinger RC, Araújo MR, Gebirim LH. Prevenção de gestações não planejadas com implante subdérmico em mulheres da Cracolândia, São Paulo. *Reprod Clim*. 2015;30(3):102-107
- 22 Bastos FI, Bertoni N. Pesquisa Nacional sobre o uso de crack: quem são os usuários de crack e/ou similares do Brasil? Quantos são nas capitais brasileiras? [Internet]. Rio de Janeiro Editora ICICT/FIO-CRUZ 2014 [citado 2016 Fev 20]. Disponível em: <https://www.iciet.fiocruz.br/sites/www.iciet.fiocruz.br/files/Pesquisa%20Nacional%20sobre%20o%20Uso%20de%20Crack.pdf>
- 23 Brito MB, Ferriani RA, Quintana SM, Yazlle ME, Silva de Sá MF, Vieira CS. Safety of the etonogestrel-releasing implant during the immediate postpartum period: a pilot study. *Contraception* 2009;80(6):519-526
- 24 Tocce KM, Sheeder JL, Teal SB. Rapid repeat pregnancy in adolescents: do immediate postpartum contraceptive implants make a difference? *Am J Obstet Gynecol* 2012;206(6):481.e1-481.e7
- 25 Cohen R, Sheeder J, Arango N, Teal SB, Tocce K. Twelve-month contraceptive continuation and repeat pregnancy among young mothers choosing postdelivery contraceptive implants or postplacental intrauterine devices. *Contraception* 2016;93(2):178-183
- 26 Committee on Adolescence. Contraception for adolescents. *Pediatrics* 2014;134(4):e1244-e1256
- 27 Population Council. International Federation of Gynecology and Obstetrics (FIGO). Reproductive Health Supplies Coalition [Internet]. 2013 Statement from the Bellagio Group on LARCs: long-acting reversible contraception in the context of full access, full choice. 2013 [cited 2015 Dec 18]. Available from: www.popcouncil.org/pdfs/2013RH_BellagioConsensus.pdf
- 28 World Health Organization [Internet]. 19th WHO model list of essential medicines. 2015 [cited 2016 Fev 8]. Available from: http://www.who.int/medicines/publications/essentialmedicines/EML2015_8-May-15.pdf?ua=1
- 29 Bahamondes L, Brache V, Meirik O, Ali M, Habib N, Landoulsi S; WHO Study Group on Contraceptive Implants for Women. A 3-year multicentre randomized controlled trial of etonogestrel- and levonorgestrel-releasing contraceptive implants, with non-randomized matched copper-intrauterine device controls. *Hum Reprod* 2015;30(11):2527-2538