Influence of Birthweight on the Prospective Stillbirth Risk in the Third Trimester: A Cross-Sectional Cohort Study

Stephen Contag, MD¹ Clayton Brown, MD, PhD² Sarah Crimmins, MD¹ Katherine Goetzinger, MD¹

¹ Department of Obstetrics, Gynecology and Reproductive Sciences, University of Maryland School of Medicine, Baltimore, Maryland

² Department of Epidemiology and Public Health, University of Maryland, Baltimore, Maryland

Address for correspondence Stephen Contag, MD, Department of Obstetrics, Gynecology and Reproductive Sciences, University of Maryland School of Medicine, Suite N6W104G, 22 South Greene Street, Baltimore, MD 20201 (e-mail: scontag@fpi.umaryland.edu).

Am J Perinatol Rep 2016;6:e287–e298.

Abstract

Objective The objective of this study was to determine the effect of birthweight on prospective stillbirth risk.

Methods Cross-sectional study of singleton births in the United States from 2010 to 2012 from 32 through 42 weeks was conducted. Stillbirth risk was stratified by birthweight and gestational age adjusted for time from death to delivery. The primary outcome was the prospective stillbirth risk for each birthweight category. Student *t*-test was used for continuous data, chi-square to compare categorical data. Binomial proportions were used to derive prospective and cumulative risks. Cox proportional hazards regression with log-rank test comparison for heterogeneity was used to compare birthweight categories and derive hazard ratios.

Results There was an increase in the risk for stillbirth as birthweight diverged from the reference group. At 40 weeks adjusted gestational age, stillbirth rate per 10,000 births for the bottom (6.17, 95% CI: 7.47–4.87) and top (2.37, 95% CI: 3.1–1.65) 5th centiles of birthweight conveyed the highest risk. Hazard ratios (HR) after adjusting for covariates were: 1.55 (1.73–1.4) <5th centile and 2.2 (2.43–1.99) > 95th centile (*p* < 0.001).

Conclusion Stillbirth risk increases as birthweight departs from the mean. Birthweight below the 5th and above the 95th centile conveyed a significantly increased risk for stillbirth which was most noticeable after 37 weeks.

Keywords

- ▶ third trimester
- ▶ birthweight
- ▶ stillbirth
- ▶ growth restriction
- ▶ prospective risk

Stillbirth rates are a sensitive indicator of quality in a country's healthcare system.¹ There are relatively limited data regarding its significance and prevention.^{1,2} In the United States, the stillbirth rate is ~3 per 1,000. The rate has decreased by 30% among preterm births between 1989 and 2000.³ This decline was partly caused by a 55% increase in medically indicated preterm births during the same period.³ Subsequently, from 2005 through 2012, medically indicated preterm birth declined by 17% indicating improvements in the identification of fetuses at risk for adverse outcomes.⁴ Aggressive management of conditions such as Rh alloimmu-

nization, diabetes, preeclampsia, and fetal growth restriction (FGR) has also contributed to this decrease.^{5–9}

Although prior case-control and cohort studies have demonstrated that low weight is a risk factor for stillbirth,^{10,11} several studies have shown that both the upper and lower extremes of birthweight are associated with an increased risk for stillbirth.^{12,13} Several reports have led to recommendations for heightened fetal surveillance of small fetuses with late preterm or early term delivery to prevent stillbirth,^{14–19} but there is no clear guidance for those in the highest birthweight centiles.

received
March 30, 2016
accepted after revision
July 1, 2016

DOI <http://dx.doi.org/10.1055/s-0036-1587322>.
ISSN 2157-6998.

Copyright © 2016 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.
Tel: +1(212) 584-4662.

License terms

Stillbirth rate is defined as the number of stillbirths in a specific week of gestation per all births in the same week of gestation. The prospective stillbirth risk is defined as the number of stillbirths in a specific week of gestation per all ongoing pregnancies in that week of gestation,²⁰ and the cumulative risk of stillbirth is the sum of the prospective stillbirth risks, and it increases with advancing gestational age. It depicts how the additive effect of prospective risk of stillbirth increases over time.^{21,22}

We sought to quantify the effect of birthweight on the prospective risk of stillbirth in the US population. Although birthweight below the 10th centile is the most commonly used definition below which the risk increases,^{23–25} low birthweight is not a dichotomous condition. We hypothesize that the risk for stillbirth increases gradually as birthweight centile decreases. The objective of this study was to quantify the risk of stillbirth within birthweight decile categories ranging from the lowest to the highest deciles of birthweight.

Methods

This is a population-based, cross-sectional analysis of singleton live births in the United States from the years 2010 through 2012. Data were accessed from the National Center for Health Statistics (NCHS).²⁶ The NCHS compiles information from the US birth data files and fetal death data files, which are required for all births. The validity of the data has been documented.²⁷ This is de-identified and publicly available information and was exempt from Institutional Review Board review.

We used the obstetric estimate of gestation which provides better estimates of gestational age than does the last menstrual period LMP.²⁸ The stillbirth certificate reports gestational age based on time of delivery and would be expected to overestimate gestational age. The gestational age for all stillbirths was randomly adjusted downwards by 0 to 4 weeks in the proportions equal to those reported for a large US stillbirth cohort.²⁹ The adjustment was performed prior to any exclusions. We performed all analyses with and without adjustment for gestational age.

We included only singleton births from women residing and delivering in the United States between 2010 and 2012. The analysis was performed at each week of gestation from 32 to 42 weeks because this is when most fetal surveillance algorithms are implemented. Exclusion criteria were gestational age less than 32 or more than 42 weeks, twins or higher order multiples, and all fetuses with congenital anomalies or known aneuploidy, which have an increased risk for stillbirth independent of growth restriction or gestational age.

The reference curves were established using birthweights from the population of livebirths in 2012.^{30–32} The mean birthweight and standard deviation for each week of gestation were used to obtain a centile for each birthweight. Birthweight centiles were stratified into 12 groups: <3rd, < 5th, 5th to 10th, 11th to 20th, 21st to 30th, 31st to 40th, 41st to 60th, 61st to 70th, 71st to 80th, 81st to 90th, 91st to 95th and > 95th centile.

We compared the descriptive data between stillborn and live-born infants. Continuous data were compared with the Student *t*-test and categorical data were compared with a chi-square test. *P* values are all two sided with $p < 0.05$ considered to be significant. Binomial proportions for prospective and cumulative stillbirth risks were obtained.²¹ The prospective risk of stillbirth is considered to be a better indicator of the population at risk than the stillbirth rate.^{33,34} An adjustment that excluded half of the deliveries occurring in the week analyzed was made to account for births that occurred during that week and would no longer have been at risk for death.²¹ For cumulative risk for stillbirth, first we calculated the cumulative probability of survival as the product of the prospective probabilities of survival. The cumulative probability of stillbirth was 1 – probability of survival.²¹ We plotted the cumulative probability of stillbirth with deliveries censored at each week of gestation as Kaplan Meier failure curves.³⁵ Risks were stratified by birthweight category and reported per 10,000 births.

The effect of demographic and clinical variables on risk of stillbirth was analyzed in a multivariate model using a Cox proportional hazards regression with log-rank test comparison for heterogeneity. Hazard ratios are the best descriptor for risk of events over a specific interval of time.³⁶ The reference group for the relative risk were infants born in the 41–60% birthweight category. For multiple analyses between birthweight categories, a Bonferroni correction was performed with *p* values being significant when <0.004 . Variables associated with both growth restriction and stillbirth, and which were available in both the live birth and stillbirth certificates, were included in the multivariate analyses.^{37,38} These were live birth order, maternal age, weight gain, and race.^{20,38} The variables maternal smoking, body mass index, the results of antepartum surveillance, and obstetrical complications were not available in the certificate of stillbirth and were not included. The unreported values were weight gain (1,036,078), delivery method (2,058,324), race (239,201), and maternal morbidities (1,868,566). These variables were not part of the primary outcome analysis were considered missing at random and no imputations were performed.³⁹ Values reported as unknown or blank were considered to be incomplete and were excluded.⁴⁰ All data analysis was performed using SAS 9.3 (Cary, NC).

Results

After exclusions, there were a total of 11,304,635 live births and 23,519 stillbirths available for analysis from 2010 through 2012 (►Fig. 1). The characteristics of the entire cohort are described comparing the stillborn and live born infants (►Table 1). Maternal age, number of prior live births, weight gain, age, race, maternal morbidities, infant weight, gender, and mode of delivery were significantly different between live and stillbirths. The total number of ongoing pregnancies, live births, stillbirths in each birthweight category is reported by adjusted gestational age in ►Table 1.

The prospective and cumulative risk of stillbirth for the entire cohort, according to clinical estimate and adjusted

Fig. 1 Flowchart demonstrating all women included in the study cohort according to outcome and exclusion category.

gestational age, are shown in ►**Fig. 2**. The prospective and cumulative stillbirth risk gradually increased up to 39 weeks, with an upwards inflection of the slope at 39 weeks among the unadjusted gestational age values. The risks after adjusting for gestational age at death demonstrated a lower risk up to 41 weeks with an inflection after 41 weeks which was observed for both adjusted and unadjusted values.

The binomial proportions for the prospective and cumulative risk of stillbirth by birthweight category were derived using the adjusted gestational age (►**Table 2**). The lowest proportions are seen in the 31st to 40th centile category, which was similar to that reported for the reference group. The highest risk was observed for birthweights below the 5th and above the 95th centiles. The cumulative stillbirth risk, which reflects the total accumulated risk, for birthweights below the 3rd centile ranged from 2.61 per 10,000 at 32 weeks to 73.33 per 10,000 at 42 weeks. For birthweights below the 5th centile, it was 2.34 per 10,000 at 32 weeks and 63.15 per 10,000 at 42 weeks. In comparison, for stillbirths with birthweights above the 95th centile, the risk was 4.03 per 10,000 at 32 weeks and 37.01 per 10,000 at 42 weeks, which was higher than in all other categories except for below the 5th centile (►**Table 2**).

Kaplan–Meier failure curves were used to plot the cumulative risk for fetal death according to birthweight categories.

The frequencies were similar to those obtained by calculating the binomial proportions (►**Figs. 3A** and **3B**). Using the clinical estimate of gestational age used in the certificate of fetal death, events with birthweights below the 5th centile had higher cumulative stillbirth risk compared with those above the 5th centile (Log rank $p < 0.001$). The lowest cumulative stillbirth risks were observed for those born between the 71st and the 95th centile. These were lower than the cumulative stillbirth risk for the reference group (41st to 60th centile). Birthweights above the 95th centile were associated with cumulative stillbirth risks similar to those of the reference group (►**Fig. 3A**).

After adjusting the gestational age for time from death to delivery, we observed a smoother transition in cumulative stillbirth risks between the birthweight categories. The lowest risk was for those in both the reference and 31st to 40th centile categories. The risk for those above the 95th centile was similar to that observed for those below the 5th centile, all of which were higher than the rest of the cohort (Log-rank test $p < 0.001$). This divergence was particularly noticeable after 37 weeks of gestation (►**Fig. 3B**).

The hazard ratio for stillbirth in each birthweight category was compared with the reference group of birthweights between the 41st and the 60th centiles. The analysis was performed using both the clinical estimate of gestational age

Table 1 Characteristics of cohort by pregnancy outcome ($n = 11,328,154$)

Characteristics	Live birth (11,304,635)	Stillbirth (23,519)
Maternal		
Age, mean(SD)	27.8 (6.0)	28.2 (6.5)
Age category, y, n (%)		
Younger than 25	3,658,180 (32.4)	7,513 (31.9)
25–34	6,024,789 (53.3)	11,659 (49.6)
Older than 35	1,621,666 (14.4)	4,347 (18.5)
Previous live births, n (%)		
0	4,566,385 (40.4)	14,026 (59.6)
1–2	5,383,399 (47.6)	6,035 (25.7)
3–4	1,058,357 (9.4)	1,522 (6.5)
> 4	296,494 (2.6)	1,936 (8.2)
Weight gain in kg, n (%)		
Less than 5	893,990 (8.7)	3,533 (15.9)
5–14	4,904,098 (47.8)	9,702 (43.7)
15–24	3,433,494 (33.4)	4,130 (18.6)
25–34	432,063 (4.2)	524 (2.4)
Greater than 34	606,235 (5.9)	4,307 (19.4)
Race–ethnicity, n (%)		
White non-Latino	6,112,340 (55.2)	10,989 (49.5)
Black non-Latino	1,636,928 (14.8)	5,148 (23.2)
White Latino	2,491,475 (22.5)	4,731 (21.3)
Black Latino	125,716 (1.1)	231 (1.0)
Other	700,290 (6.3)	1,105 (4.9)
Maternal morbidities, n (%)		
Pregestational diabetes	67,697 (0.7)	2,796 (12.5)
Chronic hypertension	123,368 (1.3)	916 (4.1)
Preeclampsia	395,304 (4.2)	1,427 (6.4)
Eclampsia	16,682 (0.2)	156 (0.7)
Infant		
Gestational age at delivery, mean (SD)	38.8 (1.5)	36.3 (2.5)
Gestational age at delivery in weeks, n (%)		
32	40,183 (0.4)	2,152 (9.2)
33	58,274 (0.5)	1,993 (8.5)
34	108,854 (1.0)	2,284 (9.7)
35	179,159 (1.6)	2,445 (10.4)
36	384,986 (3.4)	2,973 (12.6)
37	925,965 (8.2)	3,202 (13.6)
38	2,047,163 (18.1)	3,305 (14.1)
39	4,226,642 (37.4)	2,812 (12.0)
40	2,558,635 (22.6)	1,742 (7.4)
41	728,269 (6.4)	496 (2.1)
42	46,505 (0.4)	115 (0.5)

Table 1 (Continued)

Characteristics	Live birth (11,304,635)	Stillbirth (23,519)
Gender, <i>n</i> (%)		
Female	5,520,803 (48.8)	11,236 (47.8)
Male	5,783,832 (51.2)	11,236 (52.2)
Unknown	0	1,047
Birth weight in grams, mean (SD)	3,331 (526)	2,920 (1,833)
Delivery method <i>n</i> (%)		
Vaginal	6,383,296 (69.0)	16,812 (71.8)
Cesarean	2,855,512 (30.9)	5,663 (24.2)
Unknown	7,601 (0.1)	946 (4.0)

Abbreviation: SD, standard deviation.

Note: Comparisons between live births and stillbirths using Student *t*-test for continuous variables or chi square for categorical variables. *p* < 0.001.

and the adjusted gestational age (► **Table 3**). As observed in the Kaplan–Meier survival curves, the lowest cumulative risk of stillbirth was observed for infants with birthweights between the 61st and the 95th centile. After adjusting the gestational age, the risk gradually increased for all birthweight categories above or below the reference group. The exception was fetuses between the 31st and the 40th centiles. The hazard ratios for infants with birthweights above the 95th centile were similar to those below the 5th centile.

Discussion

Main Findings

The data presented demonstrate that in the US population birthweight below the 5th and above the 95th centile was associated with the highest stillbirth risk within this US population cohort. Without accounting for time from death to delivery, the lowest risk for stillbirth was in the 71st to 95th centile categories. After adjusting for the gestational age at death, the risk gradually increased as birthweight centile

category diverged from the mean. The lowest risk was observed for newborns with birthweight between the 31st and the 40th centile, who had a stillbirth risk lower than the reference group, whereas those with birthweights below the 5th and above the 95th centile had a significantly higher risk than the adjacent birthweight categories.

The clinical estimate of gestational age at birth that is reported in the fetal death certificate does not account for intrapartum deaths or time from death to delivery. Most publications regarding stillbirth rates and risks do not account for the time elapsed between actual gestational age at time of fetal death and gestational age at time of delivery. The result is an overestimation of stillbirth rates among lower birthweight categories due to assignment of gestational age at time of delivery and not gestational age at death.

Many reports on risk and odds of stillbirth use the 10th percentile or above as a reference group when calculating the relative risk or odds of stillbirths in low birthweight categories.¹⁰ Recent research has suggested that the lowest stillbirth rates are somewhere between the 80th and 95th centile of birthweight.^{13,41} The argument for this finding has been that higher birthweight is consistent with improved maternal nutrition and protective against perinatal mortality.¹³ When we analyzed our data without adjustment for time from death to delivery, we also observed the same effect, a lower stillbirth risk for larger fetuses. This effect was not seen after we adjusted gestational age for time from death to delivery suggesting that the decreased stillbirth rate observed among larger birthweight categories resulted from the assignment of those stillbirths to later gestational ages and lower birthweight categories at time of delivery. The impact of adjusting gestational age downwards among fetuses in the 80th to 95th centile would be to increase the stillbirth risk among the larger fetuses at earlier gestational ages. This may have led to the increased stillbirth rate with birthweight over the 95th centile, although increased mortality in this group has been previously reported.¹¹

We have presented prospective and cumulative risks for stillbirth across all birthweight categories. The prospective

Fig. 2 Stillbirth risk according to gestational age and adjusted gestational age. Prospective stillbirth risk numerator are stillbirths occurring in a specific week of pregnancy, and the denominator is all ongoing pregnancies. Cumulative stillbirth risk is 1- the product of the cumulative risks of survival up to a week of gestation.

Table 2 Prospective and cumulative risk of stillbirth according to birthweight category and adjusted gestational age

Week	Ongoing	Live	Stillbirths	Prospective risk ^a	Cumulative risk ^a	Ongoing	Live	Stillbirths	Prospective risk ^a	Cumulative risk ^a
<div><div><3%</div><div><5%</div><div>>10–20%</div></div>										
32	524901	1437	137	2.61 (3.06–2.17)	2.61 (6.28–0)	816537	2319	191	2.34 (2.68–2)	2.34 (5.84–0)
33	523327	2738	182	3.49 (4–2.97)	6.1 (11.44–0.76)	814027	4173	230	2.83 (3.2–2.46)	5.17 (10.13–0.22)
34	520407	5333	207	4 (4.55–3.44)	10.1 (16.82–3.37)	809624	7691	265	3.29 (3.69–2.89)	8.46 (14.66–2.26)
35	514867	9240	216	4.23 (4.81–3.66)	14.32 (22.24–6.41)	801668	13206	285	3.58 (4.01–3.16)	12.04 (19.34–4.75)
36	505411	19814	223	4.5 (5.1–3.9)	18.82 (27.81–9.82)	788177	29138	299	3.86 (4.31–3.42)	15.9 (24.21–7.59)
37	485374	51730	217	4.72 (5.36–4.08)	23.53 (33.53–13.54)	758740	76531	278	3.86 (4.32–3.4)	19.76 (28.96–10.55)
38	433427	94925	208	5.39 (6.13–4.64)	28.91 (39.93–17.89)	681931	145701	255	4.18 (4.71–3.66)	23.93 (34.01–13.86)
39	338294	183912	139	5.64 (6.6–4.68)	34.53 (46.53–22.54)	535975	293225	185	4.75 (5.45–4.05)	28.67 (39.65–17.69)
40	154243	121032	74	7.89 (9.74–6.04)	42.4 (55.64–29.17)	242565	190293	91	6.17 (7.47–4.87)	34.83 (46.88–22.78)
41	33137	30969	22	12.45 (17.93–6.97)	54.81 (69.78–39.84)	52181	48947	27	9.73 (13.59–5.88)	44.54 (58.09–30.99)
42	2146	2144	2	18.59 (48.97–0)	73.33 (90.55–56.11)	3207	3204	3	18.66 (42.86–0)	63.15 (79.17–47.12)
<div><div>5–10%</div><div>>10–20%</div></div>										
32	620674	1851	100	1.61 (1.94–1.29)	1.61 (4.6–0)	1098704	3624	4.03	1.32 (1.54–1.1)	1.32 (4.08–0)
33	618723	2770	98	1.59 (1.91–1.26)	3.2 (7.21–0)	1094935	5145	131	1.2 (1.41–0.99)	2.52 (6.13–0)
34	615855	5165	118	1.92 (2.28–1.57)	5.12 (10.06–0.19)	1089659	10888	134	1.24 (1.45–1.02)	3.76 (8.050)
35	610572	8368	107	1.76 (2.11–1.42)	6.89 (12.53–1.25)	1078637	15891	142	1.33 (1.55–1.1)	5.08 (10–0.16)
36	602097	19431	100	1.69 (2.03–1.35)	8.57 (14.81–2.34)	1062604	35520	150	1.44 (1.67–1.2)	6.52 (12.02–1.02)
37	582566	48796	112	2.01 (2.39–1.63)	10.58 (17.45–3.71)	1026934	85636	153	1.55 (1.81–1.3)	8.07 (14.14–2.01)
38	533658	104824	95	1.97 (2.38–1.57)	12.55 (19.99–5.11)	941145	192596	137	1.62 (1.9–1.34)	9.69 (16.29–3.09)
39	428739	224454	83	2.62 (3.2–2.04)	15.17 (23.3–7.04)	748412	405941	90	1.65 (2–1.3)	11.34 (18.44–4.24)
40	204202	164461	33	2.7 (3.67–1.74)	17.87 (26.65–9.09)	342381	263064	53	2.51 (3.21–1.81)	13.85 (21.64–6.06)
41	39708	37578	8	3.82 (6.71–0.94)	21.69 (31.31–12.07)	79264	75440	12	2.89 (4.64–1.13)	16.73 (25.25–8.22)
42	2122	2119	3	28.14 (64.6–30)	49.85 (64.15–35.55)	3812	3809	3	15.7 (36.07–0)	32.43 (44.08–20.79)
<div><div>>20–30%</div><div>>30–40%</div></div>										
32	1061877	3619	117	1.1 (1.31–0.9)	1.1 (3.66–0)	1035326	3687	80	0.77 (0.95–0.6)	0.77 (3–0)
33	1058141	5501	108	1.02 (1.22–0.83)	2.13 (5.48–0)	1031559	5281	107	1.04 (1.24–0.84)	1.81 (4.95–0)
34	1052532	8946	121	1.15 (1.36–0.94)	3.28 (7.33–0)	1026171	9446	111	1.09 (1.29–0.88)	2.9 (6.74–0)
35	1043465	16349	181	1.75 (2.01–1.49)	5.03 (9.92–0.13)	1016614	15486	133	1.32 (1.55–1.09)	4.22 (8.74–0)

Table 2 (Continued)

Week	Ongoing	Live	Stillbirths	Prospective risk ^a	Cumulative risk ^a	Ongoing	Live	Stillbirths	Prospective risk ^a	Cumulative risk ^a
	<3%									
36	1026935	34339	131	1.3 (1.52–1.07)	6.33 (11.75–0.9)	1000995	31166	113	1.15 (1.36–0.93)	5.36 (10.4–0.33)
37	992465	85241	130	1.37 (1.61–1.13)	7.69 (13.63–1.76)	969716	74452	118	1.27 (1.5–1.03)	6.63 (12.17–1.08)
38	907094	187551	132	1.62 (1.91–1.34)	9.32 (15.79–2.84)	895146	198353	116	1.46 (1.73–1.19)	8.09 (14.16–2.01)
39	719411	395677	78	1.5 (1.84–1.15)	10.81 (17.75–3.87)	696677	418600	68	1.4 (1.74–1.05)	9.48 (16.01–2.95)
40	323656	252870	36	1.83 (2.45–1.2)	12.63 (20.09–5.17)	278009	214742	25	1.46 (2.07–0.86)	10.94 (17.92–3.96)
41	70750	67082	8	2.15 (3.77–0.53)	14.78 (22.81–6.75)	63242	59418	7	2.09 (3.78–0.39)	13.03 (20.6–5.46)
42	3660	3657	3	16.35 (37.56–0)	31.13 (42.55–19.71)	3817	3816	1	5.24 (18.11–0)	18.26 (27.13–9.39)
	>40–60%									
32	1863832	7352	198	1.06 (1.22–0.91)	1.06 (3.59–0)	927021	4237	133	1.44 (1.69–1.19)	1.44 (4.29–0)
33	1856282	11151	194	1.05 (1.2–0.9)	2.11 (5.46–0)	922651	5136	106	1.15 (1.38–0.93)	2.59 (6.24–0)
34	1844937	19796	266	1.45 (1.63–1.27)	3.56 (7.76–0)	917409	9451	137	1.5 (1.76–1.24)	4.09 (8.55–0)
35	1824875	30639	230	1.27 (1.44–1.1)	4.83 (9.64–0.02)	907821	14714	114	1.27 (1.5–1.03)	5.36 (10.39–0.32)
36	1794006	63660	247	1.4 (1.58–1.22)	6.23 (11.63–0.84)	892993	35987	147	1.68 (1.96–1.4)	7.04 (12.73–1.34)
37	1730099	150147	233	1.41 (1.59–1.22)	7.64 (13.56–1.72)	856859	85082	139	1.71 (2–1.42)	8.74 (15.03–2.45)
38	1579719	334071	188	1.33 (1.52–1.14)	8.97 (15.34–2.6)	771638	160940	96	1.39 (1.67–1.1)	10.13 (16.86–3.39)
39	1245460	697569	100	1.12 (1.34–0.89)	10.08 (16.81–3.36)	610602	333811	52	1.17 (1.5–0.84)	11.3 (18.38–4.21)
40	547791	422381	61	1.81 (2.28–1.34)	11.89 (19.15–4.64)	276739	214184	34	2 (2.71–1.3)	13.3 (20.94–5.66)
41	125349	118391	20	3.02 (4.42–1.62)	14.91 (22.98–6.85)	62521	58919	7	2.12 (3.84–0.4)	15.42 (23.61–7.23)
42	6938	6935	3	8.64 (19.84–0)	23.54 (33.54–13.55)	3595	3594	1	5.56 (19.23–0)	20.97 (30.43–11.5)
	>70–80%									
32	967482	3823	114	1.18 (1.4–0.96)	1.18 (3.81–0)	1061514	3718	180	1.7 (1.95–1.45)	1.7 (4.75–0)
33	963545	5233	165	1.72 (1.98–1.45)	2.9 (6.73–0)	1057616	4934	183	1.73 (1.99–1.48)	3.43 (7.56–0)
34	958147	9742	142	1.49 (1.74–1.24)	4.39 (8.99–0)	1052499	9532	201	1.92 (2.19–1.65)	5.35 (10.38–0.32)
35	948263	16046	170	1.81 (2.08–1.53)	6.19 (11.57–0.82)	1042766	16186	182	1.76 (2.02–1.5)	7.11 (12.83–1.38)
36	932047	28782	158	1.72 (2–1.45)	7.91 (13.93–1.9)	1026398	38204	234	2.32 (2.63–2.02)	9.43 (15.95–2.91)
37	903107	69974	130	1.5 (1.76–1.23)	9.41 (15.92–2.9)	987960	89739	143	1.52 (1.77–1.26)	10.94 (17.93–3.96)
38	833003	181259	109	1.47 (1.75–1.19)	10.88 (17.84–3.92)	898078	185972	105	1.3 (1.56–1.05)	12.25 (19.6–4.89)
39	651635	370384	76	1.63 (2.01–1.25)	12.51 (19.93–5.08)	712001	385278	66	1.27 (1.59–0.95)	13.52 (21.22–5.82)
40	281175	211756	19	1.08 (1.6–0.57)	13.59 (21.31–5.87)	326657	247646	42	2.07 (2.72–1.42)	15.58 (23.82–7.35)

(Continued)

Table 2 (Continued)

Week	Ongoing	Live	Stillbirths	Prospective risk ^a	Cumulative risk ^a	Ongoing	Live	Stillbirths	Prospective risk ^a	Cumulative risk ^a
	<3%					<5%				
41	69400	65147	10	2.71 (4.53–0.9)	16.3 (24.71–7.89)	78969	74397	12	2.87 (4.62–1.13)	18.45 (27.36–9.54)
42	4243	4241	2	9.41 (24.8–0)	25.7 (36.13–15.28)	4560	4559	1	4.38 (15.16–0)	22.83 (32.68–12.98)
	>90–95%					>95%				
32	605137	1685	167	2.76 (3.19–2.34)	2.76 (6.52–0)	1059143	2551	426	4.03 (4.41–3.64)	4.03 (8.46–0)
33	603285	2668	132	2.19 (2.57–1.81)	4.96 (9.82–0.09)	1056166	4236	489	4.64 (5.05–4.22)	8.66 (14.93–2.4)
34	600485	5687	169	2.83 (3.26–2.39)	7.78 (13.75–1.82)	1051441	8706	494	4.72 (5.14–4.3)	13.38 (21.04–5.71)
35	594629	9231	123	2.08 (2.46–1.71)	9.87 (16.52–3.21)	1042241	17007	495	4.79 (5.21–4.36)	18.16 (27–9.32)
36	585275	20125	146	2.54 (2.96–2.12)	12.4 (19.8–5)	1024739	37066	424	4.21 (4.62–3.81)	22.37 (32.13–12.61)
37	565004	47338	89	1.64 (1.99–1.29)	14.04 (21.88–6.2)	987249	88846	297	3.15 (3.51–2.79)	25.51 (35.9–15.12)
38	517577	111481	72	1.56 (1.93–1.19)	15.6 (23.83–7.36)	898106	202545	161	2.02 (2.34–1.7)	27.52 (38.29–16.76)
39	406024	233307	39	1.35 (1.79–0.91)	16.95 (25.51–8.38)	695400	402240	106	2.14 (2.56–1.73)	29.66 (40.82–18.5)
40	172678	128604	16	1.48 (2.25–0.71)	18.42 (27.32–9.52)	293054	215424	44	2.37 (3.1–1.65)	32.03 (43.61–20.46)
41	44058	40736	6	2.53 (4.77–0.3)	20.95 (30.41–11.49)	77586	71556	7	1.67 (3.03–0.31)	33.7 (45.56–21.84)
42	3316	3316	0			6023	6022	1	3.32 (11.48–0)	37.01 (49.41–24.61)

^aPer 10,000 pregnancies.

Fig. 3 (A) Cumulative stillbirth risk by birthweight category according to unadjusted gestational age. Stillbirths are per 10,000 births. Unadjusted gestational age is according to best clinical estimate registered in the birth or fetal death certificate. (B) Cumulative stillbirth risk by birthweight category according to adjusted gestational age. Stillbirths are per 10,000 births. Stillbirth gestational age is adjusted for interval between gestational age at time of diagnosis of fetal death and gestational age at time of delivery.

risks and odds for stillbirth have been reported for the U.S. population, and are higher than the gestational age adjusted rates we present.¹⁰ Odds ratios can compare birthweight categories but cannot account for the effect over time or gestational age. We have reported hazard ratios which allowed us to compare the cumulative risk of stillbirth over time between birthweight categories. The magnitude of the hazard ratio gradually increased as birthweight percentile diverged from the reference group, although there was no significant difference for those with birthweights in the 61st

to 70th or in the 31st to 40th birthweight centile categories. The hazard ratio was not as high as previously reported prospective risks or odds ratios for fetuses below the 5th centile, but higher than the odds ratios reported for fetuses with birthweights above the 95th centile.^{10–12}

Strengths and Limitations

This study used a large national vital statistics database with over 12,000,000 deliveries in three years. The large number of events included allowed us to robustly quantify the

Table 3 Hazard ratio for stillbirth according to birthweight category

Birthweight	Clinical estimate gestational age		Adjusted gestational age	
	Hazard ratio (95% CI)	<i>p</i>	Hazard ratio (95% CI)	<i>p</i>
< 3%	4.69 (4.96–4.43)	<0.0001	2.85 (3.07–2.66)	<0.0001
< 5%	2.33 (2.53–2.15)	<0.0001	1.55 (1.73–1.4)	<0.0001
5–10%	1.95 (2.09–1.82)	<0.0001	1.35 (1.47–1.24)	<0.0001
11–20%	1.51 (1.61–1.42)	<0.0001	1.04 (1.13–0.96)	0.329
21–30%	1.26 (1.35–1.18)	<0.0001	1.02 (1.1–0.94)	0.697
31–40%	1.01 (1.08–0.94)	0.817	0.89 (0.97–0.82)	0.008
61–70%	0.94 (1.01–0.87)	0.088	1.13 (1.23–1.04)	0.0042
71–80%	0.91 (0.98–0.84)	0.011	1.26 (1.37–1.17)	<0.0001
81–90%	0.79 (0.85–0.73)	<0.0001	1.45 (1.56–1.34)	<0.0001
91–95%	0.81 (0.89–0.74)	<0.0001	1.83 (1.99–1.68)	<0.0001
>95%	0.89 (1.01–0.79)	0.066	2.2 (2.43–1.99)	<0.0001

differences in prospective and cumulative stillbirth rates across birthweight categories by week of gestation. The prospective risk for stillbirth we applied best describes the population at risk for stillbirth at 32 to 42 weeks of gestation. It included all women who were still pregnant and at risk of having a stillborn infant, a clinically useful concept. In addition, we adjusted the gestational age among all stillbirths to account for delays between time of death and delivery. This adjustment has not previously been applied in reports of stillbirth rates from a large US population cohort.

An important limitation regarding the validity of using administrative data to derive prospective and cumulative stillbirth risk is the quality of the estimates of gestational age as well as information regarding the specific cause of death and timing of death. We used the obstetrical estimate of gestational age. This has been defined as “the estimate of the infant’s gestation in completed weeks based on the birth attendant’s final estimate of gestation which should be determined by all perinatal factors and assessments such as ultrasound, but not the neonatal exam”⁴², and it appears to provide a better estimate than the last menstrual period.⁴³

The stillbirth certificate does not account for the time transpired from the occurrence and diagnosis of stillbirth to the delivery.²⁹ The time elapsed is reported to be less than one week in 50%, and two or more weeks in 40% stillbirths beyond 24 weeks.²⁹ As a result, stillborns would be registered at a more advanced gestational age making them appear smaller for gestational age. To adjust for gestational age, we applied proportions from a previously validated algorithm developed to assign time of death in a cohort of stillbirths which were part of the Stillbirth Collaborative Research Network.²⁹ On the basis of their reported data, 25.2% of stillbirths occur 0–2 days before delivery (this includes both intrapartum deaths and recent fetal deaths), 12.4% occur at approximately 1 week, 17.4% at approximately 2 weeks, 8.4% at approximately 3 weeks and 20% at approximately 4 or more weeks prior to delivery. After adjusting for these predicted variations, we found a gradual increase in stillbirth rates with progressively lower or higher birthweight categories.

A third limitation is that in this database, stillbirths are all coded equally with no differentiation between intrapartum and antepartum deaths. When analyzing antepartum deaths, the correct denominator to use is the number of on-going pregnancies. However, when analyzing the risk of intrapartum deaths the correct denominator is the number of deliveries during a specific week and the overall risk is the addition of both. The frequency of intrapartum fetal death rates from previous reports show that the rates are lower than neonatal and antepartum death rates, remain stable from 37 to 42 weeks²¹, and would not be expected to modify the trends we have observed between birthweight categories.

Interpretation

Population-based analyses have reported on stillbirth rates according to absolute birthweight but not according to birthweight centile for gestational age.³³ Although odds ratios for rates according to birthweight centile have been reported,^{11,12} the increased prospective and cumulative still-

birth rates below the 5th centile and above the 95th centile have not been reported by gestational age in the US population. These results support our hypothesis that the prospective stillbirth rates gradually increase as birthweight centile decreases or increases, with the largest increase among births below the 5th centile or above the 95th centile.^{11,44,45} The decision to analyze stillbirth rates from 32 to 42 weeks of gestation was based on the uncertainty regarding timing of delivery and previous work demonstrating low birthweight to be a risk factor for stillbirth.^{11,44,45} Stillbirths prior to 30 weeks more often include aneuploidy, obstetrical, placental and infectious complications compared with deaths occurring after 30 weeks.⁴⁶ After 32 weeks, maternal, and obstetrical factors play a more significant role, including maternal age and parity.⁴⁷

We acknowledge that growth restriction is not a cause of stillbirth but a risk factor for stillbirth.⁴⁵ The information presented highlights the impact that growth restriction and excessive fetal growth have on stillbirth rates and risk. This effect is continuous and not a dichotomous effect observed below a certain threshold. It supports the current recommendations for surveillance among growth restricted fetuses, particularly those below the 5th centile.⁴⁸

An important finding in our analysis is that having a birthweight above the 95th centile was also associated with a significant increase in the risk for stillbirth when compared with infants within the median birthweight category. The magnitude of the increased risk was similar to that observed for birthweights below the 5th centile. In clinical practice, it is not uncommon to increase surveillance or intervene in cases where the fetus is considered to be large for gestational age, macrosomic or with birthweights above the 90th centile. This study defines the birthweight at which the risk increases as well as the magnitude of the risk.

Conclusion

Stillbirth risk gradually increases as birthweights diverge from the mean. Birthweights above the 95th centile or below the 5th centile convey the highest risk for stillbirth. The increased risk for stillbirth associated with birthweights above the 95th centile would qualify it as a risk factor for stillbirth similar to that attributed to birthweight below the 5th centile.¹⁴

Conflict of Interest

The authors have no competing interests to disclose.

Authors' Contribution

All three authors contributed in the conception, planning, carrying out analysis of the data, and writing up of the article.

Disclosure on Ethics Approval

The study was performed with publicly available administrative databases and did not require approval from our institutional review committee.

Funding
None.

Acknowledgments
None.

References

- 1 Goldenberg RL, McClure EM, Bhutta ZA, et al; Lancet's Stillbirths Series steering committee. Stillbirths: the vision for 2020. *Lancet* 2011;377(9779):1798–1805
- 2 Darmstadt GL, Kinney MV, Chopra M, et al; Lancet Every Newborn Study Group. Who has been caring for the baby? *Lancet* 2014;384(9938):174–188
- 3 Ananth CV, Joseph KS, Oyelese Y, Demissie K, Vintzileos AM. Trends in preterm birth and perinatal mortality among singletons: United States, 1989 through 2000. *Obstet Gynecol* 2005;105(5 Pt 1):1084–1091
- 4 Gyamfi-Bannerman C, Ananth CV. Trends in spontaneous and indicated preterm delivery among singleton gestations in the United States, 2005–2012. *Obstet Gynecol* 2014;124(6):1069–1074
- 5 Lees CC, Marlow N, van Wassenhaer-Leemhuis A, et al; TRUFFLE study group. 2 year neurodevelopmental and intermediate perinatal outcomes in infants with very preterm fetal growth restriction (TRUFFLE): a randomised trial. *Lancet* 2015;385(9983):2162–2172
- 6 Claussn B, Gardosi J, Francis A, Cnattingius S. Perinatal outcome in SGA births defined by customised versus population-based birth-weight standards. *BJOG* 2001;108(8):830–834
- 7 Getahun D, Ananth CV, Kinzler WL. Risk factors for antepartum and intrapartum stillbirth: a population-based study. *Am J Obstet Gynecol* 2007;196(6):499–507
- 8 Malhotra A, Yahya Z, Sasi A, et al. Does fetal growth restriction lead to increased brain injury as detected by neonatal cranial ultrasound in premature infants? *J Paediatr Child Health* 2015;51(11):1103–1108
- 9 Murray E, Fernandes M, Fazel M, Kennedy SH, Villar J, Stein A. Differential effect of intrauterine growth restriction on childhood neurodevelopment: a systematic review. *BJOG* 2015;122(8):1062–1072
- 10 Pilliod RA, Cheng YW, Snowden JM, Doss AE, Caughey AB. The risk of intrauterine fetal death in the small-for-gestational-age fetus. *Am J Obstet Gynecol* 2012;207(4):318.e1–318.e6
- 11 Bukowski R, Hansen NI, Willinger M, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Stillbirth Collaborative Research Network. Fetal growth and risk of stillbirth: a population-based case-control study. *PLoS Med* 2014;11(4):e1001633
- 12 Moraitis AA, Wood AM, Fleming M, Smith GC. Birth weight percentile and the risk of term perinatal death. *Obstet Gynecol* 2014;124(2 Pt 1):274–283
- 13 Vasak B, Koenen SV, Koster MP, et al. Human fetal growth is constrained below optimal for perinatal survival. *Ultrasound Obstet Gynecol* 2015;45(2):162–167
- 14 Spong CY, Mercer BM, D'Alton M, Kilpatrick S, Blackwell S, Saade G. Timing of indicated late-preterm and early-term birth. *Obstet Gynecol* 2011;118(2 Pt 1):323–333
- 15 Figueras F, Gardosi J. Intrauterine growth restriction: new concepts in antenatal surveillance, diagnosis, and management. *Am J Obstet Gynecol* 2011;204(4):288–300
- 16 Hecher K, Bilardo CM, Stigter RH, et al. Monitoring of fetuses with intrauterine growth restriction: a longitudinal study. *Ultrasound Obstet Gynecol* 2001;18(6):564–570
- 17 Romero R, Kalache KD, Kadar N. Timing the delivery of the preterm severely growth-restricted fetus: venous Doppler, cardiotocography or the biophysical profile? *Ultrasound Obstet Gynecol* 2002;19(2):118–121
- 18 Turan OM, Turan S, Gungor S, et al. Progression of Doppler abnormalities in intrauterine growth restriction. *Ultrasound Obstet Gynecol* 2008;32(2):160–167
- 19 Crimmins S, Desai A, Block-Abraham D, Berg C, Gembruch U, Baschat AA. A comparison of Doppler and biophysical findings between liveborn and stillborn growth-restricted fetuses. *Am J Obstet Gynecol* 2014;211(6):669.e1–669.e10
- 20 MacDorman MF, Reddy UM, Silver RM. Trends in Stillbirth by Gestational Age in the United States, 2006–2012. *Obstet Gynecol* 2015;126(6):1146–1150
- 21 Smith GCS. Life-table analysis of the risk of perinatal death at term and post term in singleton pregnancies. *Am J Obstet Gynecol* 2001;184(3):489–496
- 22 Smith GC. Re: "A proportional hazards model with time-dependent covariates and time-varying effects for analysis of fetal and infant death". *Am J Epidemiol* 2005;161(1):100–101, author reply 100–101
- 23 American College of Obstetricians and Gynecologists. ACOG Practice bulletin no. 134: fetal growth restriction. *Obstet Gynecol* 2013;121(5):1122–1133
- 24 Battaglia FC, Lubchenco LO. A practical classification of newborn infants by weight and gestational age. *J Pediatr* 1967;71(2):159–163
- 25 Resnik R. Intrauterine growth restriction. *Obstet Gynecol* 2002;99(3):490–496
- 26 National Center for Health Statistics. Rockville, MD. Available at: http://www.cdc.gov/nchs/data_access/Vitalstatsonline.htm. Last accessed September 10, 2014.
- 27 Martin JA, Wilson EC, Osterman MJ, Saadi EW, Sutton SR, Hamilton BE. Assessing the quality of medical and health data from the 2003 birth certificate revision: results from two states. *Natl Vital Stat Rep*. 2013 Jul 22;62(2):1–19
- 28 Duryea EL, McIntire DD, Leveno KJ. The rate of preterm birth in the United States is affected by the method of gestational age assignment. *Am J Obstet Gynecol* 2015;213(2):231.e1–231.e5
- 29 Conway DL, Hansen NI, Dudley DJ, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Stillbirth Collaborative Research Network. An algorithm for the estimation of gestational age at the time of fetal death. *Paediatr Perinat Epidemiol* 2013;27(2):145–157
- 30 Altman DG, Chitty LS. Design and analysis of studies to derive charts of fetal size. *Ultrasound Obstet Gynecol* 1993;3(6):378–384
- 31 Altman DG, Chitty LS. Charts of fetal size: 1. Methodology. *Br J Obstet Gynaecol* 1994;101(1):29–34
- 32 Royston P, Wright EM. How to construct 'normal ranges' for fetal variables. *Ultrasound Obstet Gynecol* 1998;11(1):30–38
- 33 MacDorman MF, Kirmeyer SE, Wilson EC. Fetal and perinatal mortality, United States, 2006. *Natl Vital Stat Rep* 2012 Aug 28;60(8):1–22
- 34 Yudkin PL, Wood L, Redman CW. Risk of unexplained stillbirth at different gestational ages. *Lancet* 1987;1(8543):1192–1194
- 35 Smith GC, Pell JP, Dobbie R. Caesarean section and risk of unexplained stillbirth in subsequent pregnancy. *Lancet* 2003;362(9398):1779–1784
- 36 K.G. M. Stallard E. Medical Demography: Interaction of Disability Dynamics and Mortality. In: Martin LG, Preston SH eds, National Research Council (US) Committee on Population: Demography of Aging. Washington, DC: National Academies Press; 1994
- 37 Greenland S, Neutra R. Control of confounding in the assessment of medical technology. *Int J Epidemiol* 1980;9(4):361–367
- 38 Macdorman MF, Kirmeyer S. The challenge of fetal mortality. NCHS data brief 2009(16):1–8. Available at: <http://www.cdc.gov/nchs/data/databriefs/db16.pdf> Accessed September 15, 2015.
- 39 Altman DG, Bland JM. Missing data. *BMJ* 2007;334(7590):424

- 40 Society for Research in Child Development. Missing data: What to do with or without them. *Monogr Soc Res Child Dev* 2006;71(3):42–64
- 41 Morales-Roselló J, Khalil A, Morlando M, Papageorgiou A, Bhide A, Thilaganathan B. Changes in fetal Doppler indices as a marker of failure to reach growth potential at term. *Ultrasound Obstet Gynecol* 2014;43(3):303–310
- 42 Center for Disease Control and Prevention. Birth edit specifications for the 2003 proposed revision of the United States standard certificate of birth. Available at: http://www.cdc.gov/nchs/data/dvs/birth_edit_specifications.pdf> Accessed: September 30, 2015
- 43 Duryea EL, Hawkins JS, McIntire DD, Casey BM, Leveno KJ. A revised birth weight reference for the United States. *Obstet Gynecol* 2014;124(1):16–22
- 44 Gardosi J, Madurasinghe V, Williams M, Malik A, Francis A. Maternal and fetal risk factors for stillbirth: population based study. *BMJ* 2013;346:f108
- 45 Edmunds SF, Silver RM. Stillbirth reduction efforts and impact on early births. *Clin Perinatol* 2013;40(4):611–628
- 46 Stillbirth Collaborative Research Network Writing Group. Causes of death among stillbirths. *JAMA* 2011;306(22):2459–2468
- 47 Waldenström U, Cnattingius S, Norman M, Schytt E. Advanced Maternal Age and Stillbirth Risk in Nulliparous and Parous Women. *Obstet Gynecol* 2015;126(2):355–362
- 48 Darmstadt GL, Shiffman J, Lawn JE. Advancing the newborn and stillbirth global agenda: priorities for the next decade. *Arch Dis Child* 2015;100(Suppl 1):S13–S18