J. A. M. MERCER, C. M. COHEN, S. R. SHUKEN, A. M. WAGNER, M. W. SMITH, F. R. MOSS, III, M. D. SMITH, R. VAHALA, A. GONZALEZ-MARTINEZ,* S. G. BOXER,* N. Z. BURNS* (STANFORD UNIVERSITY, USA AND AALTO UNIVERSITY, FINLAND)
Chemical Synthesis and Self-Assembly of a Ladderane Phospholipid
J. Am. Chem. Soc. **2016**, *138*, 15845–15848.

Total Synthesis of a Ladderane Phospholipid

Significance: Gonzalez-Martinez, Boxer, Burns and co-workers report an impressive total synthesis of a ladderane phospholipid based on strategic [2+2] cycloadditions of bicyclohexene **B**, which is obtained by means of a Ramberg–Bäcklund ring contraction of sulfoxide **A**.

SYNFACTS Contributors: Erick M. Carreira, Leonardo J. Nannini
Synfacts 2017, 13(02), 0117 Published online: 18.01.2017
DOI: 10.1055/s-0036-1589874; Reg-No.: C08416SF

Comment: Bicyclohexene **B** irradiated in the presence of CuOTf gave pentacycle **F**, which was subjected to an oxidative chlorination–elimination sequence to give cyclobutene **I**. Enantioselective hydroboration and four further steps yielded [5]-ladderanoic acid.

Category

Synthesis of Natural Products and Potential Drugs

Key words

[2+2] cycloaddition

[5]-ladderanoic acid

[3]-ladderanol

ladderane phospholipids

Ramberg–Bäcklund reaction

Synfact
of the month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.