L. J. CHENG, A. P. N. BROWN, C. J. CORDIER* (IMPERIAL COLLEGE LONDON, UK)

Enantioselective Propargylic [1,3]-Rearrangements: Copper-Catalyzed O-to-N Migrations toward C–N Bond Formation
Chem. Sci. **2017**, *8*, 4299–4305.

Copper-Catalyzed Enantioselective [1,3]-Rearrangement to Chiral Propargylic Amines

Selected examples:

Synthetic utility:

Significance: Thermal [3,3]-sigmatropic rearrangements are well preceded and can be achieved at ambient temperature with high levels of enantioselectivity (see, for example: M. P. Watson, L. E. Overman, R. G. Bergman *J. Am. Chem. Soc.* **2007**, *129*, 5031). In contrast, [1,3]-sigmatropic rearrangements are thermally disallowed, but can be promoted at high temperatures with the use of transition-metal catalysts. No examples of [1,3]-rearrangements affording propargylic products or their enantioselective variants have been previously reported.

Comment: The present report details the first example of an enantioselective [1,3]-rearrangement. The alkyne starting materials **1** can be converted into pyridone alkynes **3** with a diverse substrate scope, affording products in high yields and with high er. Detailed mechanistic studies indicate that the reaction proceeds through allene intermediate **2** with copper coordination, followed by a formal [1,3]-rearrangement. The enantioselectivity is derived from a BINAP phosphoryl ligand **2** through CuTC coordination. The synthetic utility of the products was shown by carrying **4** forward to the synthesis of cysteine protease inhibitor **11**. This reaction will find utility due to the mild reaction conditions and the ability to modify the pyridone core to generate chiral amines similar to **10**.

SYNFACTS Contributors: Victor Snieckus, Vince M. Lombardo (Pfizer)

Synfacts 2017, 13(07), 0686 Published online: 19.06.2017

DOI: 10.1055/s-0036-1590587; Reg-No.: V07017SF