The Brain after Cardiac Arrest

Jonathan Elmer, MD, MS^{1,2} Clifton W. Callaway, MD, PhD¹

¹ Department of Emergency Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania

² Department of Critical Care Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania

Address for correspondence Clifton W. Callaway, MD, PhD, Iroquois Building, Suite 400A, 3600 Forbes Avenue, Pittsburgh, PA 15260 (e-mail: callawaycw@upmc.edu).

Semin Neurol 2017;37:19–24.

Abstract

Cardiac arrest is common and deadly. Most patients who are treated in the hospital after achieving return of spontaneous circulation still go on to die from the sequelae of anoxic brain injury. In this review, the authors provide an overview of the mechanisms and consequences of postarrest brain injury. Special attention is paid to potentially modifiable mechanisms of secondary brain injury including seizures, hyperpyrexia, cerebral hypoxia and hypoperfusion, oxidative injury, and the development of cerebral edema. Finally, the authors discuss the outcomes of cardiac arrest survivors with a focus on commonly observed patterns of injury as well as the scales used to measure patient outcome and their limitations.

Keywords

- cardiac arrest
- brain injury
- outcomes

Cardiac arrest (CA) is the most common cause of death in the developed world, claiming over 17 million lives annually.^{1–3} Despite advances in care, the majority of patients who are treated after the return of spontaneous circulation (ROSC) will succumb to the sequelae of hypoxic-ischemic brain injury before hospital discharge. Similar to traumatic brain injury, hypoxic-ischemic brain injury after CA results in both primary and secondary injuries, with distinct mechanisms and treatment strategies (► **Fig. 1**). High-quality cardiopulmonary resuscitation (CPR) initiated as soon as possible after collapse, and early defibrillation to rapidly restore normal perfusion, are the most important interventions to reduce the severity of primary brain injury.⁴ It remains unclear whether care provided after ROSC can reduce the severity of primary brain injury or if therapies after CPR improve outcomes only by reducing the risk of secondary brain injury. Evidence-based therapies in the hours to days after ROSC that improve neurologic outcomes include targeted temperature management; early coronary revascularization; delayed, multimodal neurologic prognostication; and postacute care rehabilitation.⁵ In this review, we provide an overview of the mechanisms and consequences of postarrest brain injury as an introduction to the many specific therapies that are reviewed in depth elsewhere in this issue.

Neuronal Death after Cardiac Arrest

The physiological and molecular events associated with brain injury after CA are complex and have been described in detail.^{6–8} In the clinical setting, hypoxic-ischemic brain injury cannot be ascribed to any single mechanism. It is perhaps unsurprising then that well-controlled experiments which block or promote a single molecular pathway are ineffective for improving brain recovery,^{9,10} but pleiotropic, relatively nonspecific interventions may improve recovery. Examples of the latter include reduction of brain temperature and general anesthesia.^{11,12} Substantial current research focuses on understanding the fact that histological signs of post-CA neuronal death are delayed for hours or days after ischemia-reperfusion. This observation has prompted an analogy to other types of delayed, programmed cell death such as apoptosis, autophagy, necroptosis, and ferroptosis.^{6,13,14} No pure model of cell death is exactly like the delayed neuronal death observed after global ischemia-reperfusion, and it is probably best to consider each of these mechanisms as a potential contributor to post-CA brain injury. In patients, other macroscopic disturbances of multiple organ systems, endothelial function, and homeostatic processes also contribute to the progression of brain injury and neuronal death.

Issue Theme Hypoxic-Ischemic Encephalopathy; Guest Editors, Hans Friberg, MD, PhD, and Tobias Cronberg, MD, PhD

Copyright © 2017 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.
Tel: +1(212) 584-4662.

DOI <http://dx.doi.org/10.1055/s-0036-1597833>.
ISSN 0271-8235.

Fig. 1 Conceptual map of brain injury after cardiac arrest. ASA, aspirin; BBB, blood–brain barrier; ROS, reactive oxygen species.

Cerebral Homeostasis after Cardiac Arrest

Potentially modifiable mechanisms of secondary brain injury include seizures, hyperpyrexia, cerebral hypoxia, oxidative stress, the development of cerebral edema, microcirculatory dysfunction, impaired autoregulation of cerebral blood flow, and increased cerebral vascular resistance.^{15–24} Not all of these phenomena occur to the same degree in every patient. However, like the molecular events, each one has the potential to add to brain injury. The provision of high-quality postarrest critical care can minimize the risk from these potentially modifiable factors.

Seizures occur in 10 to 20% of comatose patients after CA, may worsen excitotoxicity, and are associated with worse outcomes.^{23,25} It remains unknown whether seizures are simply an epiphenomenon of more severe injury or by themselves produce secondary injury.^{23,24} Regardless, monitoring of the electrical state of the brain provides both prognostic information as well as guidance for intensive care treatment.²⁶

Cerebral autoregulation of blood flow can be absent or right-shifted after cardiac arrest.¹⁹ Thus, the brain is particularly sensitive to hypotension and may require higher than normal arterial pressure to maintain normal blood flow. Combined with the fact that systemic hypotension and shock after CA are also common,²⁷ and microvascular dysfunction can lead to areas of no-reflow despite return of a perfusing rhythm,²⁸ significant cerebral hypoperfusion is a real risk. Aggressive early resuscitation, coronary revascularization, and vasopressor use to increase mean arterial pressure may reduce this risk. At a population level, higher arterial pressures after CA are associated with improved patient outcomes even after adjusting for the use of vasopressors to achieve these goals.^{29,30} Individual need may vary depending on baseline

blood pressure and severity of postarrest illness. Direct invasive monitoring of cerebral oxygenation and blood flow can reveal substantial interpatient heterogeneity (→ Fig. 2). Population-based hemodynamic goals may lead to hyperemic cerebral perfusion in some patients, theoretically worsening the risk of vasogenic edema or oxidative injury (see below) and clinically significant brain tissue hypoxia in others. Unfortunately, broadly applicable strategies do not yet exist to determine the optimal cerebral perfusion pressure for an individual patient as it changes over time, or alternatively to detect in real-time the presence of hyperemia or hypoperfusion.

Respiratory failure and dependence on mechanical ventilation are also common after CA, and imprudent ventilator management may potentiate secondary brain injury. Hyperventilation-associated hypocapnia may lead to cerebral vasoconstriction and exacerbate hypoperfusion, whereas normocapnia or even mild permissive hypercapnia are associated with improved neurologic outcomes.^{31,32} This effect relies on intact chemoregulation of cerebral blood flow, which may or may not be present after severe brain injury.³³ Because broadly applicable real-time measures of individual need and treatment responsiveness are not available, clinicians now treat individual patients using population-based estimates. Finally, severe hyperoxemia may increase oxidative stress through formation of reactive oxygen species, whereas hypoxemia causes unnecessary cellular hypoxia.^{34,35} In most patients, both hypocapnia and prolonged hyperoxemia are avoidable with careful ventilator management.

Cerebral Edema

Cerebral edema occurs early after ROSC in some patients or during rewarming in others, and regardless of timing is an

Fig. 2 Direct monitoring of brain tissue oxygenation or cerebral blood flow in two patients after cardiac arrest reveals significant between-patient variation in the necessary mean arterial pressure (MAP) to maintain adequate brain oxygenation. In the left panel, brain tissue hypoxia at a threshold of 20 mmHg (red line) is present despite intact autoregulation at blood MAPs below ~110 mmHg, but can be overcome by increasing MAP >110 mmHg. In the right panel, autoregulation is mostly intact and brain oxygenation is adequate across a range of MAPs.

ominous marker of severe brain injury. However, favorable outcomes can occur in patients with mild-to-moderate early edema.¹⁵ The timing of early cerebral edema and mechanistic studies suggest a vasogenic rather than cytotoxic mechanism for early edema,³⁶ and indeed hyperosmolar therapy may reduce radiographic signs of edema and intracranial hypertension (►Fig. 3). Reducing gross edema theoretically will improve microvascular blood flow to vulnerable brain regions, but it is unknown whether treating postanoxic cerebral edema improves patient outcomes.

Clinical Assessment of Brain Injury and Prognosis

Multimodal neurologic prognostication of coma after CA is also reviewed in detail in this issue. From the perspective of cerebral vulnerability after CA, it should be noted that inaccurate or inappropriately early neurologic prognostication after CA can

contribute substantially to avoidable mortality after CA.³⁷ Guidelines recommend delaying withdrawal of life-sustaining therapy based on neurologic prognosis until at least 72 hours after ROSC because the accuracy of prognostic information available prior to this time is limited.^{38–40} Even patients who remain comatose 72 hours after ROSC may go on to awaken and have favorable recoveries.⁴¹ Unfortunately, the withdrawal of life-sustaining therapy in the first 24 hours after ROSC based on perceived neurologic prognosis remains common.³⁷ Even the best postarrest critical care can be undermined by therapeutic nihilism and inappropriate care withdrawal.

Regional Heterogeneity Produces Different Brain Injury Phenotypes

Several neuronal subtypes and brain regions are particularly sensitive to the physiological and cellular effects of anoxic injury and circulatory arrest. This results in distinct

Fig. 3 Cerebral edema may develop early after cardiac arrest or during rewarming. In this patient, baseline computed tomography (CT) scan shortly after cardiac arrest was unremarkable (left). Just at the time of the second CT scan, the patient showed clinical signs of herniation, prompting empiric therapy with hypertonic saline. Basilar and quadrigeminal cisterns were collapsed (center, arrows). After 24 hours, the subject had recovery of brainstem reflexes and improved appearance on CT scan (right). Ultimately, this patient had no recovery of cortical function.

phenotypes among survivors of CA. For example, vulnerable cell populations include the hippocampal CA1 pyramidal neurons in the mesial temporal lobe.^{42,43} As a consequence, memory impairments, particularly the ability to consolidate short-term memory, are common after CA even among patients with otherwise favorable functional outcomes.^{44,45} Cerebellar Purkinje cell and basal ganglia injury with cortical sparing may lead to postanoxic myoclonus and a range of other movement disorders in patients who are cognitively intact.^{46–48} Cortical pyramidal neuron injury can cause impaired attention, processing speed, and/or executive function depending on the region of injury.^{44,45,49} Of note, many of these deficits also occur in patients with other critical illnesses who have not suffered CA, supporting the concern that a combination of critical illness, intensive care, and patient comorbidities can potentiate brain injury itself.^{50,51} Regardless, many patients experience substantial improvement in symptom severity in the months after CA, and rehabilitation improves recovery. Randomized trials have demonstrated that focused rehabilitation can improve social engagement, quality of life, and emotional outcomes as well as reduce medical complications in those discharged after CA.^{52,53}

Quantifying Neurologic Recovery

The common deficits described above are challenging to quantify. Indeed, most trials to date have relied on gross measures of functional or neurologic impairment such as Cerebral Performance Category (CPC) or modified Rankin Scale Score (mRS) as a primary outcome, in part because of their simplicity compared with more detailed neurocognitive testing (→Table 1). In awake patients, CPC and mRS capture varying aspects of functional recovery across levels, and are influenced not only by the severity of illness, but also by environmental and personal factors that modify the interplay

between neurologic disability (e.g., arm paresis), activity (ability to carry out activities of daily living [ADLs]), and participation (ability to return to work).^{54,55} This means that the use of CPC or mRS as a primary trial outcome risks a failure to detect true physiological differences between treatment arms because of environmental or personal modifiers. These scales are dynamic over the course of recovery and sensitive to setting, adding an additional level of complexity when considering the timing of outcome assessment. For example, early after hospital discharge, functional measures may worsen as patients move from a sheltered hospital environment to home where there is less assistance for ADLs, then improve over the following 6 to 12 months.^{56,57} In addition, these scales are relatively insensitive to deficits in memory, executive function, and processing speed that develop commonly after CA and that are important to patient experience.^{45,58,59} More sensitive measures have been used and validated after CA, but have not thus far been implemented as clinical trial endpoints.^{45,50,59} Despite their limitations, short-term measures of functional recovery at hospital discharge do predict long-term survival. Those discharged with moderate or severe disability as measured by CPC are at substantially higher hazard of death compared with those with mild or no disability.⁶⁰

Conclusion

Post-CA brain injury results in multiple molecular and physiological changes over the hours to days after CPR. In practice, monitoring the status of the brain after CA and the prevention of secondary injury are central organizing principles for postarrest care. Reducing the deleterious effects of impaired homeostasis guides blood pressure and ventilator, fluid, and temperature management, as well as neurophysiological monitoring during postarrest intensive care. Proper multimodal assessment of

Table 1 Commonly used measures of neurologic recovery after cardiac arrest

Modified Rankin Scale	Cerebral Performance Category
0– <u>No symptoms at all</u>	1–Good recovery: Conscious, alert, able to work , <u>might have mild neurologic or psychological deficit.</u>
1– No significant disability despite symptoms ; <u>able to carry out all usual duties and activities</u>	2–Moderate disability: Conscious, sufficient cerebral function for independent activities of daily life. <u>Able to work in sheltered environment</u>
2– Slight disability ; <u>unable to carry out all previous activities, but able to look after own affairs without assistance</u>	3–Severe disability: Conscious , dependent on others for daily support because of impaired brain function; ranges from ambulatory state to severe dementia or paralysis
3– Moderate disability ; requiring some help, but able to walk without assistance	4–Coma or vegetative state: Any degree of coma without the presence of all brain death criteria
4– Moderately severe disability ; <u>unable to walk without assistance and unable to attend to own bodily needs without assistance</u>	5–Death: Apnea, electroencephalographic silence, etc.
5– Severe disability ; <u>bedridden, incontinent, requiring constant nursing care and attention</u>	
6–Death: Apnea, electroencephalographic silence, etc.	

Note: Cerebral Performance Category and modified Rankin Scale measure different aspects of recovery. Neurologic impairments are listed in bold, function capability is in italics, and participation ability is underlined.

coma is critical to avoid premature withdrawal of life support resulting in death. The pattern of brain injury that remains after intensive care varies from mild to severe, with some brain regions being particularly susceptible. Proper evaluation of cognition and other functions after emergence from coma is critical for guiding postacute rehabilitation and support services. Traditional outcome scales (MRS, CPC) may be too coarse to detect cognitive issues that affect patients' quality of life.

Acknowledgments

Dr. Elmer's research time is supported by the National Heart, Lung and Blood Institute through grant number 5K12HL109068.

References

- Mozaffarian D, Benjamin EJ, Go AS, et al; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2015 update: a report from the American Heart Association. *Circulation* 2015;131(4):e29–e322
- Lozano R, Naghavi M, Foreman K, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380(9859):2095–2128
- Nolan JP, Hazinski MF, Aickin R, et al. Part 1: Executive summary: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation* 2015;95:e1–e31
- Kleinman ME, Brennan EE, Goldberger ZD, et al. Part 5: Adult Basic Life Support and Cardiopulmonary Resuscitation Quality: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2015;132(18, Suppl 2):S414–S435
- Callaway CW, Donnino MW, Fink EL, et al. Part 8: Post-cardiac arrest care: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2015;132(18, Suppl 2):S465–S482
- Conrad M, Angeli JP, Vandenabeele P, Stockwell BR. Regulated necrosis: disease relevance and therapeutic opportunities. *Nat Rev Drug Discov* 2016;15(5):348–366
- Safar P, Behring W. Brain resuscitation after cardiac arrest. In: Layon AJ, Gabrielli A, Friedman WA, eds. *Textbook of neuro-intensive care*. Philadelphia: Saunders; 2004:457–498
- Greer DM. Mechanisms of injury in hypoxic-ischemic encephalopathy: implications to therapy. *Semin Neurol* 2006;26(4):373–379
- Callaway CW, Ramos R, Logue ES, Betz AE, Wheeler M, Repine MJ. Brain-derived neurotrophic factor does not improve recovery after cardiac arrest in rats. *Neurosci Lett* 2008;445(1):103–107
- Teschendorf P, Vogel P, Wippel A, et al. The effect of intracerebroventricular application of the caspase-3 inhibitor zDEVD-FMK on neurological outcome and neuronal cell death after global cerebral ischaemia due to cardiac arrest in rats. *Resuscitation* 2008;78(1):85–91
- Derwall M, Timper A, Kottmann K, Rossaint R, Fries M. Neuroprotective effects of the inhalational anesthetics isoflurane and xenon after cardiac arrest in pigs. *Crit Care Med* 2008;36(11, Suppl):S492–S495
- Hicks SD, DeFranco DB, Callaway CW. Hypothermia during reperfusion after asphyxial cardiac arrest improves functional recovery and selectively alters stress-induced protein expression. *J Cereb Blood Flow Metab* 2000;20(3):520–530
- Horn M, Schlote W. Delayed neuronal death and delayed neuronal recovery in the human brain following global ischemia. *Acta Neuropathol* 1992;85(1):79–87
- Au AK, Chen Y, Du L, et al. Ischemia-induced autophagy contributes to neurodegeneration in cerebellar Purkinje cells in the developing rat brain and in primary cortical neurons in vitro. *Biochim Biophys Acta* 2015;1852(9):1902–1911
- Metter RB, Rittenberger JC, Guyette FX, Callaway CW. Association between a quantitative CT scan measure of brain edema and outcome after cardiac arrest. *Resuscitation* 2011;82(9):1180–1185
- Oku K, Kuboyama K, Safar P, et al. Cerebral and systemic arteriovenous oxygen monitoring after cardiac arrest. Inadequate cerebral oxygen delivery. *Resuscitation* 1994;27(2):141–152
- Wolfson SK Jr, Safar P, Reich H, et al. Dynamic heterogeneity of cerebral hypoperfusion after prolonged cardiac arrest in dogs measured by the stable xenon/CT technique: a preliminary study. *Resuscitation* 1992;23(1):1–20
- Nishizawa H, Kudoh I. Cerebral autoregulation is impaired in patients resuscitated after cardiac arrest. *Acta Anaesthesiol Scand* 1996;40(9):1149–1153
- Sundgreen C, Larsen FS, Herzog TM, Knudsen GM, Boesgaard S, Aldershvile J. Autoregulation of cerebral blood flow in patients resuscitated from cardiac arrest. *Stroke* 2001;32(1):128–132
- Schaafsma A, de Jong BM, Bams JL, Haaxma-Reiche H, Pruim J, Zijlstra JG. Cerebral perfusion and metabolism in resuscitated patients with severe post-hypoxic encephalopathy. *J Neurol Sci* 2003;210(1–2):23–30
- Buunk G, van der Hoeven JG, Meinders AE. Cerebrovascular reactivity in comatose patients resuscitated from a cardiac arrest. *Stroke* 1997;28(8):1569–1573
- Hickey RW, Kochanek PM, Ferimer H, Alexander HL, Garman RH, Graham SH. Induced hyperthermia exacerbates neurologic neuronal histologic damage after asphyxial cardiac arrest in rats. *Crit Care Med* 2003;31(2):531–535
- Rittenberger JC, Popescu A, Brenner RP, Guyette FX, Callaway CW. Frequency and timing of nonconvulsive status epilepticus in comatose post-cardiac arrest subjects treated with hypothermia. *Neurocrit Care* 2012;16(1):114–122
- Rossetti AO, Carrera E, Oddo M. Early EEG correlates of neuronal injury after brain anoxia. *Neurology* 2012;78(11):796–802
- Youn CS, Callaway CW, Rittenberger JC. Post Cardiac Arrest Service. Combination of initial neurologic examination and continuous EEG to predict survival after cardiac arrest. *Resuscitation* 2015;94:73–79
- Friberg H, Westhall E, Rosén I, Rundgren M, Nielsen N, Cronberg T. Clinical review: Continuous and simplified electroencephalography to monitor brain recovery after cardiac arrest. *Crit Care* 2013;17(4):233
- Coppler PJ, Elmer J, Calderon L, et al; Post Cardiac Arrest Service. Validation of the Pittsburgh Cardiac Arrest Category illness severity score. *Resuscitation* 2015;89:86–92
- Ames A III, Wright RL, Kowada M, Thurston JM, Majno G. Cerebral ischemia. II. The no-reflow phenomenon. *Am J Pathol* 1968;52(2):437–453
- Beylin ME, Perman SM, Abella BS, et al. Higher mean arterial pressure with or without vasoactive agents is associated with increased survival and better neurological outcomes in comatose survivors of cardiac arrest. *Intensive Care Med* 2013;39(11):1981–1988
- Kilgannon JH, Roberts BW, Jones AE, et al. Arterial blood pressure and neurologic outcome after resuscitation from cardiac arrest*. *Crit Care Med* 2014;42(9):2083–2091
- Schneider AG, Eastwood GM, Bellomo R, et al. Arterial carbon dioxide tension and outcome in patients admitted to the intensive care unit after cardiac arrest. *Resuscitation* 2013;84(7):927–934
- Roberts BW, Kilgannon JH, Chansky ME, Mittal N, Wooden J, Trzeciak S. Association between postresuscitation partial pressure of arterial carbon dioxide and neurological outcome in patients with post-cardiac arrest syndrome. *Circulation* 2013;127(21):2107–2113

- 33 Carmona Suazo JA, Maas AI, van den Brink WA, van Santbrink H, Steyerberg EW, Avezaat CJ. CO₂ reactivity and brain oxygen pressure monitoring in severe head injury. *Crit Care Med* 2000;28(9):3268–3274
- 34 Elmer J, Scutella M, Pullalarevu R, et al; Pittsburgh Post-Cardiac Arrest Service (PCAS). The association between hyperoxia and patient outcomes after cardiac arrest: analysis of a high-resolution database. *Intensive Care Med* 2015;41(1):49–57
- 35 Kilgannon JH, Jones AE, Shapiro NI, et al; Emergency Medicine Shock Research Network (EMShockNet) Investigators. Association between arterial hyperoxia following resuscitation from cardiac arrest and in-hospital mortality. *JAMA* 2010;303(21):2165–2171
- 36 Xiao F, Arnold TC, Zhang S, et al. Cerebral cortical aquaporin-4 expression in brain edema following cardiac arrest in rats. *Acad Emerg Med* 2004;11(10):1001–1007
- 37 Elmer J, Torres C, Aufderheide TP, et al; Resuscitation Outcomes Consortium. Association of early withdrawal of life-sustaining therapy for perceived neurological prognosis with mortality after cardiac arrest. *Resuscitation* 2016;102:127–135
- 38 Cronberg T, Brizzi M, Liedholm LJ, et al. Neurological prognostication after cardiac arrest—recommendations from the Swedish Resuscitation Council. *Resuscitation* 2013;84(7):867–872
- 39 Wijdicks EF, Hijdra A, Young GB, Bassetti CL, Wiebe S; Quality Standards Subcommittee of the American Academy of Neurology. Practice parameter: prediction of outcome in comatose survivors after cardiopulmonary resuscitation (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2006;67(2):203–210
- 40 Sandroni C, Cariou A, Cavallaro F, et al. Prognostication in comatose survivors of cardiac arrest: an advisory statement from the European Resuscitation Council and the European Society of Intensive Care Medicine. *Intensive Care Med* 2014;40(12):1816–1831
- 41 Gold B, Puertas L, Davis SP, et al. Awakening after cardiac arrest and post resuscitation hypothermia: are we pulling the plug too early? *Resuscitation* 2014;85(2):211–214
- 42 Larsson E, Lindvall O, Kokaia Z. Stereological assessment of vulnerability of immunocytochemically identified striatal and hippocampal neurons after global cerebral ischemia in rats. *Brain Res* 2001;913(2):117–132
- 43 Björklund E, Lindberg E, Rundgren M, Cronberg T, Friberg H, Englund E. Ischaemic brain damage after cardiac arrest and induced hypothermia—a systematic description of selective eosinophilic neuronal death. A neuropathologic study of 23 patients. *Resuscitation* 2014;85(4):527–532
- 44 Tiainen M, Poutiainen E, Oksanen T, et al. Functional outcome, cognition and quality of life after out-of-hospital cardiac arrest and therapeutic hypothermia: data from a randomized controlled trial. *Scand J Trauma Resusc Emerg Med* 2015;23:12
- 45 Sabedra AR, Kristan J, Raina K, et al; Post Cardiac Arrest Service. Neurocognitive outcomes following successful resuscitation from cardiac arrest. *Resuscitation* 2015;90:67–72
- 46 Harper SJ, Wilkes RG. Posthypoxic myoclonus (the Lance-Adams syndrome) in the intensive care unit. *Anaesthesia* 1991;46(3):199–201
- 47 Welsh JP, Yuen G, Placantonakis DG, et al. Why do Purkinje cells die so easily after global brain ischemia? Aldolase C, EAAT4, and the cerebellar contribution to posthypoxic myoclonus. *Adv Neurol* 2002;89:331–359
- 48 Venkatesan A, Frucht S. Movement disorders after resuscitation from cardiac arrest. *Neurol Clin* 2006;24(1):123–132
- 49 Sieber FE, Palmon SC, Traystman RJ, Martin LJ. Global incomplete cerebral ischemia produces predominantly cortical neuronal injury. *Stroke* 1995;26(11):2091–2095, discussion 2096
- 50 Lilja G, Nielsen N, Friberg H, et al. Cognitive function in survivors of out-of-hospital cardiac arrest after target temperature management at 33°C versus 36°C. *Circulation* 2015;131(15):1340–1349
- 51 Hopkins RO, Weaver LK, Collingridge D, Parkinson RB, Chan KJ, Orme JF Jr. Two-year cognitive, emotional, and quality-of-life outcomes in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2005;171(4):340–347
- 52 Moolaert VR, van Heugten CM, Winkens B, et al. Early neurologically-focused follow-up after cardiac arrest improves quality of life at one year: A randomised controlled trial. *Int J Cardiol* 2015;193:8–16
- 53 Takahashi K, Sasanuma N, Itani Y, et al. Impact of early interventions by a cardiac rehabilitation team on the social rehabilitation of patients resuscitated from cardiogenic out-of-hospital cardiopulmonary arrest. *Intern Med* 2015;54(2):133–139
- 54 WHO. Towards a common language for function, disability, and health. International Classification of Functioning, Disability and Health (ICF). Geneva, Switzerland: World Health Organization; 2002
- 55 Rittenberger JC, Raina K, Holm MB, Kim YJ, Callaway CW. Association between Cerebral Performance Category, Modified Rankin Scale, and discharge disposition after cardiac arrest. *Resuscitation* 2011;82(8):1036–1040
- 56 Raina KD, Callaway C, Rittenberger JC, Holm MB. Neurological and functional status following cardiac arrest: method and tool utility. *Resuscitation* 2008;79(2):249–256
- 57 Raina KD, Rittenberger JC, Holm MB, Callaway CW. Functional outcomes: one year after a cardiac arrest. *BioMed Res Int* 2015;2015:283608
- 58 Moolaert VR, Verbunt JA, van Heugten CM, Wade DT. Cognitive impairments in survivors of out-of-hospital cardiac arrest: a systematic review. *Resuscitation* 2009;80(3):297–305
- 59 Cronberg T, Lilja G, Horn J, et al; TTM Trial Investigators. Neurologic function and health-related quality of life in patients following targeted temperature management at 33°C vs 36°C after out-of-hospital cardiac arrest: a randomized clinical trial. *JAMA Neurol* 2015;72(6):634–641
- 60 Phelps R, Dumas F, Maynard C, Silver J, Rea T. Cerebral Performance Category and long-term prognosis following out-of-hospital cardiac arrest. *Crit Care Med* 2013;41(5):1252–1257