Sports and Recreational Activities following Total Ankle Replacement

Federico Giuseppe Uselli¹ Andrea Pantalone² Camilla Maccario¹ Matteo Guelfi² Vincenzo Salini²

¹Casco, Foot and Ankle Unit, IRCCS Galeazzi, Milan, Italy

²Orthopaedic and Traumatology Division, Department of Medicine and Science of Aging, University G. d'Annunzio, Chieti-Pescara, Chieti, Italy

Address for correspondence Matteo Guelfi, MD, Orthopaedic and Traumatology Division, Department of Medicine and Science of Aging, University G. d'Annunzio, Chieti-Pescara, Via dei Vestini 35, 66013 Chieti, Italy (e-mail: matteoguel@hotmail.com).

Joints 2017;5:12–16.

Abstract

Purpose In literature, there is a controversy regarding whether patients who have undergone total ankle replacement (TAR) can participate in sports and recreational activities. The purpose of this study was to report change in sports activity level after TAR.

Methods A retrospective study was performed, enrolling 76 patients with symptomatic end-stage ankle arthritis who underwent TAR from May 2011 to October 2014. Patients were mainly males (44/76; 58%) and 56 years old on average (range: 22.3–79.6 years) at the time of surgery. They were treated with mobile-bearing prosthesis implanted with an anterior approach. Patients were evaluated preoperatively and 12 months postoperatively. Pain and function were assessed using the American Orthopaedic Foot and Ankle Society (AOFAS) ankle and hindfoot score, the visual analog scale (VAS) pain score, and the 12-Item Short Form Health Survey (SF-12) – physical component summary (PCS) and mental component summary (MCS). Activity level was assessed with the University of California at Los Angeles (UCLA) activity scale.

Results At 12 months postoperatively, statistically significant increase was reported for AOFAS scores (from 32.8 ± 12.7 preoperatively to 72.6 ± 13.3 ; $p < 0.001$), SF-12 PCS (from 34.3 ± 5.1 preoperatively to 45.4 ± 6.4 ; $p < 0.001$), and SF-12 MCS (from 39.8 ± 7.5 preoperatively to 51.4 ± 6.1 ; $p < 0.001$). A statistically significant decrease was detected in VAS pain score (from 8.7 ± 1.6 preoperatively to 2.2 ± 1.6 ; $p < 0.001$). The UCLA activity levels increased significantly from 2.4 ± 0.8 to 6.3 ± 2.3 ($p < 0.001$).

Conclusion Pain and function significantly improved in patients affected by ankle osteoarthritis, who underwent TAR, at 1-year follow-up. In addition, activity level showed a significant increase respect to preoperative condition.

Level of Evidence Level IV, retrospective case series.

Keywords

- ankle
- arthroplasty
- osteoarthritis
- replacement
- sport

Introduction

Ankle osteoarthritis (OA) incidence is significantly lower than knee and hip arthritis.¹ It is mostly posttraumatic (65–80% of the cases), and this may explain a different epidemiology with a younger population affected.² Historically, design issues are related to early failures of total ankle

replacement (TAR) rather than patient age or high-impact activities. Furthermore, it has been showed that TAR is a high demanding surgery with a long learning curve. This may contribute to raise the rate of failures in small cohorts, rather than patient age and high-impact activities. In the past, ankle arthrodesis was considered the first surgical treatment for end-stage arthritis of the ankle. However, during the years,

some important disadvantages related to this surgery became clear: compensatory overload, gait change, high rates of nonunion, long rehabilitation period, and eventual development of adjacent joint OA.³ These led to a new increasing interest in TAR, inducing important development in implants design and materials.

Mobile-bearing implants represented a revolutionary and successful step forward in Europe. With their consistent use, it has been clarified that primary implant stability should be the main focus of the surgeon rather than any cement fixation. Kofoed and Lundberg-Jensen⁴ evaluated early generation cemented and uncemented mobile-bearing TAR and concluded that the results of TAR were equivalent in patients younger and older than 50 years, pointing out cement fixation as a major reason for failures rather than patient age or high-impact activities. Recently, some authors proposed TAR as the gold standard treatment for ankle OA, even in young and active patients. Demetracopoulos et al⁵ claimed comparable results with TAR in younger (<55 years) and older patients (>70 years) at early follow-up. The incidences of complications and the revision rate were similar between the two groups. However, the authors did not describe physical activities allowed for these patients. There is still a controversy regarding whether patients who underwent TAR can participate in sport and recreational activities.⁶

Today, there is a new trend toward better results in terms of quality of life, and physical and athletic ability after TAR than after tibiotalar arthrodesis.⁷

The aim of this study was to analyze sports activity before and after TAR. The hypothesis of the study was that sports activity of patients affected by ankle OA increases after TAR.

Methods

This study is a retrospective review of 76 patients who were treated with Hintegra total ankle prosthesis (Newdeal, Lyon, France; Integra LifeSciences, Plainsboro, New Jersey, United States) between May 2011 and October 2014. All patients were followed up 12 months after surgery. The indications for symptomatic end-stage ankle OA were posttraumatic OA (70 patients), revision surgery of an osteochondral lesion procedure failure (one patient), posterior tibial tendon dysfunction (PTTD) stage IV according to the Bluman–Myerson⁸ classification (one patient), drop foot (one patient), and rheumatoid arthritis (three patients).

All patients, except for those with the rheumatoid arthritis and drop foot, had prior ankle surgeries: patients with posttraumatic OA previously had fixation of fractures and, in a few cases, removal of internal fixation, the patient with osteochondral lesion underwent two arthroscopic treatments performed by other surgeons, and the patient with PTTD was previously treated with subtalar and talonavicular arthrodesis.

All TAR procedures were performed by one of the co-authors according to the technique by Barg et al⁹: an anterior ankle approach of 10 to 12 cm length was made. The retinaculum was dissected along the lateral border of the

Fig. 1 Anterior ankle approach for total ankle replacement.

anterior tibial tendon, avoiding the neurovascular bundle that runs behind the long extensor hallucis tendon (► **Fig. 1**).

Patients were evaluated preoperatively and 12 months after surgery. Pain and function were assessed using the American Orthopaedic Foot and Ankle Society (AOFAS) ankle and hindfoot score, the visual analog scale (VAS) pain score, and the 12-Item Short Form Health Survey (SF-12) – physical component summary (PCS) and mental component summary (MCS). Furthermore, the patients were radiologically examined preoperatively and at follow-up using standard weight-bearing radiographs (anteroposterior and lateral views)¹⁰ and the Saltzman alignment view¹¹ (► **Fig. 2**).

Pre- and postoperative activity level was determined using the University of California at Los Angeles (UCLA) activity scale.¹² The UCLA scale is based on the patient's ranking of her/his level of physical activity. The score ranges from 1 to 10. The patient had to indicate her/his most appropriate activity level, with level 1 defined as “no physical activity, dependent on others” and level 10 defined as “regular participation in impact sports.”^{13,14}

All data were expressed as mean and standard deviation. Statistical analysis was performed with MATLAB statistical toolbox version 2008 (MathWorks, Natick, Massachusetts, United States). Paired Student's *t*-test was used to compare baseline and follow-up values. *P*-values less than 0.05 were considered statistically significant.

Results

A total of 76 patients were included in this review. Patients were mainly males (44/76; 58%), were 56 years old on average (range: 22.3–79.6 years) at the time of surgery, and were treated with Hintegra prosthesis (Allegra Orthopaedics, Lane Cove West, NSW, Australia) with an anterior approach.

Forty right and 36 left TARs were performed. The prosthesis components sizes were mismatched (talus component

Fig. 2 Painful end-stage varus osteoarthritis of the left ankle in a 54-year-old man. Preoperative standard weight-bearing X-rays in (A) anteroposterior and (B) lateral views. (C) Saltzman alignment view shows a significant varus malalignment of the heel. Postoperative radiographs in (D) anteroposterior, (E) lateral, and (F) Saltzman alignment view show a significant correction of the malalignment of the heel.

size smaller than tibial component size) in 32 patients. Nineteen patients underwent the following concurrent operative procedure at the time of TAR: Achilles tendon lengthening (seven patients), realignment calcaneal osteotomy (three patients), and subtalar arthrodesis (three patients). Ligament balancing was performed as required through an isolated deltoid release (one patient), a medial malleolar lengthening osteotomy (three patients), and a combination of medial malleolar lengthening osteotomy, fibular shortening osteotomy, and peroneus longus to brevis transfer (two patients).

At 12 months postoperatively, there was a statistically significant increase in AOFAS scores from 32.8 ± 12.7 preoperatively to 72.6 ± 13.3 ($p < 0.001$), in SF-12 PCS from 34.3 ± 5.1 preoperatively to 45.4 ± 6.4 ($p < 0.001$), and in the SF-12 MCS from 39.8 ± 7.5 preoperatively to 51.4 ± 6.1 ($p < 0.001$). In addition, there was also a statistically significant decrease in the VAS pain score from 8.7 ± 1.6 preoperatively to 2.2 ± 1.6 ($p < 0.001$). The UCLA activity levels increased significantly from 2.4 ± 0.8 to 6.3 ± 2.3 ($p < 0.001$) (–Table 1).

Before surgery, 11.7% of the patients were active in sports; after surgery, this percentage rose to 49.4%. The most frequent sports activities were jogging, dancing, biking, and skiing; a fraction of patients ($n = 14$) participated in impact sports such as jogging (13 patients) and martial arts (1 patient) in spite of adverse medical advice (–Table 2).

Table 1 UCLA score

	Pre-Operative		Post- TAR	
	Arthritic Condition			
UCLA Score	%	N° of Pts	%	N° of Pts
1	5,3	4	0	0
2	61,8	47	7,9	6
3	23,7	18	7,9	6
4	6,6	5	6,6	5
5	2,6	2	9,2	7
6	0	0	17,1	13
7	0	0	14,5	11
8	0	0	18,4	14
9	0	0	14,5	11
10	0	0	3,9	3
Total	100	76	100	76
Mean UCLA Score	2,4		6,3	

Discussion

Ankle OA has a deeply different epidemiology from hip and knee. These may be related to biomechanics reasons¹⁵: the loading area of the ankle is approximately one-third smaller

Table 2 Recommendations of sports activities

Recommended	Possible	Not recommended
Aerobic activity	Cross-country skiing	Basketball
Cycling	Sailing	Beach volleyball
Dancing	Alpine skiing	Climbing
Golf	Tennis	Horseback riding
Hiking or nordic walking	Hunting	Ice hockey
Swimming		Ice skating and skating
		Jogging
		Martial arts
		Mountain biking
		Paragliding and skydiving
		Rugby
		Soccer
		Volleyball
		Water ski
		Windsurfing

than the corresponding ones for knee or hip¹⁶; the thinner (1–2 mm) cartilage layer of the ankle offer a far more resilient tissue than those of the other joints.¹⁷ Posttraumatic changes are the main reasons for developing ankle OA (~65–80%).

Ankle trauma is more likely to affect younger population. This can also explain a younger population for ankle OA, which usually affect higher demanding patients if compared with hip and knee. In this respect, as a consequence of the impressive progress achieved in ankle prosthesis designs and clinical outcome following TAR, ankle arthrodesis is no longer the standard treatment for end-stage ankle OA.¹⁸

In our study, patient's recovery and joint restoration in terms of balance and movement suggest that TAR may be considered the golden standard at least for highly active patients. This statement finds a confirmation in our UCLA scale results. The UCLA activity score is described as a valid instrument for assessment of activity levels in patients undergoing total joint arthroplasties¹³: we detected a significant increase in UCLA activity levels from 2.4 to 6.3.

The most frequent sports activities were jogging, dancing, biking, and skiing; a group of patients¹⁴ participated in impact sports such as jogging¹³ and martial arts¹ in spite of adverse medical advice.

Despite of our early results, high-impact sports activities may increase polyethylene wear¹⁹ and the incidence of the fracture of polyethylene²⁰ at a long-term follow-up. In addition, young age, high body mass index, and poor bone quality are additional risk factors that led orthopaedic surgeons to remain more cautious and restrictive.⁶

Nevertheless, the first clinical study on sport and recreation activities in patients who underwent TAR published in

2006 found that the rate of sports participation increased from 36% preoperatively to 56% postoperatively.²¹ Furthermore, some authors pointed out that patients active in sports were not at higher risk for failure compared with other groups¹⁸; Naal et al reported that neither sports participation nor activity levels were identified as risk factors for development of periprosthetic radiolucencies.¹⁴

A critical factor in sport activities is the ankle range motion. Although the major advantages of TAR versus ankle arthrodesis is the preservation of ankle mobility, there is still no consensus whether patients with TAR have better results in sport activities than patients with ankle fusion. In fact, Schuh et al²² revealed no significant difference between these two groups concerning activity levels, participation in sports activities, and UCLA and AOFAS scores. Anyway, the patients after ankle arthrodesis significantly decreased their activities. It may be related to the loss of movement due to the degenerative changes in the ipsilateral, tarsometatarsal, and subtalar joints.^{23–25}

Recently, Flavin et al²⁶ demonstrated that patients with TAR showed higher speed during walk. In fact, the restored functionality of the ankle allowed a comparatively increased stride length and faster cadence, as well as more normalized first and second rockers at the gait analysis.

Bonnin et al²⁷ in their retrospective series of 140 mobile-bearing ankle arthroplasties showed that TAR improved the quality of life and that return to recreational activities was generally possible but the return to impact sport was rarely possible. As our results suggest, less pain and better range of motion can encourage patients to increase their participation in sports (including high impact ones) and recreation activities, but this observation should be checked at a longer follow-up.

Although TAR has become an increasingly popular procedure to treat symptomatic and recalcitrant ankle arthritis, orthopaedic surgeons have little guidance concerning physical activity and sport participation recommendations following this procedure.

Our results suggest, though preliminary, that TAR is possibly a better choice with respect to arthrodesis, at least for highly active individuals. This consideration comes from both patients' perception and, more objectively, the reduced stress imposed by TAR to the nearest joints with respect to OA.

To date, further studies on the durability and complications of prostheses, as well as patient activity and return to specific sports and activities after surgery, are needed to reach a unique guideline based on stronger levels of evidence.

References

- Huch K. Knee and ankle: human joints with different susceptibility to osteoarthritis reveal different cartilage cellularity and matrix synthesis in vitro. *Arch Orthop Trauma Surg* 2001;121(06):301–306
- Horisberger M, Valderrabano V, Hintermann B. Posttraumatic ankle osteoarthritis after ankle-related fractures. *J Orthop Trauma* 2009;23(01):60–67
- Boobbyer GN. The long-term results of ankle arthrodesis. *Acta Orthop Scand* 1981;52(01):107–110

- 4 Kofoed H, Lundberg-Jensen A. Ankle arthroplasty in patients younger and older than 50 years: a prospective series with long-term follow-up. *Foot Ankle Int* 1999;20(08):501–506
- 5 Demetracopoulos CA, Adams SB Jr, Queen RM, DeOrio JK, Nunley JA II, Easley ME. Effect of age on outcomes in total ankle arthroplasty. *Foot Ankle Int* 2015;36(08):871–880
- 6 Macaulay AA, VanValkenburg SM, DiGiovanni CW. Sport and activity restrictions following total ankle replacement: a survey of orthopaedic foot and ankle specialists. *Foot Ankle Surg* 2015; 21(04):260–265
- 7 Dalat F, Trouillet F, Fessy MH, Bourdin M, Besse JL. Comparison of quality of life following total ankle arthroplasty and ankle arthrodesis: retrospective study of 54 cases. *Orthop Traumatol Surg Res* 2014;100(07):761–766
- 8 Bluman EM, Title CI, Myerson MS. Posterior tibial tendon rupture: a refined classification system. *Foot Ankle Clin* 2007;12(02): 233–249, v
- 9 Barg A, Knupp M, Henninger HB, Zwicky L, Hintermann B. Total ankle replacement using HINTEGRA, an unconstrained, three-component system: surgical technique and pitfalls. *Foot Ankle Clin* 2012;17(04):607–635
- 10 Hintermann B, Valderrabano V, Dereymaeker G, Dick W. The HINTEGRA ankle: rationale and short-term results of 122 consecutive ankles. *Clin Orthop Relat Res* 2004;(424):57–68
- 11 Saltzman CL, el-Khoury GY. The hindfoot alignment view. *Foot Ankle Int* 1995;16(09):572–576
- 12 Zehiri CA, Schmalzried TP, Szuszczewicz ES, Amstutz HC. Assessing activity in joint replacement patients. *J Arthroplasty* 1998; 13(08):890–895
- 13 Naal FD, Impellizzeri FM, Leunig M. Which is the best activity rating scale for patients undergoing total joint arthroplasty? *Clin Orthop Relat Res* 2009;467(04):958–965
- 14 Naal FD, Impellizzeri FM, Loibl M, Huber M, Rippstein PF. Habitual physical activity and sports participation after total ankle arthroplasty. *Am J Sports Med* 2009;37(01):95–102
- 15 Labib SA, Raikin SM, Lau JT, et al. Joint preservation procedures for ankle arthritis. *Foot Ankle Int* 2013;34(07):1040–1047
- 16 Kimizuka M, Kurosawa H, Fukubayashi T. Load-bearing pattern of the ankle joint. Contact area and pressure distribution. *Arch Orthop Trauma Surg* 1980;96(01):45–49
- 17 Thomas RH, Daniels TR. Ankle arthritis. *J Bone Joint Surg Am* 2003;85-A(05):923–936
- 18 Hörterer H, Miltner O, Müller-Rath R, Phisitkul P, Barg A. Sports activity in patients with total ankle replacement. *Sport-Orthop-Traumatol* 2015;31:34–40
- 19 Harris NJ, Brooke BT, Sturdee S. A wear debris cyst following S.T.A. R. Total Ankle Replacement—surgical management. *Foot Ankle Surg* 2009;15(01):43–45
- 20 Scott AT, Nunley JA. Polyethylene fracture following STAR ankle arthroplasty: a report of three cases. *Foot Ankle Int* 2009;30(04): 375–379
- 21 Valderrabano V, Pagenstert G, Horisberger M, Knupp M, Hintermann B. Sports and recreation activity of ankle arthritis patients before and after total ankle replacement. *Am J Sports Med* 2006; 34(06):993–999
- 22 Schuh R, Hofstaetter J, Krismer M, Bevon R, Windhager R, Trnka HJ. Total ankle arthroplasty versus ankle arthrodesis. Comparison of sports, recreational activities and functional outcome. *Int Orthop* 2012;36(06):1207–1214
- 23 Glazebrook M. End-stage ankle arthritis: magnitude of the problem and solutions. *Instr Course Lect* 2010;59: 359–365
- 24 Fuchs S, Sandmann C, Skwara A, Chylarecki C. Quality of life 20 years after arthrodesis of the ankle. A study of adjacent joints. *J Bone Joint Surg Br* 2003;85(07):994–998
- 25 Buchner M, Sabo D. Ankle fusion attributable to posttraumatic arthrosis: a long-term followup of 48 patients. *Clin Orthop Relat Res* 2003;(406):155–164
- 26 Flavin R, Coleman SC, Tenenbaum S, Brodsky JW. Comparison of gait after total ankle arthroplasty and ankle arthrodesis. *Foot Ankle Int* 2013;34(10):1340–1348
- 27 Bonnin MP, Laurent JR, Casillas M. Ankle function and sports activity after total ankle arthroplasty. *Foot Ankle Int* 2009;30(10): 933–944