Editorial

Progressive Resistance Training as Complementary Therapy for Polycystic Ovarian Syndrome

Treinamento de Resistência Progressiva como Terapia Complementar Para Síndrome de Ovário Policístico

Gislaine Satyko Kogure¹ Rosana Maria dos Reis¹

¹ Department of Gynecology and Obstetrics, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, Brazil

Rev Bras Ginecol Obstet 2017;39:255–257.

Polycystic ovarian syndrome (PCOS) significantly impacts women, since the broad spectrum of clinical manifestations associated with it are significant and include reproductive dysfunction, menstrual irregularities, and an increased risk of infertility. However, the consequences of PCOS go beyond the reproductive axis, with psychological and social impairments, including stress, depression, anxiety, and sexual dissatisfaction.¹ There is also a high prevalence of dyslipidemia, hyperinsulinemia, obesity, hypertension, and glucose intolerance, which are risk factors that predispose women to cardiovascular disease (CVD) and diabetes mellitus type 2 (DM2).² Therefore, PCOS assumes aspects of a chronic disease, as these factors extend throughout life. If not prevented and treated, they can lead to increased morbidity and mortality.³ The etiology of PCOS has not been fully elucidated, but it is known to be linked to excess androgens.^{4,5} Insulin resistance (IR) is a common feature of PCOS,⁶ and although it is not considered a diagnostic criterion, it is a key factor in the syndrome's etiology and evolution.⁷

In 2008, the European Society of Human Reproduction and Embryology (ESHRE) and the American Society for Reproductive Medicine (ASRM) published a consensus suggesting lifestyle changes as the first line of treatment for women with PCOS.⁸ This consensus was reinforced in 2009 by The Androgen Excess and Polycystic Ovary Syndrome Society,⁹ which investigated evidence of lifestyle management (dietary, exercise, or behavioral interventions) for obesity in women with PCOS. Since then, the benefits of lifestyle changes resulting from PCOS therapy have been well documented.^{10,11} The rationale for this non-pharmacological therapy is based on regular exercise and a healthy diet, as well as combined interventions that aim to

achieve and maintain a healthy weight to minimize hormonal and reproductive complications, reduce the long-term risks of chronic diseases such as CVD and DM2, and consequently improve quality of life. In this way, weight loss has been considered the main goal of PCOS therapy in obese women.

Moderate- to high-intensity aerobic physical exercise¹² has been predominantly recommended as a treatment for PCOS.¹³ Preliminary data from our group with an interval aerobic training protocol (exercises alternating in intensity from moderate to heavy effort with low-effort recovery periods) showed a decreased central obesity index measured by anthropometric measures and improved testosterone levels. However, other training programs, such as aerobic exercises, alone or in combination with resistance training, with or without dietary restriction, have also effectively reduced total and abdominal body fat or body fat percentile,^{12,14–16} leading to improved menstrual frequency and/or ovulation,^{12,17} reduced serum testosterone concentrations and fasting plasma glucose levels,¹² and improved insulin sensitivity.^{18,19}

More recently, a review presented evidence that progressive resistance training (PRT), or strength training, may also be beneficial for women with PCOS, promoting changes in body composition and associated factors, especially IR.²⁰ However, this type of physical exercise has not been well explored in terms of its therapeutic purposes. Faced with the phenotypic characteristics of PCOS, we set out to perform a periodic protocol of resistance exercises in lean, overweight, and obese women with the intention of evaluating the results of this therapy. The PRT improved hyperandrogenism and the menstrual cycle, as well as the functional capacity with increased muscle strength, and resulted in changes in body

Address for correspondence
Rosana Maria dos Reis, PhD, MD,
Department of ObGyn, Ribeirão
Preto Medical School,
Universidade de São Paulo,
Av. Bandeirantes, 3900,
14049-900 - Ribeirão Preto, SP, Brazil
(e-mail: romareis@fmrp.usp.br).

DOI <https://doi.org/10.1055/s-0037-1602705>.
ISSN 0100-7203.

Copyright © 2017 by Thieme Revinter
Publicações Ltda, Rio de Janeiro, Brazil

License terms

composition with increased lean muscle mass and decreased central obesity, without a reduced total weight.²¹ There were also improvements in quality of life and sexual function as complementary benefits.^{22,23}

Evidence in the literature suggests that a PRT protocol associated with the practice of calisthenics (exercises that use the body's own weight as the primary tool) promoted significant changes in body composition, even with the increase of total weight and lean muscle strength in overweight and obese individuals. It also promoted improvements in several health-related domains, such as anxiety, depression, and quality of life.²⁴ There was also a significant improvement in body composition without reductions in the total weight of lean women, regardless of the type of training when high-intensity interval aerobic physical exercise and PRT were compared.²⁵ These studies are pioneers in the evaluation of PRT alone, in women with PCOS, and attest that strength training can be an excellent and effective exercise option.

The improvement in body composition, with little or no effective change in total body weight, may be related to the concomitance of the increase in lean muscle mass and reduced body fat promoted by this type of exercise.²⁶ The loss of fat mass is probably mediated by an increase in the basal metabolic rate that results from an increase in lean muscle mass, which is considered a metabolically active tissue that causes the body to increase its caloric expenditure.^{27,28} This anabolic action adds to the improved insulin sensitivity,^{27,29,30} since the skeletal muscles are predominantly involved in insulin-mediated glucose uptake,^{31,32} while muscle contractile activity may stimulate the translocation of glucose transporter type 4 (GLUT4) molecules in the absence of insulin.^{33,34} This is a key consideration for women with PCOS, since IR is implicated in the etiology of the disease.^{6,7}

The obesity that affects most women with PCOS³⁵ is associated with reduced muscle strength, difficulties in postural control, and changes in the biomechanical behavior of the lower limbs,³⁶ and may be a limiting factor for some physical activities. In particular, those physical activities with a great cyclic impact, such as walking, running, and those that require great joint amplitude are affected by obesity, since the excess weight alone puts a significant amount of stress on the joints. In addition, rapid exhaustion due to reduced physical fitness and functional capacity, even with low physical effort, contributes to the non-compliance with this type of exercise. In addition to the aforementioned benefits, as a therapy for obese women, PRT improves daily functional capacity, increases resistance to joint impact, promotes muscle strengthening, reduces the risk of injury, and favors subsequent aerobic exercises³⁷ within the recommended levels.¹²

Intervention programs with aerobic training or strength training performed exclusively induce favorable adaptations in women with PCOS. Although aerobic exercises are more highly recommended, a consistent training protocol including aerobic and strength exercises, either in the same session or on alternate days, can be both efficient and capable of improving the variables of the components of physical fitness related to health, such as muscle strength, and of preventing loss of lean mass. This protocol can also improve

disease-related characteristics, such as central obesity, hyperandrogenism, and insulin sensitivity.

Physical performance depends not only on the factors inherent to the suggested training program, but also on the degree of motivation for certain activities. With the proven effectiveness of the different physical training modalities, the possibility that it will provide personal satisfaction is increased, which can promote better adherence to the training program. Evidently, the positive effects of exercise may vary significantly among lifestyles as well as exercise program levels, such as intensity, frequency, and duration.³⁸ These should be prescribed individually by a physical educator with a focus on the expectations and motivations of women with PCOS and, above all, the safety of the proposed exercises. It is believed that therapeutic orientations based on non-pharmacological therapy may favor behavioral changes and the adoption of healthy lifestyle habits for women with PCOS.

References

- Himelein MJ, Thatcher SS. Polycystic ovary syndrome and mental health: A review. *Obstet Gynecol Surv* 2006;61(11):723-732
- American Association of Clinical Endocrinologists Polycystic Ovary Syndrome Writing Committee. American Association of Clinical Endocrinologists position statement on metabolic and cardiovascular consequences of polycystic ovary syndrome. *Endocr Pract* 2005;11(02):126-134
- Azziz R, Carmina E, Dewailly D, et al; Androgen Excess Society. Positions statement: criteria for defining polycystic ovary syndrome as a predominantly hyperandrogenic syndrome: An Androgen Excess Society guideline. *J Clin Endocrinol Metab* 2006; 91(11):4237-4245
- Barth JH, Yasmin E, Balen AH. The diagnosis of polycystic ovary syndrome: the criteria are insufficiently robust for clinical research. *Clin Endocrinol (Oxf)* 2007;67(06):811-815
- Svendsen PF, Nilas L, Nørgaard K, Madsbad S. [Polycystic ovary syndrome. New pathophysiological discoveries—therapeutic consequences]. *Ugeskr Laeger* 2005;167(34):3147-3151 Danish
- Stepto NK, Cassar S, Joham AE, et al. Women with polycystic ovary syndrome have intrinsic insulin resistance on euglycaemic-hyperinsulinaemic clamp. *Hum Reprod* 2013;28(03):777-784
- Diamanti-Kandaraki E, Papavassiliou AG. Molecular mechanisms of insulin resistance in polycystic ovary syndrome. *Trends Mol Med* 2006;12(07):324-332
- Thessaloniki ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Consensus on infertility treatment related to polycystic ovary syndrome. *Hum Reprod* 2008;23(03):462-477
- Moran LJ, Pasquali R, Teede HJ, Hoeger KM, Norman RJ. Treatment of obesity in polycystic ovary syndrome: a position statement of the Androgen Excess and Polycystic Ovary Syndrome Society. *Fertil Steril* 2009;92(06):1966-1982
- Haqq L, McFarlane J, Dieberg G, Smart N. Effect of lifestyle intervention on the reproductive endocrine profile in women with polycystic ovarian syndrome: a systematic review and meta-analysis. *Endocr Connect* 2014;3(01):36-46
- Moran LJ, Hutchison SK, Norman RJ, Teede HJ. Lifestyle changes in women with polycystic ovary syndrome. *Cochrane Database Syst Rev* 2011;(02):CD007506
- Thomson RL, Buckley JD, Noakes M, Clifton PM, Norman RJ, Brinkworth GD. The effect of a hypocaloric diet with and without exercise training on body composition, cardiometabolic risk profile, and reproductive function in overweight and obese women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2008;93(09):3373-3380

- 13 Teede HJ, Misso ML, Deeks AA, et al; Guideline Development Groups. Assessment and management of polycystic ovary syndrome: summary of an evidence-based guideline. *Med J Aust* 2011;195(06):S65–S112
- 14 Huber-Buchholz MM, Carey DG, Norman RJ. Restoration of reproductive potential by lifestyle modification in obese polycystic ovary syndrome: role of insulin sensitivity and luteinizing hormone. *J Clin Endocrinol Metab* 1999;84(04):1470–1474
- 15 Thomson RL, Brinkworth GD, Noakes M, Clifton PM, Norman RJ, Buckley JD. The effect of diet and exercise on markers of endothelial function in overweight and obese women with polycystic ovary syndrome. *Hum Reprod* 2012;27(07):2169–2176
- 16 Thomson RL, Buckley JD, Moran LJ, et al. Comparison of aerobic exercise capacity and muscle strength in overweight women with and without polycystic ovary syndrome. *BJOG* 2009;116(09):1242–1250
- 17 Aubuchon M, Laughbaum N, Poetker A, Williams D, Thomas M. Supervised short-term nutrition and exercise promotes weight loss in overweight and obese patients with polycystic ovary syndrome. *Fertil Steril* 2009;91(4, Suppl):1336–1338
- 18 Harrison CL, Stepto NK, Hutchison SK, Teede HJ. The impact of intensified exercise training on insulin resistance and fitness in overweight and obese women with and without polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2012;76(03):351–357
- 19 Hutchison SK, Stepto NK, Harrison CL, Moran LJ, Strauss BJ, Teede HJ. Effects of exercise on insulin resistance and body composition in overweight and obese women with and without polycystic ovary syndrome. *J Clin Endocrinol Metab* 2011;96(01):E48–E56
- 20 Cheema BS, Vizza L, Swaraj S. Progressive resistance training in polycystic ovary syndrome: can pumping iron improve clinical outcomes? *Sports Med* 2014;44(09):1197–1207
- 21 Kogure GS, Miranda-Furtado CL, Silva RC, et al. Resistance exercise impacts lean muscle mass in women with polycystic ovary syndrome. *Med Sci Sports Exerc* 2016;48(04):589–598
- 22 Lara LA, Ramos FK, Kogure GS, et al. Impact of physical resistance training on the sexual function of women with polycystic ovary syndrome. *J Sex Med* 2015;12(07):1584–1590
- 23 Ramos FK, Lara LA, Kogure GS, et al. Quality of life in women with polycystic ovary syndrome after a program of resistance exercise training. *Rev Bras Ginecol Obstet* 2016;38(07):340–347
- 24 Vizza L, Smith CA, Swaraj S, Agho K, Cheema BS. The feasibility of progressive resistance training in women with polycystic ovary syndrome: a pilot randomized controlled trial. *BMC Sports Sci Med Rehabil* 2016;8:14
- 25 Almenning I, Rieber-Mohn A, Lundgren KM, Shetelig Løvvik T, Garnæs KK, Moholdt T. Effects of high intensity interval training and strength training on metabolic, cardiovascular and hormonal outcomes in women with polycystic ovary syndrome: a pilot study. *PLoS One* 2015;10(09):e0138793
- 26 Hunter GR, Bryan DR, Wetzstein CJ, Zuckerman PA, Bamman MM. Resistance training and intra-abdominal adipose tissue in older men and women. *Med Sci Sports Exerc* 2002;34(06):1023–1028
- 27 Treserras MA, Balady GJ. Resistance training in the treatment of diabetes and obesity: mechanisms and outcomes. *J Cardiopulm Rehabil Prev* 2009;29(02):67–75
- 28 Strasser B, Siebert U, Schobersberger W. Resistance training in the treatment of the metabolic syndrome: a systematic review and meta-analysis of the effect of resistance training on metabolic clustering in patients with abnormal glucose metabolism. *Sports Med* 2010;40(05):397–415
- 29 Holten MK, Zacho M, Gaster M, Juel C, Wojtaszewski JF, Dela F. Strength training increases insulin-mediated glucose uptake, GLUT4 content, and insulin signaling in skeletal muscle in patients with type 2 diabetes. *Diabetes* 2004;53(02):294–305
- 30 Strasser B, Schobersberger W. Evidence for resistance training as a treatment therapy in obesity. *J Obes* 2011;2011:482564
- 31 DeFronzo RA, Jacot E, Jequier E, Maeder E, Wahren J, Felber JP. The effect of insulin on the disposal of intravenous glucose. Results from indirect calorimetry and hepatic and femoral venous catheterization. *Diabetes* 1981;30(12):1000–1007
- 32 Smith AG, Muscat GE. Skeletal muscle and nuclear hormone receptors: implications for cardiovascular and metabolic disease. *Int J Biochem Cell Biol* 2005;37(10):2047–2063
- 33 Hayashi T, Hirshman MF, Fujii N, Habinowski SA, Witters LA, Goodyear LJ. Metabolic stress and altered glucose transport: activation of AMP-activated protein kinase as a unifying coupling mechanism. *Diabetes* 2000;49(04):527–531
- 34 Goodyear LJ, Kahn BB. Exercise, glucose transport, and insulin sensitivity. *Annu Rev Med* 1998;49:235–261
- 35 Vrbikova J, Hainer V. Obesity and polycystic ovary syndrome. *Obes Facts* 2009;2(01):26–35
- 36 Wearing SC, Hennig EM, Byrne NM, Steele JR, Hills AP. The biomechanics of restricted movement in adult obesity. *Obes Rev* 2006;7(01):13–24
- 37 Ramos AT. Atividade física: diabéticos, gestantes, 3ª idade, crianças, obesos. São Paulo: Sprint; 1996
- 38 Marx JO, Ratamess NA, Nindl BC, et al. Low-volume circuit versus high-volume periodized resistance training in women. *Med Sci Sports Exerc* 2001;33(04):635–643