Lumbar Spinal Hydatid Cyst with Intracanal and Paravertebral Invasion: A Rare Entity in an Unusual Location

Elias Antoniadēs¹ Ioannis Magras¹ Konstantinos Sapalidis²

¹ Department of Neurosurgery, AHEPA Hospital, Aristotle University of Thessaloniki, Thessaloniki, Greece

² Department of General Surgery, AHEPA Hospital, Aristotle University of Thessaloniki, Thessaloniki, 54621 Greece

Address for correspondence Elias Antoniadēs, MD, Department of Neurosurgery, AHEPA Hospital, Aristotle University of Thessaloniki, Thessaloniki, 54621, Greece (e-mail: eliasantoniad@yahoo.gr).

Indian J Neurosurg 2017;6:151–152.

Hydatid disease is a parasitic infection associated with *Echinococcus granulosus* tapeworm. The spine may be affected in the context of hematogenous dissemination in 1% of the cases.¹ According to its manifestation, hydatid disease is categorized as

1. Intramedullary
2. Extradural
3. Intradural extramedullary
4. Hydatid disease of the vertebra
5. Paravertebral²

An intradural disease without vertebral column involvement is extremely rare.² Thoracolumbar spine (63.9%) is the predominantly affected region, as *Echinococcus* typically inserts vascular segments of the spine.³

We aim to present an uncommon case of lumbar manifestation of hydatid disease, reviewing the literature as well.

A 65-year-old patient presented with lumboischialgia, progressive paraparesis 3/5 of his right leg and 2/5 of his left one accompanied by reflexes absence, hypaesthesia, and vegetative disorders. Symptoms were initially attributed to lumbar stenosis with unsuccessful conservative treatment having no amelioration by inclination. Plain radiographs of the lumbar spine were uneventful. Magnetic resonance imaging (MRI) of the lumbar spine revealed a multilobulated cystic formation with spatial dimensions 10 cm × 4 cm between the second and fifth lumbar spine extending intracanalally and compressing the dural sac leftward. The rest MRI neuraxis control was unremarkable. Computer tomographic (CT) scans of the chest and abdomen revealed no infiltration (► Fig. 1). The labor control revealed no eosinophilia, whereas indirect hemagglutination test for *Echinococcus* was positive in 1/256 dilution. Finally the echocardiogram was unremarkable too.

With a raised suspicion of hydatid cyst, the surgical excision was warranted.

We performed a posterior laminectomy with resection of the intracanal cysts. Postoperative pain symptoms were relieved and muscle strengths restituted to 4/5 right and 3/5 left, respectively. Hypaesthesia also retreated but unfortunately impaired micturition and defecation remained. Postoperative MRI control showed total resection. Albendazole was also administered for the following 4 months.

The mechanism of intracanalally parasites seeding is attributed to hematogenic dissemination.⁴ The mechanism is imputed to the unique feature of vertebral plexus. Groen et al⁵ have divided it in their studies in three parts: the internal vertebral plexus surrounding the dura matter, the basivertebral veins of vertebral bodies, and the external vertebral plexus surrounding the column. The internal vertebral plexus communicates with the intraspinal and radicular veins that constitute valves, thus permitting a reflux and explaining the primary presence of hydatid cyst in the spine.

Symptoms of spinal disease usually manifest insidiously years after the primary infestation of cysts. Slow development of cysts over several years and the surrounding bone resistance support a rationale for this. Most patients refer a progressive low back pain with radiculopathy.⁶

MRI remains the diagnostic modality of choice. Findings suggestive of hydatid cyst on T1-weighted images include the isointense signal and hyperintensity on T2-weighted images.³

By deteriorating neurologic dysfunction, posterior laminectomy with intact cyst enucleation remains the horseshoe, which should also be combined with adjuvant anthelmintic therapy.⁷

Hanci et al⁸ and Prabhakar et al⁹ have also presented cases of lumbar spine hydatid cysts with deteriorating neurologic deficits treated with laminectomy and anthelmintic therapy with recession of deficits after decompression.

received
May 4, 2017
accepted
July 3, 2017
published online
August 29, 2017

DOI <https://doi.org/10.1055/s-0037-1606309>.
ISSN 2277-954X.

© 2017 Neurological Surgeons' Society of India

License terms

Fig. 1 (a) Preoperative sagittal sequences of MRI of lumbar spine, (b) axial sequences of the same examination, and (c) postoperative MRI T2-sagittal sequences of lumbar spine.

The unique features in this case were the initial deluding clinical and radiologic combination of findings regarding the extension of the lesion, extended intracranially and paravertebrally without significant vertebral body infiltration, and concomitant infection signs. Thus diagnostic suspicion should be raised by a cystic lesion type I or II according to Nabors' classification¹⁰ concerning hydatid disease as a rare but possible diagnosis.

Note

The patient has given his informed consent for publishing this case report.

Funding

None.

Conflict of Interest

None.

References

- 1 Fares Y, Khazim R, El Zaatari MM, Haddad GF, Barnes PR. Spinal hydatid disease and its neurological complications. *Scand J Infect Dis* 2003;35(6-7):394-396
- 2 Braithwaite PA, Lees RF. Vertebral hydatid disease: radiological assessment. *Radiology* 1981;140(03):763-766
- 3 Kafaji A, Al-Zain T, Lemcke J, Al-Zain F. Spinal manifestation of hydatid disease: a case series of 36 patients. *World Neurosurg* 2013;80(05):620-626
- 4 Kaen A, Lagares A, Perez-Nuñez A, Rivas JJ, Ramos A, Lobato RD. Intradural extramedullary spinal hydatidosis: case report. *Neurocirugia (Astur)* 2009;20(03):282-287
- 5 Groen RJ, du Toit DF, Phillips FM, et al. Anatomical and pathological considerations in percutaneous vertebroplasty and kyphoplasty: a reappraisal of the vertebral venous system. *Spine* 2004;29(13):1465-1471
- 6 Salduz A, Koyuncu LO, Dikici F, Talu U. [Long-term result of treatment for paraspinal and extradural hydatid cyst: a case report] [in Turkish]. *Acta Orthop Traumatol Turc* 2009;43(03):267-271
- 7 Lam KS, Faraj A, Mulholland RC, Finch RG. Medical decompression of vertebral hydatidosis. *Spine* 1997;22(17):2050-2055
- 8 Hanci M, Aydingoz O, Tanriverdi T, Kocer N. Epidural hydatid cyst of the lumbar spine. A case report. *Turkish J Spinal Surg* 2001;12:38-43
- 9 Prabhakar MM, Acharya AJ, Modi DR, Jadav B. Spinal hydatid disease: a case series. *J Spinal Cord Med* 2005;28(05):426-431
- 10 Nabors MW, Pait TG, Byrd EB, et al. Updated assessment and current classification of spinal meningeal cysts. *J Neurosurg* 1988;68(03):366-377