Intralesional Endoscopy and Septectomy as a Diagnostic Tool and Treatment Method for Lymphatic Malformations

Anne-Sophie Holler¹ Jan Gödeke¹ Veronika Engel¹ Oliver J. Muensterer¹

¹Department of Pediatric Surgery, University Medicine of the Johannes Gutenberg University Mainz, Mainz, Germany

Eur J Pediatr Surg Rep 2017;5:e60–e61.

Address for correspondence Anne-Sophie Holler, MD, Department of Pediatric Surgery, University Medicine of the Johannes Gutenberg University Mainz, Langenbeckstr. 1, Mainz 55131, Germany (e-mail: anne-sophie.holler@unimedizin-mainz.de).

New Insights and the Importance for the Pediatric Surgeon

Current treatment options for lymphatic malformations are expectant management, sclerotherapy, and surgical resection. Intralesional endoscopy and intercystic septectomy constitute a promising supplementary treatment method for subcutaneous lymphatic malformations. By creating a single, communicating cavity, the efficacy of subsequent sclerotherapy may be increased, and thus the need for multiple treatment sessions may be avoidable.

Case Report

Sclerotherapy and surgery are both effective treatment methods for lymphatic malformations.^{1,2} However, recurrence due to incomplete resection is a common problem, often necessitating multiple treatment sessions.² Intralesional

endoscopy has been described as a diagnostic approach and potential therapeutic tool.^{1,3}

We report a case of a 12-year-old male patient who presented with a mainly subcutaneous mixed lymphatic malformation located on the right flank (→Fig. 1). After suffering a direct trauma, the lesion had increased markedly in size and surgical intervention was indicated due to the associated pain. Intralesional endoscopy was performed that showed a mixed macro-/microcystic lymphatic malformation with hemorrhage (→Fig. 2). Intercystic septa were dissected under endo-

Fig. 1 Coronal T2 magnetic resonance imaging (MRI) with a subcutaneous mixed macro-/microcystic lymphatic malformation on the patient's right flank (white arrow).

Fig. 2 Intraoperative image showing a macrocystic cavity and many small fluid-filled cysts (asterisks), which are separated by thin septae.

received
January 3, 2017
accepted after revision
February 23, 2017

DOI <https://doi.org/10.1055/s-0037-1606388>.
ISSN 2194-7619.

© 2017 Georg Thieme Verlag KG
Stuttgart · New York

License terms

scopic visualization. At the end of the procedure, a single macrocystic cavity had been artificially created. Picibanil (OK-432) was inserted into the cavity and left in situ for 24 hours (► **Video 1**). The patient had no visible swelling, no pain, and merely two small, well-healed scars at 2 months of follow-up.

Video 1

Under sonographic guidance, the largest cyst of the lymphatic malformation was cannulated, and an 8 French pigtail catheter was advanced into the cyst over a guidewire. The cyst was filled with 0.9% sodium chloride solution to facilitate the placement of two 3-mm trocars into the cyst. Intralesional endoscopy showed a mixed macro-/microcystic lymphatic malformation with hemorrhage. Dissection of intercystic septa was performed under endoscopic visualization using hook electrocautery and blunt dissection. At the end of the procedure, a single macrocystic cavity had been artificially created. Picibanil (OK-432) was instilled through the formerly inserted pigtail catheter into the now solitary cyst and left in situ for 24 hours. Online content including video sequences viewable at: www.thieme-connect.com/products/ejournals/html/10.1055/s-0037-1606388.

Intralesional endoscopy and intercystic septectomy constitute an interesting novel approach for the diagnosis and treatment of mixed lymphatic malformations. By creating a single, communicating cavity, the efficacy of subsequent sclerotherapy may be increased, and thus the need for multiple treatment sessions may be avoidable.

Conflict of Interest

None.

References

- 1 Defnet AM, Bagrodia N, Hernandez SL, Gwilliam N, Kandel JJ. Pediatric lymphatic malformations: evolving understanding and therapeutic options. *Pediatr Surg Int* 2016;32(05):425–433
- 2 Cheng J. Doxycycline sclerotherapy in children with head and neck lymphatic malformations. *J Pediatr Surg* 2015;50(12):2143–2146
- 3 Eivazi B, Teymoortash A, Wiegand S, et al. Intralesional endoscopy of advanced lymphatic malformations of the head and neck: a new diagnostic approach and a potential therapeutic tool. *Arch Otolaryngol Head Neck Surg* 2010;136(08):790–795