L. QIU, R. MCCAFFREY, Y. JIN, Y. GONG, Y. HU, H. SUN, W. PARK, W. ZHANG* (YUNNAN UNIVERSITY, KUNMING, P. R. OF CHINA AND UNIVERSITY OF COLORADO AT BOULDER, USA)

Cage-Templated Synthesis of Highly Stable Palladium Nanoparticles and their Catalytic Activities in Suzuki–Miyaura Coupling

Chem. Sci. **2018**, *9*, 676–680.

Suzuki–Miyaura Reaction Catalyzed by Cage-Templated Palladium Nanoparticles

Category

Polymer-Supported Synthesis

Key words

palladium catalysis

nanoparticles

Suzuki–Miyaura coupling

biaryls

Synfact
of the month

Significance: A palladium nanocatalyst (PdNP@1) prepared by a cage-templated synthesis method (eq. 1), catalyzed the Suzuki–Miyaura coupling of phenylboronic acid with iodo- or bromobenzenes under microwave irradiation to give the corresponding biaryl products in 96–99% yield (eq. 2).

SYNFACTS Contributors: Yasuhiro Uozumi, Anggi Eka Putra
Synfacts 2018, 14(04), 0431 Published online: 16.03.2018
DOI: 10.1055/s-0037-1609379; Reg-No.: Y02018SF

Comment: PdNP@1 was characterized by means of UV–Vis, ^1H NMR (DOSY), HR-TEM, and EDS analyses. In the reaction of 4-iodotoluene with phenylboronic acid, the catalyst was reused three times without loss of its catalytic activity. The reactivity of PdNP@1 was higher than that of commercially available palladium catalysts such as $\text{Pd}(\text{PPh}_3)_4$ or Pd/C .