M. W. GRIBBLE, JR, S. GUO, S. L. BUCHWALD* (MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, USA)

Asymmetric Cu-Catalyzed 1,4-Dearomatization of Pyridines and Pyridazines without Preactivation of the Heterocycle or Nucleophile

J. Am. Chem. Soc. **2018**, *140*, 5057–5060.

Enantioselective Dearomatization of Pyridines with Styrene-Derived Nucleophiles

Selected examples:

Significance: Dearomatization of heteroaromatics is a valuable transformation in organic chemistry to create new carbon–carbon bonds. Herein, Buchwald and co-workers report the dearomatization of pyridines with a chiral benzylic copper species generated through an enantioselective copper hydride addition to the corresponding styrene.

SYNFACTS Contributors: Mark Lautens, Andrew Whyte
Synfacts 2018, 14(07), 0707 Published online: 18.06.2018
DOI: 10.1055/s-0037-1609533; Reg-No.: L06218SF

Comment: The intermediate dihydropyridine could be oxidized to regenerate the pyridine or reduced to give the piperidine scaffold. A variety of substituents could be tolerated on either the styrene or pyridine, including halogens and alkyl or alkynyl groups. The alkene could also be a β -substituted styrene containing a heteroatom in the alkyl chain.

Category

Metal-Catalyzed
Asymmetric
Synthesis and
Stereoselective
Reactions

Key words

copper catalysis
dearomatization
pyridines

Synfact
of the month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.