G. PUPO, F. IBBA, D. M. H. ASCOUGH, A. C. VICINI, P. RICCI, K. E. CHRISTENSEN, L. PFEIFER, J. R. MORPHY, J. M. BROWN, R. S. PATON, V. GOUVERNEUR* (UNIVERSITY OF OXFORD AND ELI LILLY AND CO. LTD., WINDLESHAM, UK)

Asymmetric Nucleophilic Fluorination under Hydrogen Bonding Phase-Transfer Catalysis

Science **2018**, *360*, 638–642.

Hydrogen-Bonding Phase-Transfer Catalyst Enabled Asymmetric Fluorination

Significance: Gouverneur and co-workers merge two fundamental strategies of modern catalyst design, hydrogen-bonding interactions and phase-transfer catalysis, to report a chiral bis-urea catalyzed asymmetric nucleophilic fluorination of episulfonium salts formed in situ. The fluorinated products are formed in good to excellent yields and moderate to good enantioselectivities by using CsF.

Comment: Hydrogen-bonding phase-transfer catalysis (HB PTC) has been previously reported in highly enantioselective nitro-Mannich reactions (B. Wang, Y. Liu, C. Sun, Z. Wei, J. Cao, D. Liang, Y. Lin, H. Duan *Org. Lett.* **2014**, *16*, 6432). Gouverneur and co-workers develop such catalysts to solve a synthetic challenge; namely, to enable an asymmetric fluorination reaction using a safe and readily accessible inorganic fluorine source. Henceforth, the expansion of HB PTC to other systems is expected.

SYNFACTS Contributors: Benjamin List, Jennifer L. Kennemur
Synfacts 2018, 14(08), 0861 Published online: 18.07.2018
DOI: 10.1055/s-0037-1609893; **Reg-No.:** B05418SF

2018 © THIEME STUTTGART • NEW YORK

Category

Organo- and Biocatalysis

Key words

phase-transfer catalysis

nucleophilic fluorination

hydrogen bonding

bis-urea catalysis

Synfact
of the month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.