Synthesis of a Bruton's Tyrosine Kinase Inhibitor

Significance: The target molecule **N** is a Bruton's tyrosine kinase (BTK) inhibitor that is of interest for the treatment of rheumatoid arthritis. A major challenge in the synthesis depicted is the construction of the two axes of chirality marked α and β . Rotation about these chiral axes gives four possible atropisomeric diastereoisomers. While the higher-energy C–C chirality axis α is relatively stable (rotational barrier 28 kcal/mol), the lower-energy C–N bond β (rotational barrier 26 kcal/mol) has a rotational half-life of hours to days, with a risk of epimerization.

Comment: The chiral axis α was constructed using an asymmetric Suzuki–Miyaura reaction at 5 °C to afford biaryl **K** (*dr* = 16:1). After crystallization from *n*-BuOH, **K** was obtained with *dr* = 65:1. The chiral axis β was created by crystallization-induced diastereoselection at 25 °C. Compound **N** was obtained in 90% yield and *dr* > 99:1. Thus the higher barrier diastereoisomer **K** acted as a chiral template to construct the stereochemical configuration of lower barrier bond β . This route delivered >200 kg of the target compound **N** in 28.3% overall yield.

SYNFACTS Contributors: Philip Kocienski
 Synfacts 2018, 14(09), 0885 Published online: 20.08.2018
DOI: 10.1055/s-0037-1610562; **Reg-No.:** K05018SF