M. WANG, X. PU, Y. ZHAO, P. WANG, Z. LI, C. ZHU, Z. SHI* (NANJING UNIVERSITY, P. R. OF CHINA)

Enantioselective Copper-Catalyzed Defluoroalkylation Using Arylboronate-Activated Alkyl Grignard Reagents
J. Am. Chem. Soc. **2018**, *140*, 9061–9065.

Asymmetric Copper-Catalyzed Defluoroalkylation

Category

Metal-Catalyzed
Asymmetric
Synthesis and
Stereoselective
Reactions

Key words

defluorination

gem-difluoro-
alkenes

copper catalysis

Synfact
of the month

Selected examples:

Proposed mechanism:

Significance: Defluorination of trifluoromethyl alkenes has become a popular strategy in the past few years to introduce a new stereocenter and synthesize *gem*-difluoroalkenes. *gem*-Difluoroalkenes are useful handles for further functionalizations and they are also useful moieties in medicinal chemistry. Shi and co-workers report the enantioselective addition of alkyl Grignard reagents to trifluoromethyl alkenes with the assistance of an arylboronate.

SYNFACTS Contributors: Mark Lautens, Andrew Whyte
 Synfacts 2018, 14(09), 0929 Published online: 20.08.2018
 DOI: 10.1055/s-0037-1610578; Reg-No.: L09118SF

Comment: The reaction uses alkyl magnesium bromide reagents which initially react with the organoboron species to form a tetraorganoborate. The tetraalkylboron species reacts with the copper catalyst to undergo an enantioselective addition to the trifluoromethyl alkene followed by β -fluoride elimination. The reaction was successful with a variety of substituents on the trifluoromethyl alkenes including aryl, alkenyl, alkynyl, and alkyl groups. Unfunctionalized Grignard reagents were most extensively investigated in the reaction.