H. Y. BAE, D. HÖFLER, P. S. J. KAIB, P. KASAPLAR, C. K. DE, A. DÖHRING, S. LEE, K. KAUPMEES, I. LEITO, B. LIST* (MAX-PLANCK-INSTITUT FÜR KOHLENFORSCHUNG, MÜLHEIM AN DER RUHR, GERMANY AND UNIVERSITY OF TARTU, ESTONIA)
 Approaching Sub-ppm-Level Asymmetric Organocatalysis of a Highly Challenging and Scalable Carbon–Carbon Bond Forming Reaction
Nat. Chem. **2018**, *10*, 888–894.

Activating Ketones with Low Catalyst Loadings of a Potent Chiral Lewis Acid

Significance: List and co-workers report the use of chiral organic Lewis acids at ppm- and sub-ppm levels as catalysts for the challenging asymmetric Mukaiyama aldol reaction of ketones with silyl ketene acetals. The corresponding products were obtained in excellent yields and enantioselectivities, even on a decagram scale. The extremely low catalyst loading is unprecedented in catalytic asymmetric C–C bond-forming reactions.

Comment: Aldehydes are commonly used as electrophiles in the catalytic asymmetric Mukaiyama aldol reaction, whereas ketones have remained a challenge as substrates for this transformation. The presented work shows the excellent performance of silylated chiral imidodiphosphorimidates (*Angew. Chem. Int. Ed.* **2016**, *55*, 13200) as powerful Lewis acid catalysts in the aforementioned reaction of both aryl- and alkyl-substituted ketones.

SYNFACTS Contributors: Benjamin List, David Díaz-Oviedo
Synfacts 2018, 14(10), 1089 Published online: 17.09.2018
DOI: 10.1055/s-0037-1610913; **Reg-No.:** B07418SF