K. LIAO, Y.-F. YANG, Y. LI, J. N. SANDERS, K. N. HOUK, D. G. MUSAIEV, H. M. L. DAVIES*
(EMORY UNIVERSITY, ATLANTA AND UNIVERSITY OF CALIFORNIA, LOS ANGELES, USA;
ZHEIJANG UNIVERSITY OF TECHNOLOGY, HANGZHOU, P. R. OF CHINA)
Design of Catalysts for Site-Selective and Enantioselective Functionalization of Non-activated Primary C–H Bonds
Nat. Chem. **2018**, *10*, 1048–1055.

Enantioselective Functionalization of Primary C–H Bonds

Category

**Metal-Catalyzed
Asymmetric
Synthesis and
Stereoselective
Reactions**

Key words

rhodium catalysis

C–H activation

carbenes

Synfact
of the month

Significance: The authors previously described Rh-catalyzed functionalizations of secondary or tertiary C–H bonds (Liao et al. *Nature* **2016**, 533, 230; Liao et al. *Nature* **2017**, 551, 609). By tuning the catalyst, non-activated primary C–H bonds are functionalized with excellent site and enantioselectivities.

SYNFACTS Contributors: Mark Lautens, José F. Rodríguez
Synfacts 2018, 14(11), 1143 Published online: 18.10.2018
DOI: 10.1055/s-0037-1611078; **Reg-No.:** L13218SF

Comment: Subjecting chiral substrates to the reaction conditions allows access to either set of diastereomeric products when the appropriate catalyst enantiomer is employed. Stigmasteryl acetate could be selectively functionalized at its most sterically accessible primary C–H bond.