Bay-Region-Selective Annulative π -Extension (APEX) of Perylene Diimides with Arynes

Takayuki Nakamuro^{a,1} Kazushi Kumazawa^aHideto Ito^{*a,b} Kenichiro Itami^{*a,b,c}

^a Graduate School of Science, Nagoya University, Chikusa, Nagoya 464-8602, Japan
ito.hideto@g.mbox.nagoya-u.ac.jp

^b JST-ERATO, Itami Molecular Nanocarbon Project, Nagoya University, Chikusa, Nagoya 464-8602, Japan

^c Institute of Transformative Bio-Molecules (WPI-ITbM), Nagoya University, Chikusa, Nagoya 464-8602, Japan
itami@chem.nagoya-u.ac.jp

Published as part of the 30 Years SYNNETT – Pearl Anniversary Issue

Bay-region-selective annulative π -extension (APEX)

- One-step π -extension at unfunctionalized bay-region
- No halogenation/oxidation
- Two-directional APEX
- 5 examples of π -extended PDI and CDI with moderate yields

Received: 30.11.2018

Accepted after revision: 10.01.2019

Published online: 07.02.2019

DOI: 10.1055/s-0037-1611668; Art ID: st-2018-b0780-I

License terms:

Abstract A bay-region-selective annulative π -extension (APEX) reaction of perylene diimides (PDIs) has been achieved by means of in-situ generated reactive aryne intermediates. This method provides an efficient one-pot π -extension at the short axis of PDIs in a sequential manner. Mechanistically, an inverse-electron-demand Diels–Alder reaction might be operative for the transformation.

Key words annulative π -extension, perylene diimides, arynes, Diels–Alder reaction, coronene diimides, polycyclic aromatic hydrocarbons

Perylene diimides (PDIs)² are useful chemical scaffolds because of their rigid perylene π -conjugated systems and their electron-withdrawing imide groups. These unique π -systems bestow attractive optoelectronic properties³ that have contributed to versatile applications, for example, as fluorescent materials,^{3a} as laser compositions,^{3b} and even in supramolecular chemistry.^{3c} Therefore, a variety of PDI derivatives, together with methods for their functionalization, have been developed in attempts to improve and/or modify their optoelectronic and self-assembling properties.⁴ The π -character of PDIs can be modulated by π -extensions of PDI in two ways: along the longitudinal (long-axis)⁵ or along the transverse (short-axis)⁶ molecular axis. Although longitudinal modifications of PDIs have been well studied for the development of near-infrared-absorbing materials, the corresponding transverse extensions are less developed due to difficulties in synthesis. Müllen and co-workers demonstrated a π -extension of PDIs in the bay-regions (the transverse concave armchair edges) through several steps, in-

cluding bromination, Suzuki–Miyaura coupling, and cyclization [Scheme 1(a)].^{6a} As demonstrated by other researchers, lateral π -extension of PDIs relies on stepwise halogenation, coupling reactions, and cyclization or oxidation reactions, which reduces the synthetic efficiency and availability of the π -extended PDIs and loses the opportunity for further π -extension and functionalization in later stages of the process. Therefore, the development of direct and step-economical methods for obtaining π -extended PDIs is in high demand.

Recently, the annulative π -extension (APEX)⁷ reaction has attracted much interest in relation to the synthesis of polycyclic aromatic compounds (PACs), because the APEX reaction permits a one-step π -extension of nonfunctionalized polycyclic aromatic hydrocarbons (PAHs) or heteroaromatics without any prior functionalization, such as halogenation, thereby providing a variety of nanographenes⁸ or heteroatom-containing PACs⁹ that are difficult to access by conventional methods. For example, we have developed *K*-region (concave armchair edge)-selective APEX reactions of PAHs such as phenanthrenes,^{8a} pyrenes,^{8b} corannulene,^{8c} and chrysene,^{8d} catalyzed by cationic palladium complexes. In addition, APEX reactions of PAHs in the bay-region¹⁰ have also been demonstrated by Clar and Zander,^{10a,b} and by the groups of Scott,^{10c–f,j} Wu,^{10g} Matsuda and Stork,^{10h,i} Kubo,^{10k} Peña,^{10l} and Hoye.^{10m} These reactions involve Diels–Alder-type reactions with dienophiles such as alkynes, quinones, or arynes [Scheme 1(b)]. However, the range of available substrates is limited to perylene, benzoperylene, and bisanthene, exclusively, due to the harsh reaction conditions, and no examples of bay-region-selective APEX of PDIs have been reported despite the high demand for such processes.

(a) π -extension of perylene diimides (b) APEX reactions of perylene with arynes (Stork/Matsuda, Scott, Kubo, Peña, Hoye)¹⁰ **Scheme 1** (a) The classical stepwise π -extension method and the bay-region-selective annulative π -extension (APEX) method (this work) for π -extended perylene diimides. (b) Previously developed APEX reaction of perylene with arynes.**Table 1** Screening of the reaction conditions for bay-region-selective APEX reactions of perylene diimide **1** with benzyne precursors **2**^a

Entry	Deviations from the standard conditions for the reaction of 1a with 2a	Yield ^b (%) of 3aa	Yield ^b (%) of 4aa
1	–	63 (37) ^c	13
2	120 °C	55	10
3	24 h	58	9
4	2a (5.0 equiv)	38	25
5	2b instead of 2a	24	n.d. ^d
6	MeCN instead of PhCN	trace	n.d.
7	DMF instead of PhCN	n.d.	n.d.
8	toluene instead of PhCN	n.d.	n.d.
9	CsF instead of KF	35	trace
10	TBAT instead of KF	13	trace
11	1b instead of 1a	68 (46) ^e	9 ^f

^a Standard reaction conditions: **1a** (0.10 mmol, 1.0 equiv), **2a** (2.0 equiv), KF (5.0 equiv), PhCN (2.0 mL), 160 °C, 48 h.^b ¹H NMR yield determined by using CH₂Br₂ as an internal standard.^c Isolated yield.^d n.d. = not detected.^e ¹H NMR yield of **3ba**.^f ¹H NMR yield of **4ba**.

Here, we report the first bay-region-selective APEX reactions of PDIs with arynes as π -extending agents for the synthesis of π -extended PDIs [Scheme 1(a); bottom].

Optimization of the reaction conditions was performed by using the dimesityl-substituted PDI **1a** as a standard substrate (Table 1). PDI **1a** was treated with the benzyne¹¹ precursor 2-(trimethylsilyl)phenyl triflate (**2a**) (2.0 equiv) and KF (5.0 equiv) in benzonitrile as the solvent at 160 °C. After 48 hours, **1a** was almost consumed (>90% conversion) to afford the single-APEX product **3aa** and the double-APEX product **4aa** in 63% and 13% NMR yields, respectively (Table 1, entry 1). However, the separation of product **3aa** from **4aa** was difficult, and **3aa** was obtained in only 37% isolated yield. Reactions at a lower temperature or for a shorter reaction time resulted in slightly lower yields of the product (entries 2 and 3). The amount of **2a** affected the ratio of **3aa** and **4aa**; however, an exclusive synthesis of the doubly- π -extended PDI **4aa** was not achieved by changing the reaction conditions (entry 4). The use of 2-diazonobenzoate (**2b**) instead of **2a** gave an inferior result, probably due to the thermal lability of **2b** (entry 5; 24% NMR yield of **3aa**). An examination of various solvents revealed that benzonitrile is the optimal solvent [for details, see the Supporting Information (SI)]. In reactions using CsF or tetrabutylammonium difluorotriphenylsilicate (TBAT) instead of KF, **3aa** was exclusively obtained, but in a lower yields of 35% and 15%, respectively (entries 9 and 10); none of the double-APEX product **4aa** was generated. The reaction of PDI **1b**, which had 6-undecyl groups on the nitrogen atoms, was also examined, and comparable results to those for **1a** were obtained (entry 11).

Next, we explored the substrate scope of arynes in the bay-region-selective APEX reactions of PDIs **1a** and **1b** (Scheme 2). When the commercially available 2,3-naphthalene precursor **2c** was subjected to the APEX reaction with **1a** under the optimized reaction conditions shown in entry 1 of Table 1, only traces of the desired APEX product **3ac** were observed in the crude mixture. This situation was not markedly improved by changing various reaction conditions, such as the temperature, the amounts of reagents, or the solvent. This might be due to the instability of the 2,3-naphthalene intermediate under severe thermal reaction conditions.¹² However, the reaction of the highly soluble 6-undecyl-substituted PDI **1b** proceeded smoothly in MeCN at 120 °C to afford the APEX product **3bc** in 49% NMR yield and 36% isolated yield. On the other hand, the 9,10-phenanthryne precursor **2d** reacted with PDIs **1a** and **1b** with good efficiency to give the corresponding single-APEX products **3ad** and **3bd**, respectively. In these reactions, although traces of the corresponding double-APEX products were also detected in the crude mixture, products **3bc**, **3ad**, and **3bd** were easily isolated by simple chromatography techniques. When other arynes, such as electron-deficient 2,3-pyri-

dyne¹³ or alkyl- or aryl-substituted alkynes were subjected to the above reaction conditions, the starting PDIs were recovered without the formation of any APEX products. These results suggest that highly reactive triple bond, such as those of aromatic-hydrocarbon-based arynes are necessary for bay-region-selective APEX reaction of PDIs (see SI for details).

Scheme 2 Substrate scope of arynes in APEX reactions of PDIs **1a** and **1b**. Standard reaction conditions: **1a** (0.050 mmol, 1.0 equiv), **2a** (2.0 equiv), KF (5.0 equiv), PhCN (1.0 mL), 160 °C, 48 h. ¹H NMR yields were determined by using CH₂Br₂ as an internal standard. Isolated yields are shown in parentheses.

To clarify the reaction profiles and to explore the potential of our bay-region APEX reaction, we performed sequential APEX reactions (Scheme 3). First, the single-APEX product **3aa** was further subjected to the optimized APEX reaction conditions with **2a** in an attempt to obtain the double-APEX product **4aa**. However, a quite-low conversion of **3aa** and the formation of a small amount of **4aa** were observed [Scheme 3(a)], which is inconsistent with the result obtained in entry 1 of Table 1. This result implies that the actual intermediate for the second bay-region APEX reaction in entry 1 of Table 1 might not be **3aa**, but instead **3aa-H₂**, the primary product of the Diels–Alder reaction before the ejection of two hydrogen atoms that completes the APEX reaction through rearomatization. On the contrary, the APEX reaction of **3ad** with **2a** afforded the π -extended PDI **5**, along with a good recovery of **3ad** [Scheme 3(b)]. The resulting nonsymmetrical nanographene diimide structure might provide a new entry to π -extended coronene diimides (CDIs) for future photophysical and electrochemical applications.¹⁴

Scheme 3 APEX reactions of **3aa** and **3ad** for the synthesis of further π -extended PDIs **4aa** and **5**

Stimulated by these difference in reactivities between PDIs in the bay-region-selective APEX reaction, we performed DFT calculations at the B3LYP/6-31G(d) level of theory. Scott and co-workers reported that the bay-region-selective APEX reaction of unsubstituted PAHs such as phenanthrene, perylene and bisanthene with acetylene occurs through a Diels–Alder reaction with subsequent aromatization.^{10e} Furthermore, they showed experimentally and computationally that perylene is the smallest PAH that affords the APEX product, with an activation energy of 30.0 kcal/mol for the Diels–Alder reaction step. In our calcula-

tions using PDIs **1a**, **3aa**, and **3ad** as dienes with benzyne as a dienophile, inverse-electron-demand Diels–Alder reactions¹⁵ successfully afford the dihydro-APEX products **3aa-H₂**, **4aa-H₂**, and **5-H₂** through transition states **TS1–TS3**, respectively (Scheme 4). The activation barrier to **TS1** was calculated to be 14.9 kcal/mol, which is relatively low and supports the smooth reaction progress observed in Table 1.¹⁶ On the contrary, the activation energies from **3aa** and **3ad** to **TS2** and **TS3** were calculated to be 17.1 and 15.7 kcal/mol, respectively, probably due to the increased LUMO energy of **3aa** (−3.24 eV) and **3ad** (−3.16 eV) compared with

Scheme 4 DFT calculation of the activation energies in Diels–Alder reaction of **1a**, **3aa**, and **3ad** with benzyne

that of **1a** (−3.42 eV). Although the reason for the lower activation energies in **TS3** than **TS2** is unclear, the present calculations well reflect the experimental results that Diels–Alder reactions occur not in **3aa**, but also in **1a** and **3ad**. Further tuning of arynes (HOMO) and PDIs (LUMO) should lead to even-more-efficient Diels–Alder APEX reactions at nonfunctionalized bay-regions of PDIs.

In summary, we have demonstrated the first bay-region-selective APEX reaction of PDIs by Diels–Alder reactions with arynes.¹⁷ This method gave the transversally π -extended PDIs in a single operation from unfunctionalized PDIs. Even nonsymmetrical π -extended PDIs, which are difficult to synthesize by conventional methods, were obtained with high efficiency. DFT calculations suggested that the bay-region APEX reaction probably proceeds through an inverse-electron-demand Diels–Alder reaction with a barrier of about 14.9–15.7 kcal/mol. We believe that our APEX method will provide an alternative synthetic tool for preparing various π -extended PDIs, thereby accelerating research in PDI-based materials science.

Funding Information

This work was supported by the ERATO program from JST (JPM-JER1302 to K.I.), JSPS KAKENHI Grants 18J01322 to T.N. and JP26810057, JP16H00907, JP17K19155, and JP18H02019 to H.I., the SUMITOMO Foundation (141495 to H.I.), and the DAIKO Foundation (H.I.).

Acknowledgment

The computations were performed at the Research Center for Computational Science, Okazaki, Japan. ITbM, which is supported by the World Premier International Research Center Initiative (WPI), Japan.

Supporting Information

Supporting information for this article is available online at <https://doi.org/10.1055/s-0037-1611668>.

References and Notes

- (1) Current address: Graduate School of Science, The University of Tokyo, Hongo, Tokyo 113-0033, Japan.
- (2) For a review on perylene diimides, see: Chen, L.; Li, C.; Müllen, K. *J. Mater. Chem. C* **2014**, *2*, 1938.
- (3) (a) Soh, N.; Ueda, T. *Talanta* **2011**, *85*, 1233. (b) Ramírez, M. G.; Morales-Vidal, M.; Navarro-Fuster, V.; Boj, P. G.; Quintana, J. A.; Villalvilla, J. M.; Retolaza, A.; Merino, S.; Díaz-García, M. A. *J. Mater. Chem. C* **2013**, *1*, 1182. (c) Würthner, F.; Saha-Möller, C. R.; Fimmel, B.; Ogi, S.; Leowanawat, P.; Schmidt, D. *Chem. Rev.* **2016**, *116*, 962. (d) Kojima, M.; Tamoto, A.; Aratani, N.; Yamada, H. *Chem. Commun.* **2017**, *53*, 5698. (e) Nagarajan, K.; Mallia, A. R.; Muralledharan, K.; Hariharan, M. *Chem. Sci.* **2017**, *8*, 1776.
- (4) (a) Battagliarin, G.; Li, C.; Enkelmann, V.; Müllen, K. *Org. Lett.* **2011**, *13*, 3012. (b) Sun, J.; Zhong, F.; Zhao, J. *Dalton Trans.* **2013**, *42*, 9595. (c) Ito, S.; Hiroto, S.; Shinokubo, H. *Chem. Lett.* **2014**, *43*, 1309.
- (5) (a) Pschirer, N. G.; Kohl, C.; Nolde, F.; Qu, J.; Müllen, K. *Angew. Chem. Int. Ed.* **2006**, *45*, 1401. (b) Zhao, X.; Xiong, Y.; Ma, J.; Yuan, Z. *J. Phys. Chem. A* **2016**, *120*, 7554.
- (6) (a) Avlasevich, Y.; Müller, S.; Erk, P.; Müllen, K. *Chem. Eur. J.* **2007**, *13*, 6555. (b) Li, Y.; Xu, L.; Liu, T.; Yu, Y.; Liu, H.; Li, Y.; Zhu, D. *Org. Lett.* **2011**, *13*, 5692. (c) Chaolumen Enno, H.; Murata, M.; Wakamiya, A.; Murata, Y. *Chem. Asian. J.* **2014**, *9*, 3136. (d) Calbo, J.; Doncel-Giménez, A.; Aragón, J.; Ortí, E. *Theor. Chem. Acc.* **2018**, *137*, 27.
- (7) For reviews on APEX reaction of aromatics, see: (a) Ito, H.; Ozaki, K.; Itami, K. *Angew. Chem. Int. Ed.* **2017**, *56*, 11144. (b) Ito, H.; Segawa, Y.; Murakami, K.; Itami, K. *J. Am. Chem. Soc.* **2019**, *141*, 3.
- (8) For our contributions towards APEX reactions of nonfunctionalized PAHs, see: (a) Ozaki, K.; Kawasumi, K.; Shibata, M.; Ito, H.; Itami, K. *Nat. Commun.* **2015**, *6*, 6251. (b) Yano, Y.; Ito, H.; Segawa, Y.; Itami, K. *Synlett* **2016**, *27*, 2081. (c) Kato, K.; Segawa, Y.; Itami, K. *Can. J. Chem.* **2017**, *95*, 329. (d) Ozaki, K.; Zhang, H.; Ito, H.; Lei, A.; Itami, K. *Chem. Sci.* **2013**, *4*, 3416. (e) Ozaki, K.; Murai, K.; Matsuoka, W.; Kawasumi, K.; Ito, H.; Itami, K. *Angew. Chem. Int. Ed.* **2017**, *56*, 1361. (f) Matsuoka, W.; Ito, H.; Itami, K. *Angew. Chem. Int. Ed.* **2017**, *56*, 12224. For a definition of the APEX reaction, see: (g) Segawa, Y.; Ito, H.; Itami, K. *Nat. Rev. Mater.* **2016**, *1*, 15002.
- (9) For selected APEX reactions of nonfunctionalized heteroaromatics, see: (a) Paria, S.; Reiser, O. *Adv. Synth. Catal.* **2014**, *356*, 557. (b) Ozaki, K.; Matsuoka, W.; Ito, H.; Itami, K. *Org. Lett.* **2017**, *19*, 1930. (c) Kitano, H.; Matsuoka, W.; Ito, H.; Itami, K. *Chem. Sci.* **2018**, *9*, 7556.
- (10) For selected bay-region selective APEX reactions, see: (a) Clar, E. *Ber. Dtsch. Chem. Ges.* **1932**, *65*, 846. (b) Clar, E.; Zander, M. *J. Chem. Soc.* **1957**, 4616. (c) Fort, E. H.; Donovan, P. M.; Scott, L. T. *J. Am. Chem. Soc.* **2009**, *131*, 16006. (d) Fort, E. H.; Scott, L. T. *J. Mater. Chem.* **2011**, *21*, 1373. (e) Fort, E. H.; Jeffreys, M. S.; Scott, L. T. *Chem. Commun.* **2012**, *48*, 8102. (f) Fort, E. H.; Scott, L. T. *Angew. Chem. Int. Ed.* **2010**, *49*, 6626. (g) Li, J.; Jiao, C.; Huang, K.-W.; Wu, J. *Chem. Eur. J.* **2011**, *17*, 14672. For the utilization of arynes, see: (h) Stork, G.; Leonia, N. J.; Matsuda, K. US 3364274, **1968**. (i) Stork, G.; Leonia, N. J.; Matsuda, K. US 3364275, **1968**. (j) Fort, E. H.; Scott, L. T. *Tetrahedron Lett.* **2011**, *52*, 2051. (k) Konishi, A.; Hirao, Y.; Matsumoto, K.; Kurata, H.; Kubo, T. *Chem. Lett.* **2013**, *42*, 592. (l) Schuler, B.; Collazos, S.; Gross, L.; Meyer, G.; Pérez, D.; Guitián, E.; Peña, D. *Angew. Chem. Int. Ed.* **2014**, *53*, 9004. (m) Xu, F.; Xiao, X.; Hoye, T. R. *Org. Lett.* **2016**, *18*, 5636.
- (11) For reviews on benzyne chemistry, see: (a) Takikawa, H.; Nishii, A.; Sakai, T.; Suzuki, K. *Chem. Soc. Rev.* **2018**, *47*, 8030. (b) Roy, T.; Biju, T. T. *Chem. Commun.* **2018**, *54*, 2580. (c) Shi, J.; Li, Y.; Li, Y. *Chem. Soc. Rev.* **2017**, *46*, 1707. (d) Bhojgude, S. S.; Bhunia, A.; Biju, A. T. *Acc. Chem. Res.* **2016**, *49*, 1658. (e) Pérez, D.; Peña, D.; Guitián, E. *Eur. J. Org. Chem.* **2013**, 5981.
- (12) Cioslowski, J.; Piskorz, P.; Moncrieff, D. J. *Am. Chem. Soc.* **1998**, *120*, 1695; and references cited therein.
- (13) (a) Saito, N.; Nakamura, K.; Shibano, S.; Ide, S.; Minami, M.; Sato, Y. *Org. Lett.* **2013**, *15*, 386. (b) For a review on heterocyclic arynes, see: Goetz, A. E.; Shah, T. K.; Garg, N. K. *Chem. Commun.* **2015**, *51*, 34.

- (14) (a) Sanyal, S.; Manna, A. K.; Pati, S. K. *J. Phys. Chem. C* **2013**, *117*, 825. (b) Zhang, C.; Shi, K.; Jiajun, X.; Lei, T.; Yan, Q.; Wang, J.-Y.; Pei, J.; Zhao, D. *Chem. Commun.* **2015**, *51*, 7144. (c) Yang, M.; Zhou, H.; Li, Y.; Zhang, Q.; Li, J.; Zhang, C.; Zhou, C.; Yu, C. *J. Mater. Chem. B* **2017**, *5*, 6572. (d) Paul, S. C.; Cammarata, V. J. *Electrochem. Soc.* **2018**, *165*, G116.
- (15) For a review on recent inverse-electron-demand Diels–Alder reactions, see: Jiang, X.; Wang, R. *Chem. Rev.* **2013**, *113*, 5515.
- (16) We also calculated each stationary point and transition state by other basis sets (See SI for details).
- (17) **3a; Typical Procedure**
A screw-capped glass tube containing a magnetic stirrer bar was charged sequentially with the dimesityl PDI **1a** (100 μ mol, 1.0 equiv, 62.4 mg), KF (0.51 mmol, 5.0 equiv, 29.5 mg), PhCN (2.0 mL), and 2-(trimethylsilyl)phenyl triflate (**2a**, 0.20 mmol, 2.0 equiv, 60.0 mg) under a stream of N_2 . The mixture was

stirred at 160 °C for 48 h, cooled to r.t., and passed through a short pad of silica gel (eluent: $CHCl_3$). The organic solvent was removed under reduced pressure to give a crude mixture that was analyzed by 1H NMR ($CDCl_3$) with CH_2Br_2 as an internal standard. The residue was purified by flash column chromatography (silica gel) to afford a mixture of **3aa** and **4aa**, which was further purified by gel-permeation chromatography to give **3aa** as a red solid; yield: 26.4 mg (37.7 μ mol, 37% isolated). 1H NMR (400 MHz, $CDCl_3$): δ = 10.2 (s, 2 H), 9.32–9.27 (m, 2 H), 9.24 (d, J = 8.4 Hz, 2 H), 9.10 (d, J = 8.4 Hz, 2 H), 8.19–8.13 (m, 2 H), 7.13 (s, 4 H), 2.42 (s, 6 H), 2.25 (s, 12 H). ^{13}C NMR (150 MHz, $CDCl_3$): δ = 163.5, 163.3, 138.8, 135.2, 134.2, 131.1, 130.0, 129.51, 129.47, 129.3, 129.2, 129.0, 128.6, 127.9, 125.1, 124.2, 123.3, 122.9, 122.5, 21.3, 17.9. HRMS (MALDI-TOF): m/z [$M + H$] $^+$ calcd for $C_{48}H_{33}N_2O_4$: 701.2435; found: 701.2434.