Stable Carbon-Centered Radicals Based on N-Heterocyclic Carbenes

Rajendra S. Ghadwal*

Anorganische Molekülchemie und Katalyse, Anorganische Chemie und Strukturchemie, Centrum für Molekulare Materialien, Fakultät für Chemie, Universität Bielefeld, Universitätsstraße 25, 33615 Bielefeld, Germany
rghadwal@uni-bielefeld.de

Received: 25.05.2019

Accepted after revision: 23.06.2019

Published online: 12.07.2019

DOI: 10.1055/s-0037-1611887; Art ID: st-2019-a0289-a

Abstract Carbon-centered radicals and diradicaloids based on classical N-heterocyclic carbene (NHC) scaffolds are readily accessible as crystalline solids. The presence of an aryl (Ar) substituent at the C2-position is the key to the remarkable stability of these open-shell species as it provides appropriate room for the spin-density delocalization. Two catalytic as well as high-yielding protocols have been developed to install a suitable aryl group at the C2-position of NHCs. The spin-density in mono-radicals (NHC^{Ar})[•] is mostly located on the parent carbene carbon (C2) atom. The bridging of two NHCs through a phenylene spacer (C₆H₄)_n enables the isolation of various *p*-quinodimethane (*p*-QDM) derivatives, which may be considered as open-shell Kekulé diradicaloids. The diradical character of these NHC-analogues of Thiele (*n* = 1), Chichibabin (*n* = 2), and Müller (*n* = 3) hydrocarbons [(NHC)(C₆H₄)_n(NHC)] can be tuned by a rational choice of the size and/or the topology of spacers. In this account, the synthesis, structure, and properties of this new class of radical hydrocarbons is presented.

- 1 Introduction
- 2 NHC-Monoradicals
- 3 NHC-Diradicaloids
- 4 Conclusion

Key words radical, diradical, diradicaloid, carbene, hydrocarbon, heterocycle, open-shell, paramagnetic

1 Introduction

Radicals are fundamental intermediates¹ and they play crucial roles in numerous biological² and industrially important processes.³ In general, radicals are short-lived species and, therefore, in a condensed phase, spontaneously react with surrounding (solvent, precursor, or by-product) molecules, undergo self-decomposition or dimerization to eventually form closed-shell products.⁴ Consequently, syn-

thetic access to thermally stable⁵ radicals for detailed structural investigations has been a subject of intense fundamental interest for augmenting our understanding about chemical bonding and electronic structure.⁶ In addition, stable radicals are also highly promising class of compounds for their potential applications in materials science and emerging areas.⁷ This is because of their unique optical, electronic, and magnetic properties.^{6b,8} The potential of stable radicals as unique building blocks for the design of next-generation advanced functional materials for applications in optoelectronics, spintronics, and solar cells is substantial.⁹ The incorporation of stable radicals into artificial molecular machines (AMMs) is very appealing because an electron's spin can be much easier and more precisely controlled, even with a smaller energy input, than a charge.^{7c} In this context, carbon-centered radicals have special advantages because of their planar trivalent nature, which promises structural diversity and elaborate designs. Stable radicals are also of immense interest as active materials in organic radical batteries (ORBs).¹⁰ These so-called 'eco batteries' have the potential to be cheaper, safer, and longer-lasting than current metal-based batteries and, thus, are of ever-growing importance in our modern technology-based society.¹¹ This becomes evidently more appealing if such radicals are derived from 1,3-imidazoli(ni)um precursors, which are promising electrolytes for battery applications. In recent years, radical catalysis has also emerged as an important tool in organic synthesis.¹²

Stable Radicals and Radicaloids

In 1900, Gomberg isolated the first stable radical, the triphenylmethyl **I** (Scheme 1).¹³ This seminal report marked the dawn of stable radical chemistry. **I**, however, undergoes self-association in the condensed phase, yielding the σ -dimer **II**. Although the delocalization of the unpaired electron

over the phenyl rings contributes to the stability of **I**, steric shielding of the central carbon atom that retains the most spin density is considered the key factor. The use of bulky substituents at *para*- or *meta*-position(s) of the phenyl groups slows down the dimerization to some extent, but no monomeric species could be isolated. Derivatives containing *ortho*-substituted or perchlorinated aryl groups were found to be remarkably stable. Because of the synthetic modularity, triphenylmethyl derivatives remain the most well-known class of stable carbon-centered radical hydrocarbons reported to date. Nonetheless, only heavily substituted or chlorinated species are isolable as monomers.¹⁴ The presence of two unpaired electrons in a molecule adds additional complexity to the electronic structure. Molecules containing two unpaired electrons in two nearly degenerate molecular orbitals are called diradicals, which may have parallel (triplet) or antiparallel (singlet) spins.⁸ Depending on the interaction between the unpaired electrons, singlet diradicals are further classified as open-shell (OS) and close-shell (CS) singlets. It should however be noted that the nature of the singlet wave function changes progressively from an OS to a CS configuration. Thus, the region in which the triplet becomes strongly destabilized and where the OS singlet becomes a CS singlet is not clearly defined. The term 'diradicaloids' is used for molecules with partial singlet diradical nature in their ground state.¹⁵

Scheme 1 The triphenylmethyl (trityl) radical **I** and its σ -dimer **II**

After the isolation of the trityl radical **I**, Thiele reported the phenylene bridged compound **III** ($n = 1$; Scheme 2) in 1904.¹⁶ In 1907, Tschitschibabin isolated a related derivative **III** ($n = 2$) featuring a biphenylene bridge.¹⁷ In general, they can be described either as an OS diradical (**IIIa**) or a CS quinoid (**IIIb**) structure. This is, however, not very straightforward. The electronic structure and properties of these so-called Thiele's (**TH**) and Chichibabin's (**CH**) hydrocarbons had been a subject of intense scientific discussion for more than 70 years, and this issue came to be known as 'the Biradical Paradox'.¹⁸ In 1986, Montgomery et al.¹⁹ determined the solid-state molecular structures of these highly reactive species by single-crystal X-ray diffraction. Currently, **CH** is generally described as a diradicaloid (a resonance hybrid of **IIIa** and **IIIb**) with a significant diradical character, whereas **TH** is considered as a *p*-quinodimethane (*p*-QDM). This can be explained based on the recovery of one (**TH**) and two (**CH**) Clar's aromatic sextet ring(s) in the OS diradical forms **IIIa**.²⁰ Obviously, the Müller's hydrocarbon (**MH**)²¹ with a terphenylene (**I**, $n = 3$) bridge reported in 1941 has higher diradical character compared with **TH** and **CH**. So, the **MH** is extremely reactive and it remains a challenging task to obtain pure samples for reliable measurements.

Stable diradicaloids delocalized over π -conjugated systems are appealing synthetic targets because of their unique physical properties and potential applications in nonlinear optics, molecular electronics, and organic spintronics.²² In this context, **TH**, **CH**, **MH**, and related molecules serve as very promising model systems. Over the past few years, different synthetic strategies have been developed to thermodynamically and/or kinetically stabilize various an-

Biographical Sketches

Rajendra S. Ghadwal studied chemistry at the University of Rajasthan Jaipur. In 2008, he joined the research group of Professor Herbert W. Roesky at the Georg-August-Universität (GAU) Göttingen as a postdoctoral research fellow. His postdoctoral work was focused on stable silylenes and silicon-heteroatom multiple bonded compounds. He started his independent research career at the GAU Göttingen in 2011 with a special emphasis on the development of unconventional carbon-donor ligands for organometallic catalysis and for the stabilization of open-shell main-group compounds. He re-

located his research group in 2015 from Göttingen to Bielefeld in Germany. In 2016, he completed his Habilitation in inorganic chemistry under the mentorship of Professor Dietmar Stalke at the GAU Göttingen and subsequently rehabilitated (Umhabilitation) at the Bielefeld University. He is currently a Privatdozent at the Bielefeld University and leading the 'Molecular Inorganic Chemistry and Catalysis' research group (www.ghadwalgroup.de). His research endeavors on the exploration of open-shell main-group compounds, π -conjugated molecular frameworks, base-metal (Ni and Cu in particular)

catalysis with mesoionic carbenes (MICs), optoelectronic properties of MIC-metal complexes, small molecules activation and functionalization aim to advance fundamental low-valent main-group element (including carbon) chemistry for sustainable synthesis and advanced functional materials. He recently received a Feodor Lynen Research Fellowship for Experienced Researchers from the Alexander von Humboldt Foundation to collaborate with Professor Christopher C. Cummins at the Massachusetts Institute of Technology (USA).

alogues of **TH**, **CH**, and **MH**. Bis(triarylamine) dications **IV** (Scheme 2) are isoelectronic to **III**, in which the $\text{Ph}_2\text{C}^\bullet$ moiety of **III** is replaced by an Ar_2N^{++} unit.²³ The benzannulation approach has also been successfully employed to isolate stable polycyclic hydrocarbons such as **V**.^{9a,24} In 2017, Wu et al. reported the oxindolyl-based carbon-centered diradicaloid **VI**, an analogue of the **MH**.²⁵ Synthetic strategies to various carbon-centered stable radicals have been summarized in some recent review articles.^{6b,26}

N-Heterocyclic carbenes (NHCs **VII**; Scheme 3) are very versatile σ -donor neutral ligands and have been extensively explored in transition-metal chemistry and catalysis.²⁷ Similarly, numerous compounds containing a low-valent main-group element,²⁸ including radicals (**VIII**),²⁹ have been stabilized by the use of NHCs. In this context, the success of Bertrand's cyclic alkyl amino carbenes (CAACs, **IX**) in stabilizing open-shell main-group or transition-metal species is remarkable.³⁰ This is because of the favorable electronic structure of CAACs. The smaller HOMO–LUMO energy gap of CAACs compared with classical NHCs makes them not only stronger σ -donors but also very efficient π -acceptors with respect to the latter.³¹ The enhanced electrophilicity of CAACs originates from the low-lying $\pi^*(\text{C}_{\text{carbene}}-\text{N})$ orbital that allows captodative³² interactions between the carbene (**IX**) and a main-group moiety (E) in the corresponding derivatives (**X**).^{30a-d,33} Various carbon-centered stable radicals have been also isolated using CAAC-frameworks.³⁴ In contrast, stable organic radicals derived from NHCs (**VII**) have so far remained limited to some specific compounds featuring a redox active nitrogen or sulfur atom(s).³⁵

Our research foci of developing new catalytic structures for sustainable synthesis³⁶ and accessing open-shell main-group compounds^{27b,37} for advanced functional materials relies on the development as well as exploration of uncon-

ventional carbon-donor ligands/scaffolds. For this purpose, we use classical NHCs as fundamental structural motifs to design new ligand sets.³⁸ This is because of the easy synthetic accessibility and tunable steric as well as electronic properties of NHCs. The available structural variables (Figure 1) in NHCs offer a plethora of opportunities to fine-tune the properties of derived molecules in many ways. These variables clearly appear to be more important and decisive in addressing the stability issue of highly sought-after open-shell molecules. Therefore, NHCs are an ideal choice for the design of stable radicals with adaptable properties. 1,3-Imidazolium salts **XI** ($\text{R}' = \text{H}$) are typical precursors to NHCs (**VII**) (Scheme 4). Their cyclic voltammograms (CVs) exhibit a very negative reduction potential (ca. -2.3 V vs. Fc/Fc^+) that is irreversible.³⁹ Not surprisingly, chemical reduction of **XI** ($\text{R}' = \text{H}$) with potassium in refluxing THF affords free NHCs (**VII**). The formation of **VII** is believed to result from the decomposition of unstable radical **XII-H**.³⁹ Muon resonance spectroscopy (μRS) studies showed that **VII** reacts with $\text{Mu}^\bullet$ (muonium, a hydrogen atom equivalent) to generate the radical **XII-Mu**, analogous to **XII-H**.⁴⁰ The **XII-Mu** species was calculated to feature the nonplanar radical center with 74.9% of the spin density located at the carbene-carbon atom.

These early studies clearly emphasize the requirement of a suitable substituent at the C2-position of **XI** to stabilize putative NHC-radicals. We previously showed that various transition-metal³⁶ as well as main-group element^{38,41} compounds featuring a mesoionic carbene (MIC), which may also be referred to as abnormal NHCs (aNHCs),⁴² **XII** can be readily prepared by the deprotonation of C2-arylated 1,3-imidazolium salts **XII** ($\text{R}' = \text{Ar}$) and in no case was cleavage of the $\text{C2}-\text{C}_{\text{Aryl}}$ bond observed (Scheme 4). This encouraged us to reason that stable NHC-based radicals **XIV** should be synthetically accessible on one-electron reduction of the appropriate C2-arylated 1,3-imidazoli(ni)um salts. This ac-

Scheme 4 Deprotonation of XI (R = aryl groups; R' = H or an aryl group) with a base to NHCs (VII) or MICs (XII). Reduction of XI (R' = H) to an NHC via a transient radical XII-H. Reaction of VII with Mu to XII-Mu. One-electron reduction of XI (R' = Ar) to stable radicals XIV

count summarizes recent advances, particularly from this laboratory, in NHC-based carbon-centered stable radicals chemistry.

2 NHC-Monoradicals

For the synthesis of desired C2-arylated precursors **3**, two protocols based on palladium^{36e} and nickel^{36b} catalysis have been developed in this laboratory. While the scope of palladium catalysis is limited to aryl iodides, nickel catalysis protocol is equally compatible with aryl bromides and iodides. Remarkably, bis-1,3-imidazoli(ni)um salts are also accessible by two-fold C–C coupling of appropriate aryl dihalides with various NHCs (see below). The yield of reactions in most cases is more than 80%. The product precipitates out in the employed solvent (*o*-xylene, THF, toluene or 1,4-dioxane) and can be readily extracted by simple filtration. No additional purification is required and the product can be used as such. Two typical examples with unsaturated (**1a**) and saturated (**2**) NHCs are shown in Scheme 5.

Scheme 5 Direct C2-arylation of NHCs **1a** and **2** to **3a**, **3b**, **4a**, and **4b**

In contrast to the C2-protonated 1,3-imidazolium salts **XI** (R' = H), the CVs of C2-arylated 1,3-imidazoli(ni)um salts **3a**, **3b** ($E_{1/2} = -1.66$ V, -1.91 V) and **4a**, **4b** ($E_{1/2} = -1.42$ V, -1.49 V vs. Fc⁺/Fc) exhibit one-electron reduction at a very low potential (Figure 2).⁴³ Note, the reduction potential for **3** and **4** is not only significantly lower compared with **XI-H** ($E_{1/2} \geq -2.3$ V) but the process is also reversible. This indicates the accessibility and stability of the corresponding radicals.

Figure 2 Cyclic voltammograms (CVs) of **3a**, **b** and **4a**, **b** (in CH₃CN/nBu₄NPF₆, 100 mV s⁻¹ vs. Fc⁺/Fc)

Indeed, the radicals (IPr^{Ph})[•] (**5a**), (IPr^{4-DMP})[•] (**5b**), (SIPr^{Ph})[•] (**6a**), and (SIPr^{4-Tol})[•] (**6b**) are accessible as crystalline solids on chemical reduction of the corresponding cations **3a**, **b** and **4a**, **b** with KC₈ (Scheme 6). Compounds **5** and **6** are deeply colored stable solids.⁴³

Scheme 6 Reduction of **3a**, **b** and **4a**, **b** to stable radicals **5a**, **b** and **6a**, **b**

Freshly prepared samples of **5** and **6** were NMR silent but each of them exhibited a doublet EPR signal (see below), indicating their paramagnetic character. Attempts to grow single crystals of **5a** and **5b** for X-ray diffraction failed because they slowly decompose in solution to give the oxidized products. The unsaturated NHC (IPr) and its 1,3-imidazolium salts (**XI**, **3**) are 6π-electron aromates. The lone pair at the carbene carbon atom of IPr is the s-type orbital and does not take part in the π-system. The IPr-derived radicals **5** are 7π-electron systems. Thus, the loss of the aromaticity in the IPr-derived radicals **5a**, **b** may account for their low stability. This is however not the case with the SIPr-derivatives. Fortunately, the radicals **6** based on a saturated NHC (SIPr) are stable, and suitable single crystals can be

Figure 3 Solid-state molecular structure of the monoradical **6b**; H atoms are omitted for clarity

grown from a saturated *n*-hexane solution. The solid-state molecular structures of **6a** and **6b** determined by X-ray diffraction are distinct from the free NHC (SIPr) and the corresponding 1,3-imidazolinium salts **4a,b**. A solid-state molecular structure of a typical radical **6b** is shown in Figure 3.

The formation of free NHCs **VII** from their 1,3-imidazolium precursor **XI-H** is a formal two-electron reduction (Scheme 4), whereas one-electron reduction of **XI-H** gives radicals **XII-H**. The value of C–N bond lengths and the N–C–N bond angle in the free NHC (**2**) is clearly larger compared with the corresponding protonated precursor (Table 1).

Table 1 Selected Bond Lengths (Å) and Bond Angles (°) for the Free NHC (SIPr), Radical Precursors (**4a** and **4b**), and Radicals (**6a** and **6b**)

Compound	C1–C4	C4–C5	C1–N1 ^b	N1–C1–N2
SIPr (2) ^a	–	–	1.346(2)	105.0(1)
(SIPr ^{Ph}) ⁺ (4a)	1.474(2)	1.395(2)	1.329(2)	111.8(2)
(SIPr ^{Ph}) [•] (6a)	1.398(5)	1.430(5)	1.410(4)	107.7(3)
(SIPr ^{4-Tol}) ⁺ (4b)	1.477(4)	1.390(4)	1.333(4)	111.8(3)
(SIPr ^{4-Tol}) [•] (6b)	1.406(2)	1.429(2)	1.405(1)	108.5(1)

^a Free NHC.

^b The value is averaged.

An intermediate bond length and bond angle alteration trend can be expected for the respective radicals. Indeed, the C1–N1/N2 bond lengths of **6a** (av. 1.405(4) Å) and **6b** (av. 1.404(2) Å) are considerably longer than those of **4a** (av. 1.330(4) Å) and **4b** (av. 1.329(4) Å) as well as of SIPr (**2**) (av. 1.346(2) Å).⁴⁴ The N1–C1–N2 bond angle of **6a** (108.9(3)°) and **6b** (108.5(9)°) is intermediate between the free carbene SIPr (105.0(1)°)⁴⁴ and the imidazolinium salts (**2-H**)⁺ (114.6(4)°), **4a** (111.8(1)°) or **4b** (111.0(1)°). Similarly, the C1–C4 bond length in **6a** (1.398(5) Å) and **6b** (1.406(2) Å) is shortened considerably in comparison with **4a** (1.474(2) Å) and **4b** (1.477(4) Å). Some bond lengths alteration across C_{ipso}–C_{ortho} bonds of the aryl (Ph or 4-Tol) substituent of **6a,b** is also observed in comparison with **4a,b**. The structural parameters are fully consistent with the radical character of **6a,b** and the partial delocalization of the unpaired electron over the aryl substituent at the carbene carbon atom.

The localized Mulliken atomic spin densities (see Figure 4 for **6b**) clearly reveal that the unpaired electron in **5** and **6** is mainly located on the carbene carbon atom (37–45%). In addition, there are considerable spin densities on the *ortho*- (19–23%) and *para*- (19–29%) carbon atoms of the aryl group, but only 8–11% on the ring nitrogen atoms. This suggests that the radicals **5** and **6** are stabilized by delocalization of the unpaired electron on the C2-aryl substituent. The EPR spectroscopic studies clearly support the paramagnetic nature of **5** and **6**. The room-temperature EPR spectra of **5** and **6** in THF show a doublet, which is a characteristic

Figure 4 Calculated spin densities (isovalue 0.003 a.u.) and Mulliken spin density (ρ_{spin}) numbers at the BP86+D3(BJ)/def2-TZVPP//BP86-D3(BJ)/def2-SVP level of theory for **6b**; H atoms are omitted for clarity

of monoradical species (Figure 5). The doublet displays couplings with two nitrogen atoms and several hydrogen atoms of the aryl substituent. The calculated hyperfine coupling (*hfc*) constants at the PBE0/EPR-II//BP86+D3(BJ)/def2-SVP level of theory for **6a** were found to be $A_{\text{iso}}(^{14}\text{N}) = 6.08$, 5.71; $A_{\text{iso}}(^1\text{H}) = -16.12$ (*ortho*-Ph), 7.27 (*meta*-Ph), -20.76 (*para*-Ph). The significant low $A_{\text{iso}}(^{14}\text{N})$ value (4.10–6.09) for **5a,b** and **6a,b** is in good agreement with the computed spin density at the nitrogen atoms.

Figure 5 Experimental and simulated EPR spectra of **6a** and **6b**

After having isolated the monoradicals **5** and **6**, one obvious quest was to probe the feasibility of stable species featuring more than one unpaired electron. The initial challenge was to access suitable starting materials.

3 NHC-Diradical(oid)s

Compounds featuring two unpaired electrons are called biradicals, diradicals, or diradicaloids, depending on the interaction between two spin dipoles. The presence of two unpaired electrons adds additional complexity to the structure and properties of these entities. The double carbenylation of 1,4-dihaloarenes with NHCs under nickel catalysis affords the desired dicationic species [(NHC)(C₆H₄)_n(NHC)](X)₂ (X = Br or I) (Scheme 7). Salts **7–9** can be isolated as white or pale-yellow crystalline solids in good to excellent (>80%) yields. Both unsaturated (**1a** and **1b**)⁴⁵ as well as saturated (**2**)⁴⁶ NHCs can be used.

The CVs of compounds [(IPr)₂(C₆H₄)₂](Br)₂ **7a** and [(IPr)₂(C₆H₄)₂](Br)₂ **7b** (Figure 6) exhibit two one-electron reductions in the –1.04 to –1.51 V region.⁴⁵ The first reversible one-electron reduction at $E_{1/2} = -1.04$ V (**7a**) and $E_{1/2} = -1.37$ V (**7b**) vs. Fc/Fc⁺ corresponds to a radical cation, which undergoes a second reduction at $E_{1/2} = -1.29$ V (**7a**) and $E_{1/2} = -1.51$ V (**7b**) to give the corresponding neutral species. A similar trend can also be seen for the derivatives **9a** and **9b** based on the saturated NHC, SiPr (**2**).⁴⁶ Nonetheless, both redox events in **9a** ($E_{1/2} = -0.80$ and –0.95 V) and **9b** ($E_{1/2} = -1.04$ and –1.28 V) occur at a lower potential compared with **7a** and **7b**. Similarly, CVs of compounds featuring a terphenylene spacer, **7c**, **8**, and **9c** also exhibit two cycles consistent with two one-electron (quasi)reversible redox processes.⁴⁷ Masuda and co-workers reported two-electron reduction of a closely related system [(SiPr)C≡C(C₆H₄)C≡C(SiPr)](Br)₂ and isolated a stable cumulene [(SiPr)=C=C(C₆H₄)=C=C(SiPr)].⁴⁸ Moreover, Bertrand et al. also reported^{30e,30f} stable cumulenes as well as mixed-valence compounds derived from CAACs that exhibit considerable open-shell character. Very recently, some mixed-valence systems based on carbene-pyrylium/pyridinium hybrids have been also reported.⁴⁹

Figure 6 Cyclic voltammograms of [(IPr)₂(C₆H₄)₂](Br)₂ **7a** (black line) and [(IPr)₂(C₆H₄)₂](Br)₂ **7b** (red line) in CH₃CN/Bu₄NPF₆ (vs. Fc/Fc⁺) in V

As indicated by the cyclic voltammetry studies, two-electron reduction of **7–9** with a suitable reagent should lead to the formation of the corresponding neutral compounds. Indeed, treatment of **7–9** with KC₈ afforded expected compounds **10–12** in good to excellent yields (Scheme

8). Ether open-shell (singlet or triplet) diradical (**A**) or closed-shell quinoid (**B**) resonance forms can be drawn to describe the structure of compounds **10–12**. The weight of each form depends on the interaction between the unpaired electrons, which, for a perfect quinoidal structure, results in a two-center-two-electron covalent bond formation at the expense of an aromatic sextet(s). Compounds **10–12** can be regarded as NHC-analogues of **TH**, **CH**, and **MH** (Scheme 2). Therefore, it is logical to compare and analyze the structure and properties of **10–12** with these classical HCs.

3.1 NHC Analogues of Thiele's Hydrocarbon (TH)

The **TH** (1904)¹⁶ and **CH** (1907)¹⁷ were already reported in the beginning of 20th century; it was, however, only in 1986 when Montgomery et al. successfully solved the solid-state molecular structures of these highly reactive species by single-crystal X-ray diffraction.¹⁹ This provided a direct comparison of the C–C bond length of the central phenylene ring and consequently settled the issues associated with ‘the Biradical Paradox’. It should however be noted that the purification of **TH**, **CH**, and related compounds is extremely difficult and the presence of even 0.01% monoradical impurity results in a doublet EPR signal. This may mask the weak intensity triplet expected for a triplet diradical, if it exists in a thermal equilibrium with an open-shell singlet diradical.⁵⁰ Structural parameters of **TH** and **CH** along with the NHC-derivatives **10a**, **10b** and **12a**, **12b**, are given in Table 2. The calculated bond lengths and bond angles for the respective closed-shell (CS) singlet and open-shell triplet (T) are also summarized in Table 2. As can be seen from these data, the bond length alteration (BLA) for the phenylene bridge in NHC-derivatives **10a** and **12a** is comparable to that of **TH** (0.10 Å). Therefore, the formation of **10a** and **12a** is accompanied by the loss of benzenoid structure of **7a** and **9a**. Experimental structures of **10a** and **12a** determined by X-ray diffraction show close resemblance with those calculated for closed-shell systems. In-

deed, a pure sample of **12a** is EPR silent and exhibits well-resolved ^1H and ^{13}C NMR signals for the NHC and (C_6H_4) units. As described by Nakano,⁵¹ the open-shell nature of a system can be expressed in terms of its diradical character ($y = 0$ for the close-shell and $y = 1$ for the pure singlet diradical). The diradical character for compound **10a** and **12a** was calculated to be zero ($y = 0$) and thus they are closed-shell species. The estimated value for the **TH** ($y = 0.31$) is, however, considerably higher.⁵⁰ While the ground state for both **10a** and **12a** is a closed-shell singlet, the calculated singlet-triplet energy gap (ca. 27 kcal/mol) is however rather small.

3.2 NHC Analogues of Chichibabin's Hydrocarbon (CH)

The BLA for compounds **10b** (Figure 7) and **12b** (0.08 Å), NHC-analogues of the **CH**, is smaller compared with **10a** and **12a** (0.10 Å), but larger with respect to that of **CH** (0.05 Å). The calculated diradical character for **10b** ($y = 0.41$

at B3LYP/def2-SVP level of theory) and **12b** ($y = 0.43$ at B3LYP/def2-TZVPP level of theory) is higher than that of **10a** and **10b** but it is considerably lower when compared with that of **CH** ($y = 0.72$). The singlet-triplet energy gap of 10.7 kcal calculated at B3LYP/def2-SVP for **10b** and **12b** is smaller than their monophenylene counterparts **10a** and **12a** (ca. 27 kcal/mol).

Figure 7 Molecular structure of **10b**; H atoms are omitted for clarity

Table 2 Selected Bond Lengths (Å) and Angles (°) Determined by X-ray Diffraction for the Precursors (**7a**, **7b**, **9a**, **9b**), NHC-Hydrocarbons (**10a**, **10b**, **12a**, **12b**), Thiele's (**TH**) and Chichibabin's (**CH**) HCs;¹⁹ and Calculated (B3LYP/def2-SVP) for **10a,b** and **12a,b** in Close-Shell (**CS**) and Triplet (**T**) Ground States.

Compound		$\text{C}_\alpha\text{-N}$	$\text{C}_\alpha\text{-C}_i$	$\text{C}_i\text{-C}_o$	$\text{C}_o\text{-C}_{o'}$	N-C-N	$\text{C}_i\text{-C}_i$	ΣN	BLA ^b
TH	$\text{Ph}_2\text{C}(\text{C}_6\text{H}_4)_2\text{CPh}_2$	–	1.381	1.449	1.346	–	–	–	0.10
7a	$[(\text{IPr})(\text{C}_6\text{H}_4)_2(\text{IPr})](\text{Br})_2$		1.474	1.391	1.390	107.8	–	360.0	–
10a	$[(\text{IPr})(\text{C}_6\text{H}_4)_2(\text{IPr})]$		1.375	1.453	1.354	104.6	–	350.7	0.10
CS	$[(\text{IPr})(\text{C}_6\text{H}_4)_2(\text{IPr})]^a$		1.393	1.458	1.364	103.9	–	358.4	0.09
T	$[(\text{IPr})(\text{C}_6\text{H}_4)_2(\text{IPr})]^a$		1.435	1.427	1.386	105.9/104.8	–	359.7	0.04
9a	$[(\text{SIPr})(\text{C}_6\text{H}_4)_2(\text{SIPr})](\text{I})_2$	1.326	1.478	1.397	1.390	113.2	–	356.4	–
12a	$[(\text{SIPr})(\text{C}_6\text{H}_4)_2(\text{SIPr})]$	1.404	1.371	1.451	1.349	108.2	–	349.4	0.10
CS	$[(\text{SIPr})(\text{C}_6\text{H}_4)_2(\text{SIPr})]^a$	1.413	1.390	1.461	1.362	105.8	–	355.1	0.10
T	$[(\text{SIPr})(\text{C}_6\text{H}_4)_2(\text{SIPr})]^a$	1.407	1.445	1.421	1.391	109.0		353.6	0.03
CH	$\text{Ph}_2\text{C}(\text{C}_6\text{H}_4)_2\text{CPh}_2$	–	1.415	1.424	1.372	–	1.448	–	0.05
7b	$[(\text{IPr})(\text{C}_6\text{H}_4)_2(\text{IPr})](\text{Br})_2$		1.464	1.394	1.388	107.1	1.492	359.2	–
10b	$[(\text{IPr})(\text{C}_6\text{H}_4)_2(\text{IPr})]$		1.388	1.441	1.359	104.6	1.413	354.0	0.08
CS	$[(\text{IPr})(\text{C}_6\text{H}_4)_2(\text{IPr})]^a$		1.404	1.449	1.369	104.2	1.416	359.0	0.08
T	$[(\text{IPr})(\text{C}_6\text{H}_4)_2(\text{IPr})]^a$		1.436	1.428	1.391	105.1	1.470	358.0	0.04
9b	$[(\text{SIPr})(\text{C}_6\text{H}_4)_2(\text{SIPr})](\text{Br})_2$	1.331	1.466	1.395	1.383	111.5	1.494	355.5	–
12b	$[(\text{SIPr})(\text{C}_6\text{H}_4)_2(\text{SIPr})]$	1.390	1.386	1.443	1.359	107.9	1.408	347.3	0.08
CS	$[(\text{SIPr})(\text{C}_6\text{H}_4)_2(\text{SIPr})]^a$	1.403	1.398	1.450	1.367	107.0	1.415	352.9	0.08
T	$[(\text{SIPr})(\text{C}_6\text{H}_4)_2(\text{SIPr})]^a$	1.409	1.424	1.433	1.387	107.6	1.481	351.4	0.04

^a Calculated for closed-shell (**CS**) and triplet (**T**) states.

^b Bond length alteration (BLA) = $(\text{C}_i\text{-C}_o\text{-C}_o\text{-C}_{o'})$.

This suggests that compounds **10b** and **12b** should be considered as singlet diradicaloids. Multireference CASSCF(2,2)+NEVPT2/def2-SVP calculations for model systems **10b-M** and **12b-M** (where Dipp groups of **10b** and **12b** are replaced by methyl groups) also support their diradicaloid character. The extent of pyramidalization of the nitrogen atom (Table 2) of the 1,3-imidazol(in)e rings can also account for the quinoid character of **10a,b** and **12a,b**. As indicated by the sum of the angles at the nitrogen atom (ΣN), the nitrogen atoms in the precursors **7a** ($\Sigma N = 360.0^\circ$) and **9a** ($\Sigma N = 359.2^\circ$) containing the unsaturated NHC (IPr) are planar. One of the nitrogen atoms in **10a** ($\Sigma N = 350.7^\circ$) and **12a** ($\Sigma N = 354.0^\circ$) is, however, pyramidalized. A rather large pyramidalization in the former suggests its higher quinoid character compared with the latter. A similar trend can also be seen in compounds **10b** and **12b** derived from the saturated NHC (SIPr).

3.3 NHC Analogues of Müller's Hydrocarbon (MH)

The MH²¹ with a terphenylene (**III**, $n = 3$, Scheme 3) bridge has higher diradical character compared to **TH** and **CH**. This can be rationalized according to Clar's aromatic sextet rule²⁰ as the OS diradical form **IIIa** preserves three aromatic rings. Consequently, the **MH** is extremely reactive and it remains a challenging task to obtain pure samples for reliable measurements. Nonetheless, stable analogues of **MH** have been reported by using appropriate synthetic approaches (see above). The exceptional stability of **10a,b** and **12a,b** prompted us to prepare stable NHC-analogues of **MH**. Fortunately, the nickel catalysis protocol with a slightly higher loading (10 mol%) gave access to the desired dications **7c**, **8**, and **9c** ($n = 3$; Scheme 7) using three different NHCs. Consistent with the CVs of **7c**, **8**, and **9c**, their two-electron reduction with KC_8 afforded compounds **10c**, **11**, and **12c** ($n = 3$; Scheme 8) as stable crystalline solids after workup. Structural parameters (Table 3) of **10c**, **11**, and **12c** (Figure 8) clearly reveal the enhanced weight of the open-shell diradical form compared with related CH-analogues (**10b** and **12b**) (Table 2). The BLA is in the 0.05–0.07 Å range for the central and terminal phenylene rings of the ter-

phenylene unit. This suggests that compounds **10c**, **11**, and **12c** still have a considerable contribution of the quinoid resonance form, which is nonetheless smaller compared to that of **10a,b** and **12a,b**.

The BLA calculated for the RB3LYP-optimized structures for singlet diradicaloids was found to be similar to those of the solid-state data of **10c**, **11**, and **12c** obtained by X-ray diffraction (Table 3). As expected, elongation of the spacer-size leads to enhancement of the diradical character of **10c** (0.65), **11** (0.64), and **12c** (0.65). The diradical character calculated for the **TH** (0.28), **CH** (0.59), and **MH** (0.82) at the same level of theory (PUHF/6-311G*) indicate that the replacement of the Ph_2C moiety with an NHC leads to the reduction of diradical character. This may be due to the difference in the delocalization of frontier molecular orbitals concerning the diradical character.

Table 3 Selected Bond Lengths (Å) and Angles ($^\circ$) Determined for **7c**, **9c**, **10c**, and **12c** by X-ray Diffraction

	C_1-C_2	C_2-C_3	C_4-C_5	C_6-C_7	C_8-C_9	BLA ^a
7c	1.392	1.390	1.482	1.396	1.390	0.00
10c	1.436	1.364	1.431	1.425	1.367	0.07 [0.06] ^c
9c	1.398	1.387	1.481	1.396	1.389	0.01
12c ^b	1.436 (1.435)	1.364 (1.369)	1.433 (1.434)	1.425 (1.429)	1.372 (1.370)	0.07 [0.05] ^c 0.07 [0.06] ^c

^a Bond length alteration (BLA) = $(C_1-C_2-C_3-C_4)$. The values for C_1-C_2 , C_2-C_3 , and C_4-C_5 are averaged.

^b **12c** contains two molecules in the asymmetric unit (values of the second molecule).

^c BLA for the middle phenylene ring [$C_4-C_5-C_6-C_7$].

The singlet-triplet energy gap for **10c** (7.24), **11** (7.60), and **12c** (7.40 kcal/mol) is significantly smaller compared to the related monophenylene **10a** (38.28) and biphenylene **12a** (18.56 kcal/mol) derivatives at the same level of theory. While this implies the singlet ground state for compounds **10c**, **11**, and **12c**, the triplet state may be populated to some extent at moderate temperature given a relatively small ΔE_{S-T} . Indeed, the EPR spectra of **10c**, **11**, and **12c** (Figure 9a) each exhibit a half-field signal at 298 K for the forbidden $\Delta M_s = 2$ transition, a characteristic of triplet diradicals. The calculated spin-density distribution maps for the singlet state of **10c**, **11**, and **12c** (Figure 9b) suggest that the unpaired electron density is mainly located at the carbene-carbon atoms and the terminal phenylene rings.

4 Conclusion and Outlook

Synthetic access to fundamentally unique stable radicals and diradicaloids based on classical NHCs has been ac-

Figure 8 Solid-state molecular structure of **12c**; H atoms are omitted for clarity

Figure 9 (a) Solid-state X-band EPR spectrum of **12c** at 298 K with inset showing the half-field signal. (b) Odd-electron density map (yellow mesh represents the isosurface with the contour value of 0.003 a.u.) of **12c**

complicated. The results reveal the impact of an aryl substituent at the C2-position of NHCs on the stability of derived radicals. Most strikingly, the desired starting materials can be readily prepared in almost quantitative yield by the direct C2-arylation of commercially available NHCs either under nickel or palladium catalysis. The methodology is straightforward and both the pre-catalysts $\text{Ni}(\text{COD})_2$ and $\text{Pd}_2(\text{dba})_3$ are commercially available. As the nature of aryl substituents has a direct influence on the stability and property of the derived radicals, further efforts to expand the scope of C2-arylation strategy would be valuable. Direct C2-arylation of air-stable 1,3-imidazoli(ni)um salts, instead of free NHCs, to the desired C2-arylated products is more appealing as it excludes the synthesis and handling of air-sensitive carbenes. This obviously relies on the development of new catalytic protocols, which is also important to broaden the scope of this transformation to different NHCs with variable steric and electronic features.

Systematic investigations are required to identify the key parameters that direct the physical (optical, electronic, magnetic) properties of NHC-based radicals and diradical(oid)s. Instead of simple aryl groups, the use of other π -conjugated substituents at the C2-position of NHCs is an appropriate means to further enhance the stability as well as to control the properties of derived radicals. In this con-

text, furan, thiophene, pyridine, pyrazine, cyclopentadiene, and other heterocycles containing a heavier main-group element (P, As, Se or Te) are interesting candidates.

Diradicaloids are considered as very promising singlet-fission (SF) materials for their application in organic solar cells.⁵² Molecular systems with too small $\Delta E_{\text{S-T}}$ gaps and those with triplet ground states are not suitable for efficient SF because they have large deviation from $2E(\text{T}1) = E(\text{S}1)$, leading to high energy loss and slow SF rate.^{52a,53} Because of the open-shell singlet ground state and a small singlet-triplet gap, NHC-based diradicaloids hold promise for the design of SF materials. A facile access to stable NHC-diradicaloids is expected to prompt further progress in synthesis, physicochemical measurement, and theoretical analysis on the open-shell compounds with variable diradical characters. Innovative NHC-radical(oid)s based on polycyclic aromatic hydrocarbon (PAH) spacers (e.g., naphthalene, anthracene, pyrene, and other acenes) are important targets we are working on.

NHC-diradicaloids are also interesting as candidates for open-shell (non)linear optical materials.^{22,54} Future targets include mixed-valence (MV) compounds, high-spin (tri- and polyradicals) molecules, and non-Kekulé diradicals based on NHC and other singlet carbene frameworks. The incorporation of NHC-based radicals in mechanically interlocked molecules (MIMs) has enormous potential in supramolecular chemistry and materials science and thus is worth pursuing. Among MIMs, the synthesis of catenanes and rotaxanes containing NHC-based pro-radical frameworks is under way in this laboratory.

Funding Information

Funding from the Deutsche Forschungsgemeinschaft (GH 129/4) is gratefully acknowledged.

Acknowledgment

I am very thankful to my current and former co-workers whose names appear in the corresponding citations for their valuable contributions. I am grateful to my collaborators Dr. Diego M. Andrada (Universität des Saarlandes), Dr. Maurice van Gestel (MPI für Kohlenforschung Mülheim), and Prof. Dr. Masayoshi Nakano (Osaka University, Japan) for their contributions. I express my gratitude to Professor Dr. Norbert Mitzel for his generous support and encouragement.

References

- (1) Renaud, P.; Sibi, M. P. *Radicals in Organic Synthesis*; Wiley-VCH: Weinheim, **2008**.
- (2) (a) Halliwell, B.; Gutteridge, J. M. C. *Free Radicals in Biology and Medicine*, 5th ed; Oxford University Press: UK, **2015**.
(b) Chatgililoglu, C.; Studer, A. *Encyclopedia of Radicals in Chemistry, Biology and Materials*; Wiley-VCH: Weinheim, **2012**.
(c) Berliner, L. J. *Spin Labeling: A Modern Perspective*, In *Stable*

- Radicals; John Wiley & Sons: Hoboken, **2010**, 521–534.
- (d) Goldstein, S.; Samuni, A. *Biologically Relevant Chemistry of Nitroxides*, In *Stable Radicals*; John Wiley & Sons: Hoboken, **2010**, 567–576.
- (3) (a) Bielawski, C. W.; Grubbs, R. H. *Prog. Polym. Sci.* **2007**, *32*, 1–29. (b) Tomlinson, E. P.; Hay, M. E.; Boudouris, B. W. *Macromolecules* **2014**, *47*, 6145.
- (4) Newcomb, M. *Radicals*, In *Reactive Intermediate Chemistry*; Moss, R. A.; Platz, M. S.; Jones, M. Jr., Ed.; John Wiley & Sons: Hoboken, **2005**, 121–163.
- (5) The term 'stable' refers to a molecular entity that persists for many days in solution and can be isolated (or stored) at room temperature (25 °C) under an inert gas (nitrogen or argon) atmosphere without appreciable decomposition.
- (6) (a) Hicks, R. G. *Can. J. Chem.* **2004**, *82*, 1119. (b) Hicks, R. G. *Stable Radicals: Fundamentals and Applied Aspects of Odd-Electron Compounds*; John Wiley & Sons Ltd: Weinheim, **2010**. (c) Sloan, G. J.; Vaughan, W. R. *J. Org. Chem.* **1957**, *22*, 750.
- (7) (a) Ai, X.; Chen, Y.; Feng, Y.; Li, F. *Angew. Chem. Int. Ed.* **2018**, *57*, 2869. (b) Wilcox, D. A.; Agarkar, V.; Mukherjee, S.; Boudouris, B. W. *Annu. Rev. Chem. Biomol. Eng.* **2018**, *9*, 83. (c) Wang, Y.; Frasconi, M.; Stoddart, J. F. *ACS Cent. Sci.* **2017**, *3*, 927.
- (8) (a) Salem, L.; Rowland, C. *Angew. Chem. Int. Ed. Engl.* **1972**, *11*, 92. (b) Borden, W. T. *Diradicals – A Fifty Year Fascination*, In *The Foundations of Physical Organic Chemistry: Fifty Years of the James Flack Norris Award, Vol. 1209*; American Chemical Society: Washington, DC, **2015**, p 251–303. (c) Abe, M. *Chem. Rev.* **2013**, *113*, 7011.
- (9) (a) Zeng, Z.; Shi, X.; Chi, C.; Lopez Navarrete, J. T.; Casado, J.; Wu, J. *Chem. Soc. Rev.* **2015**, *44*, 6578. (b) Ratera, I.; Veciana, J. *Chem. Soc. Rev.* **2012**, *41*, 303. (c) Hu, X.; Wang, W.; Wang, D.; Zheng, Y. *J. Mater. Chem. C* **2018**, *6*, 11232.
- (10) (a) Suga, T.; Nishide, H. *Rechargeable Batteries Using Robust but Redox Active Organic Radicals*, In *Stable Radicals*; John Wiley & Sons, Ltd: Weinheim, **2010**, 507–518. (b) Luo, J.; Hu, B.; Debruler, C.; Liu, T. L. *Angew. Chem. Int. Ed.* **2018**, *57*, 231. (c) Kim, D. J.; Hermann, K. R.; Prokofjevs, A.; Otley, M. T.; Pezzato, C.; Owczarek, M.; Stoddart, J. F. *J. Am. Chem. Soc.* **2017**, *139*, 6635.
- (11) Muench, S.; Wild, A.; Friebe, C.; Häupler, B.; Janoschka, T.; Schubert, U. S. *Chem. Rev.* **2016**, *116*, 9438.
- (12) (a) Beeson, T. D.; Mastracchio, A.; Hong, J.-B.; Ashton, K.; MacMillan, D. W. C. *Science* **2007**, *316*, 582. (b) Studer, A.; Curran, D. P. *Angew. Chem. Int. Ed.* **2016**, *55*, 58.
- (13) Gomberg, M. *J. Am. Chem. Soc.* **1900**, *22*, 757.
- (14) (a) Alcon, I.; Bromley, S. T. *RSC Adv.* **2015**, *5*, 98593. (b) Tidwell, T. T. *Triarylmethyl and Related Radicals*, In *Stable Radicals*; John Wiley & Sons, Ltd: Weinheim, **2010**, 1–3. (c) Veciana, J.; Ratera, I. *Polychlorotriphenylmethyl Radicals: Towards Multifunctional Molecular Materials*, In *Stable Radicals*; John Wiley & Sons, Ltd: Weinheim, **2010**, 33–76.
- (15) (a) Jung, Y.; Head-Gordon, M. *ChemPhysChem* **2003**, *4*, 522. (b) Seierstad, M.; Kinsinger, C. R.; Cramer, C. J. *Angew. Chem. Int. Ed.* **2002**, *41*, 3894. (c) Abe, M.; Ye, J.; Mishima, M. *Chem. Soc. Rev.* **2012**, *41*, 3808. (d) Grützmacher, H.; Breher, F. *Angew. Chem. Int. Ed.* **2002**, *41*, 4006. (e) Breher, F. *Coord. Chem. Rev.* **2007**, *251*, 1007.
- (16) Thiele, J.; Balhorn, H. *Ber. Dtsch. Chem. Ges.* **1904**, *37*, 1463.
- (17) Tschitschibabin, A. E. *Ber. Dtsch. Chem. Ges.* **1907**, *40*, 1810.
- (18) (a) Kanzaki, Y.; Shiomi, D.; Sato, K.; Takui, T. *J. Phys. Chem. B* **2012**, *116*, 1053. (b) McConnell, H. M. *J. Chem. Phys.* **1960**, *33*, 1868.
- (19) Montgomery, L. K.; Huffman, J. C.; Jurczak, E. A.; Grendze, M. P. *J. Am. Chem. Soc.* **1986**, *108*, 6004.
- (20) (a) Solà, M.; *Front. Chem.*; **2013**, *1*; DOI: 10.3389/fchem.2013.00022. (b) Clar, E. *The Aromatic Sextet*, In *Mobile Source Emissions Including Polycyclic Organic Species*; Rondia, D.; Cooke, M.; Haroz, R. K., Ed.; Springer Netherlands: Dordrecht, **1983**, 49–58.
- (21) Müller, E.; Pfanz, H. *Ber. Dtsch. Chem. Ges. B* **1941**, *74*, 1051.
- (22) (a) Kamada, K.; Ohta, K.; Kubo, T.; Shimizu, A.; Morita, Y.; Nakasuji, K.; Kishi, R.; Ohta, S.; Furukawa, S.-i.; Takahashi, H.; Nakano, M. *Angew. Chem. Int. Ed.* **2007**, *46*, 3544. (b) Sun, Z.; Ye, Q.; Chi, C.; Wu, J. *Chem. Soc. Rev.* **2012**, *41*, 7857. (c) Morita, Y.; Suzuki, S.; Sato, K.; Takui, T. *Nat. Chem.* **2011**, *3*, 197. (d) Nakano, M.; Champagne, B. *J. Phys. Chem. Lett.* **2015**, *6*, 3236.
- (23) Tan, G.; Wang, X. *Acc. Chem. Res.* **2017**, *50*, 1997.
- (24) (a) Li, G.; Phan, H.; Herng, T. S.; Gopalakrishna, T. Y.; Liu, C.; Zeng, W.; Ding, J.; Wu, J. *Angew. Chem. Int. Ed.* **2017**, *56*, 5012. (b) Liu, J.; Ma, J.; Zhang, K.; Ravat, P.; Machata, P.; Avdoshenko, S.; Hennersdorf, F.; Komber, H.; Pisula, W.; Weigand, J. J.; Popov, A. A.; Berger, R.; Müllen, K.; Feng, X. *J. Am. Chem. Soc.* **2017**, *139*, 7513.
- (25) Wang, J.; Xu, X.; Phan, H.; Herng, T. S.; Gopalakrishna, T. Y.; Li, G.; Ding, J.; Wu, J. *Angew. Chem. Int. Ed.* **2017**, *56*, 14154.
- (26) Kato, K.; Osuka, A. *Angew. Chem. Int. Ed.* **2019**, *58*, 8978.
- (27) (a) Hopkinson, M. N.; Richter, C.; Schedler, M.; Glorius, F. *Nature* **2014**, *510*, 485. (b) Rottschäfer, D.; Blomeyer, S.; Neumann, B.; Stämmler, H.-G.; Ghadwal, R. S. *Chem. Eur. J.* **2018**, *24*, 380.
- (28) (a) Ghadwal, R. S.; Azhakar, R.; Roesky, H. W. *Acc. Chem. Res.* **2013**, *46*, 444. (b) Wang, Y.; Robinson, G. H. *Inorg. Chem.* **2014**, *53*, 11815. (c) Wang, Y.; Robinson, G. H. *Dalton Trans.* **2012**, 337. (d) Yao, S.; Xiong, Y.; Driess, M. *Acc. Chem. Res.* **2017**, *50*, 2026.
- (29) (a) Walton, J. C.; Brahmi, M. M.; Monot, J.; Fensterbank, L.; Malacria, M.; Curran, D. P.; Lacote, E. *J. Am. Chem. Soc.* **2011**, *133*, 10312. (b) Ueng, S. H.; Solov'yev, A.; Yuan, X.; Geib, S. J.; Fensterbank, L.; Lacote, E.; Malacria, M.; Newcomb, M.; Walton, J. C.; Curran, D. P. *J. Am. Chem. Soc.* **2009**, *131*, 11256. (c) Walton, J. C.; McFadden, T. R.; Curran, D. P. *J. Am. Chem. Soc.* **2017**, *139*, 16514.
- (30) (a) Melaimi, M.; Jazsar, R.; Soleilhavoup, M.; Bertrand, G. *Angew. Chem. Int. Ed.* **2017**, *56*, 10046. (b) Soleilhavoup, M.; Bertrand, G. *Acc. Chem. Res.* **2015**, *48*, 256. (c) Martin, C. D.; Soleilhavoup, M.; Bertrand, G. *Chem. Sci.* **2013**, *4*, 3020. (d) Mondal, K. C.; Roy, S.; Roesky, H. W. *Chem. Soc. Rev.* **2016**, *45*, 1080. (e) Hansmann, M. M.; Melaimi, M.; Munz, D.; Bertrand, G. *J. Am. Chem. Soc.* **2018**, *140*, 2546. (f) Hansmann, M. M.; Melaimi, M.; Bertrand, G. *J. Am. Chem. Soc.* **2018**, *140*, 2206.
- (31) (a) Vummaleti, S. V. C.; Nelson, D. J.; Poater, A.; Gomez-Suarez, A.; Cordes, D. B.; Slawin, A. M. Z.; Nolan, S. P.; Cavallo, L. *Chem. Sci.* **2015**, *6*, 1895. (b) Mondal, K. C.; Roy, S.; Maity, B.; Koley, D.; Roesky, H. W. *Inorg. Chem.* **2016**, *55*, 163.
- (32) Viehe, H. G.; Janousek, Z.; Merenyi, R.; Stella, L. *Acc. Chem. Res.* **1985**, *18*, 148.
- (33) (a) Li, Y.; Chan, Y.-C.; Leong, B.-X.; Li, Y.; Richards, E.; Purushothaman, I.; De, S.; Parameswaran, P.; So, C.-W. *Angew. Chem. Int. Ed.* **2017**, *56*, 7573. (b) Gu, L.; Zheng, Y.; Haldón, E.; Goddard, R.; Bill, E.; Thiel, W.; Alcarazo, M. *Angew. Chem. Int. Ed.* **2017**, *56*, 8790. (c) Kundu, S.; Sinhababu, S.; Luebben, A. V.; Mondal, T.; Koley, D.; Dittrich, B.; Roesky, H. W. *J. Am. Chem. Soc.* **2018**, *140*, 151. (d) Kundu, S.; Samuel, P. P.; Sinhababu, S.; Luebben, A. V.; Dittrich, B.; Andrada, D. M.; Frenking, G.; Stückl, A. C.; Schwederski, B.; Paretzki, A.; Kaim, W.; Roesky, H. W.

- J. Am. Chem. Soc.* **2017**, *139*, 11028. (e) Braunschweig, H.; Krummenacher, I.; Légaré, M.-A.; Matler, A.; Radacki, K.; Ye, Q. *J. Am. Chem. Soc.* **2017**, *139*, 1802.
- (34) (a) Mahoney, J. K.; Martin, D.; Moore, C. E.; Rheingold, A. L.; Bertrand, G. *J. Am. Chem. Soc.* **2013**, *135*, 18766. (b) Hansmann, M. M.; Melaimi, M.; Bertrand, G. *J. Am. Chem. Soc.* **2017**, *139*, 15620. (c) Styra, S.; Melaimi, M.; Moore, C. E.; Rheingold, A. L.; Augenstein, T.; Breher, F.; Bertrand, G. *Chem. Eur. J.* **2015**, *21*, 8441. (d) Mahoney, J. K.; Martin, D.; Thomas, F.; Moore, C. E.; Rheingold, A. L.; Bertrand, G. *J. Am. Chem. Soc.* **2015**, *137*, 7519.
- (35) (a) Eymann, L. Y. M.; Tskhovrebov, A. G.; Sienkiewicz, A.; Bila, J. L.; Živković, I.; Rønnow, H. M.; Wodrich, M. D.; Vannay, L.; Corminboeuf, C.; Pattison, P.; Solari, E.; Scopelliti, R.; Severin, K. *J. Am. Chem. Soc.* **2016**, *138*, 15126. (b) Back, J.; Park, J.; Kim, Y.; Kang, H.; Kim, Y.; Park, M. J.; Kim, K.; Lee, E. *J. Am. Chem. Soc.* **2017**, *139*, 15300. (c) Wang, Y.; Hickox, H. P.; Xie, Y.; Wei, P.; Blair, S. A.; Johnson, M. K.; Schaefer, H. F.; Robinson, G. H. *J. Am. Chem. Soc.* **2017**, *139*, 6859. (d) Kim, Y.; Lee, E. *Chem. Eur. J.* **2018**, *24*, 19110.
- (36) (a) Ghadwal, R. S.; Lamm, J.-H.; Rottschäfer, D.; Schürmann, C. J.; Demeshko, S. *Dalton Trans.* **2017**, 7664. (b) Ho, N. K. T.; Neumann, B.; Stämmler, H.-G.; Menezes da Silva, V. H.; Watanabe, D. G.; Braga, A. A. C.; Ghadwal, R. S. *Dalton Trans.* **2017**, 12027. (c) Ho, N. K. T.; Reichmann, S. O.; Rottschäfer, D.; Herbst-Irmer, R.; Ghadwal, R. S. *Catalysts* **2017**, *7*, 262. (d) Rottschäfer, D.; Schürmann, C. J.; Lamm, J.-H.; Paesch, A. N.; Neumann, B.; Ghadwal, R. S. *Organometallics* **2016**, *35*, 3421. (e) Ghadwal, R. S.; Reichmann, S. O.; Herbst-Irmer, R. *Chem. Eur. J.* **2015**, *21*, 4247.
- (37) (a) Rottschäfer, D.; Ebeler, F.; Strothmann, T.; Neumann, B.; Stämmler, H.-G.; Mix, A.; Ghadwal, R. S. *Chem. Eur. J.* **2018**, *24*, 3716. (b) Rottschäfer, D.; Neumann, B.; Stämmler, H.-G.; Ghadwal, R. S. *Chem. Eur. J.* **2017**, *23*, 9044. (c) Rottschäfer, D.; Neumann, B.; Stämmler, H.-G.; Kishi, R.; Nakano, M.; Ghadwal, R. S. *Chem. Eur. J.* **2019**, *25*, 3244. (d) Sharma, M.; Blomeyer, S.; Neumann, B.; Stämmler, H.-G.; Ghadwal, R. S. *Chem. Eur. J.* **2019**, *25*, 8249. (e) Rottschäfer, D.; Sharma, M. K.; Neumann, B.; Stämmler, H.-G.; Andrada, D. M.; Ghadwal, R. S. *Chem. Eur. J.* **2019**, *25*, 8127.
- (38) Ghadwal, R. S. *Dalton Trans.* **2016**, 16081.
- (39) Gorodetsky, B.; Ramnial, T.; Branda, N. R.; Clyburne, J. A. C. *Chem. Commun.* **2004**, 1972.
- (40) McKenzie, I.; Brodovitch, J.-C.; Percival, P. W.; Ramnial, T.; Clyburne, J. A. C. *J. Am. Chem. Soc.* **2003**, *125*, 11565.
- (41) Ghadwal, R. S.; Rottschäfer, D.; Schürmann, C. J. *Z. Anorg. Allg. Chem.* **2016**, *642*, 1236.
- (42) Aldeco-Perez, E.; Rosenthal, A. J.; Donnadieu, B.; Parameswaran, P.; Frenking, G.; Bertrand, G. *Science* **2009**, *326*, 556.
- (43) Rottschäfer, D.; Neumann, B.; Stämmler, H.-G.; van Gastel, M.; Andrada, D. M.; Ghadwal, R. S. *Angew. Chem. Int. Ed.* **2018**, *57*, 4765.
- (44) Giffin, N. A.; Hendsbee, A. D.; Masuda, J. D. *Acta Crystallogr., Sect. E: Struct. Rep. Online* **2010**, *66*, o2194.
- (45) Rottschäfer, D.; Ho, N. K. T.; Neumann, B.; Stämmler, H.-G.; van Gastel, M.; Andrada, D. M.; Ghadwal, R. S. *Angew. Chem. Int. Ed.* **2018**, *57*, 5838.
- (46) Rottschäfer, D.; Neumann, B.; Stämmler, H.-G.; Andrada, D. M.; Ghadwal, R. S. *Chem. Sci.* **2018**, *9*, 4970.
- (47) Rottschäfer, D.; Busch, J.; Neumann, B.; Stämmler, H.-G.; van Gastel, M.; Kishi, R.; Nakano, M.; Ghadwal, R. S. *Chem. Eur. J.* **2018**, *24*, 16537.
- (48) Barry, B. M.; Soper, R. G.; Hurmalainen, J.; Mansikkamaki, A.; Robertson, K. N.; McClennan, W. L.; Veinot, A. J.; Roemmele, T. L.; Werner-Zwanziger, U.; Boere, R. T.; Tuononen, H. M.; Clyburne, J. A. C.; Masuda, J. D. *Angew. Chem. Int. Ed.* **2018**, *57*, 749.
- (49) (a) Antoni, P. W.; Bruckhoff, T.; Hansmann, M. M. *J. Am. Chem. Soc.* **2019**, *141*, 9701. (b) Antoni, P. W.; Hansmann, M. M. *J. Am. Chem. Soc.* **2018**, *140*, 14823.
- (50) Ravat, P.; Baumgarten, M. *Phys. Chem. Chem. Phys.* **2015**, *17*, 983.
- (51) Kamada, K.; Ohta, K.; Shimizu, A.; Kubo, T.; Kishi, R.; Takahashi, H.; Botek, E.; Champagne, B.; Nakano, M. *J. Phys. Chem. Lett.* **2010**, *1*, 937.
- (52) (a) Ito, S.; Nagami, T.; Nakano, M. *J. Photochem. Photobiol., C* **2018**, *34*, 85. (b) Nakano, M. *Chem. Rec.* **2017**, *17*, 27. (c) Minami, T.; Ito, S.; Nakano, M. *J. Phys. Chem. Lett.* **2013**, *4*, 2133.
- (53) (a) Messelberger, J.; Grünwald, A.; Pinter, P.; Hansmann, M. M.; Munz, D. *Chem. Sci.* **2018**, *9*, 6107. (b) Chen, M.; Bae, Y. J.; Mauck, C. M.; Mandal, A.; Young, R. M.; Wasielewski, M. R. *J. Am. Chem. Soc.* **2018**, *140*, 9184. (c) Casanova, D. *Chem. Rev.* **2018**, *118*, 7164.
- (54) Fukuda, K.; Suzuki, Y.; Matsui, H.; Nagami, T.; Kitagawa, Y.; Champagne, B.; Kamada, K.; Yamamoto, Y.; Nakano, M. *ChemPhysChem* **2017**, *18*, 142.