S. M. BANIK, K. PEDRAM, S. WISNOVSKY, N. M. RILEY, C. R. BERTOZZI* (STANFORD UNIVERSITY AND HOWARD HUGHES MEDICAL INSTITUTE, STANFORD, USA)

Lysosome Targeting Chimeras (LYTACs) for the Degradation of Secreted and Membrane Proteins

ChemRxiv 2019, DOI: 10.26434/chemrxiv.7927061.v1.

LYTACs for the Degradation of Secreted and Membrane Proteins

Significance: Lysosomal degradation makes use of cell-surface lysosomal targeting receptors (LTR), which facilitate the transport of proteins into the lysosome. The authors developed so-called LYTACs, which are chimeric molecules that can bind both an extracellular protein and an LTR, allowing for degradation of non-cytosolic proteins of interest in the lysosome. LYTACs are complementary to PROTACs (proteolysis targeting chimeras), which primarily target cytosolic proteins.

Comment: Mannose-6-phosphate receptor M6PR (or IGF2R) recognizes and binds mannose-6-phosphate residues on proteins and transports them to the lysosome for degradation. The LYTACs presented herein use glycopolypeptides of varying length, which can be attached to an antibody for the POI in a modular fashion using click chemistry. The versatile applicability is demonstrated by successfully targeting different extracellular proteins.

SYNFACTS Contributors: Dirk Trauner, Nynke A. Vepřek
Synfacts 2019, 15(06), 0671 Published online: 20.05.2019
DOI: 10.1055/s-0037-1612570; Reg-No.: T04119SF

2019 © Georg Thieme Verlag Stuttgart · New York

Category

Chemistry in
Medicine and
Biology

Key words

lysosome targeting
chimeras

LYTACs

protein degradation

lysosome

Synfact
of the
Month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.