Extended Endoscopic Approach for Resection of Craniopharyngiomas

João Paulo Almeida¹ Suganth Suppiah¹ Claire Karekezi¹ Miguel Marigil-Sanchez¹ Jay S. Wong²
Allan Vescan² Fred Gentili¹ Gelareh Zadeh¹

¹ Division of Neurosurgery, Toronto Western Hospital, University of Toronto, Toronto, Ontario, Canada

² Department of Otolaryngology/Head and Neck Surgery, Mount Sinai Hospital, University of Toronto, Toronto, Ontario, Canada

Address for correspondence João Paulo Almeida, MD, Division of Neurosurgery, Toronto Western Hospital, 4th Floor 399 Bathurst Street, Toronto, ON M5T 2S8, Canada
(e-mail: jpaulocavalcante@yahoo.com.br).

J Neurol Surg B 2018;79(suppl S2):S201–S202.

Abstract

Objectives Extended endoscopic approaches are useful for resection of selected craniopharyngiomas. Midline, extraventricular, and predominantly cystic lesions are good candidates for endoscopic resection. In this video, we demonstrate the endoscopic endonasal resection of a large suprasellar craniopharyngioma and discuss the nuances of the surgical technique.

Design/Setting Surgical video of an extended endoscopic approach for resection of a suprasellar craniopharyngioma.

Results We report the case of a 56-year-old woman who presented with bitemporal hemianopsia and visual acuity deterioration secondary to a large suprasellar solid–cystic lesion. The patient underwent an extended endoscopic transtuberculum approach for resection of the lesion, which was diagnosed as a papillary craniopharyngioma. This video discusses the anatomy and surgical technique applied for endoscopic resection of such lesions.

Conclusion Endoscopic endonasal surgery is a useful technique for management of craniopharyngiomas. It is associated with good clinical outcomes in selected cases. Complications, such as postoperative CSF leak, may occur and should be carefully managed. The link to the video can be found at: <https://youtu.be/EneOCiQE7yo>.

Keywords

- craniopharyngiomas
- tuberculum
- endoscopic
- extended
- endonasal

Conflict of Interest
None.

www.thieme.com/skullbasevideos

www.thieme.com/jnlsbvideos

received
October 12, 2017
accepted
December 12, 2017
published online
January 16, 2018

DOI <https://doi.org/10.1055/s-0037-1620244>.
ISSN 2193-6331.

© 2018 Georg Thieme Verlag KG
Stuttgart · New York

License terms

Fig. 1 Preoperative MRI scan demonstrating a suprasellar tumor, suggestive of craniopharyngioma. The coronal T1 contrast-enhanced scan (left) shows the lesion, located medial to the clinoid and supraclinoid internal carotid arteries and superior to the pituitary gland (as demonstrated by the arrows), compressing the optic chiasm. The sagittal imaging (right) demonstrates the relationship of the lesion with the floor of the III ventricle and basal cisterns (right). ICA, internal carotid artery; MRI, magnetic resonance imaging.

Fig. 2 Transtuberculum approach: surgical anatomy. The anatomical direction demonstrates the anatomical structures related to the transtuberculum approach. cICA, clinoid segment of the internal carotid artery; LOCR, lateral optic carotid recess; pclICA, paraclinoid segment of the internal carotid artery; OC, optic canal. Dashed line demonstrates the area of bone opening for the transtuberculum approach (copyright © Joao Paulo Almeida, MD).

Fig. 3 Exposure of the III ventricle and interpeduncular cistern after tumor resection. The basilar tip, PCA and SCA, III nerve, and PcomA are visible. PCA, posterior cerebral; PcomA, posterior communicating artery; SCA, superior cerebellar.