BCR-ABL Tyrosine Kinase Inhibitors: Which Mechanism(s) May Explain the Risk of Thrombosis?

Hélène Haguet^{1,2} Jonathan Douxfils^{1,4} Christian Chatelain¹ Carlos Graux³ François Mullier²
Jean-Michel Dogné¹

¹University of Namur, Namur Thrombosis and Hemostasis Center (NTHC), Namur Research Institute for Life Sciences (NARILIS), Department of Pharmacy, Namur, Belgium

²Université catholique de Louvain, CHU UCL Namur, Namur Thrombosis and Hemostasis Center, Hematology Laboratory, Yvoir, Belgium

³Université catholique de Louvain, CHU UCL Namur, Namur Thrombosis and Hemostasis Center, Department of Hematology, Yvoir, Belgium

⁴QUALIBlood s.a., Namur, Belgium

Address for correspondence Hélène Haguet, MBS, Department of Pharmacy, Namur Thrombosis and Hemostasis Center (NTHC), Namur Research Institute for Life Sciences (NARILIS), University of Namur, Rue de Bruxelles, 61, B-5000 Namur, Belgium (e-mail: helene.haguet@gmail.com).

TH Open 2018;2:e68–e88.

Abstract

Imatinib, the first-in-class BCR-ABL tyrosine kinase inhibitor (TKI), had been a revolution for the treatment of chronic myeloid leukemia (CML) and had greatly enhanced patient survival. Second- (dasatinib, nilotinib, and bosutinib) and third-generation (ponatinib) TKIs have been developed to be effective against BCR-ABL mutations making imatinib less effective. However, these treatments have been associated with arterial occlusive events. This review gathers clinical data and experiments about the pathophysiology of these arterial occlusive events with BCR-ABL TKIs. Imatinib is associated with very low rates of thrombosis, suggesting a potentially protecting cardiovascular effect of this treatment in patients with BCR-ABL CML. This protective effect might be mediated by decreased platelet secretion and activation, decreased leukocyte recruitment, and anti-inflammatory or antifibrotic effects. Clinical data have guided mechanistic studies toward alteration of platelet functions and atherosclerosis development, which might be secondary to metabolism impairment. Dasatinib, nilotinib, and ponatinib affect endothelial cells and might induce atherogenesis through increased vascular permeability. Nilotinib also impairs platelet functions and induces hyperglycemia and dyslipidemia that might contribute to atherosclerosis development. Description of the pathophysiology of arterial thrombotic events is necessary to implement risk minimization strategies.

Keywords

- BCR-ABL
- arterial thrombotic events
- tyrosine kinase inhibitors
- chronic myeloid leukemia

Introduction

In 2001, the approval of *imatinib*, the first-in-class tyrosine kinase inhibitor (TKI) targeting BCR-ABL, transformed the prognosis of patients with chronic-phase (CP) chronic myeloid leukemia (CML) from a life-threatening condition to a manageable and chronic disease.¹ Yet, despite satisfactory outcomes, 33% of patients did not achieved optimal response because of treatment resistance or intolerance.¹ The identi-

fication of the predominant resistance mechanism (i.e., point mutations in the kinase domain of Bcr-Abl) led to the development of second-generation BCR-ABL TKIs (dasatinib, nilotinib, and bosutinib, respectively, approved in 2006, 2007, and 2012) active against most of the BCR-ABL mutated forms.^{2,3} Second-generation TKIs demonstrated no or little improvement of the overall survival compared with imatinib,^{4–6} but two of these (i.e., dasatinib and nilotinib) improve

received
September 13, 2017
accepted after revision
November 27, 2017

DOI <https://doi.org/10.1055/s-0038-1624566>.
ISSN 2512-9465.

© 2018 Georg Thieme Verlag KG
Stuttgart · New York

License terms

surrogate outcomes and permit quicker and deeper achievement of molecular response, which is criteria to try treatment cessation (i.e., MR⁴ or higher molecular response stable for at least 2 years).⁷ Based on these results, dasatinib and nilotinib were approved in 2010 for frontline management of CML, whereas bosutinib is used only after failure or intolerance of first-line BCR-ABL TKIs. Unfortunately, these treatments were ineffective against a common mutation (14% of all mutations) in the gatekeeper residue of BCR-ABL (i.e., the T315I^a mutation),^{8–10} requiring the development of a third-generation TKI (ponatinib), efficient against this mutation. Ponatinib is currently the only treatment active against the T315I mutation and is therefore reserved for patients with this mutation or for patients resistant to frontline treatments.¹¹

Since its approval, the first-generation TKI, imatinib, has demonstrated reassuring safety profile, with low rate of grade 3/4 adverse events and excellent tolerability.^{12,13} Conversely, new-generation BCR-ABL TKIs—nilotinib, dasatinib, bosutinib, and ponatinib—are more recent and display different safety profile. Dasatinib, nilotinib, and ponatinib are largely associated with fluid retention and dasatinib specifically induces high rate of pleural effusions.^{14–18} Nilotinib induces metabolic disorders such as dyslipidemia and hyperglycemia, whereas bosutinib safety profile is mainly characterized by gastrointestinal events (i.e., diarrhea, nausea, vomiting).^{19,20} Finally, ponatinib has been rapidly associated with high rate of vascular occlusion.²¹

Recently, meta-analyses of randomized clinical trials established that ponatinib is not the only new-generation TKI that increases the cardiovascular risk.^{22,23} The four new-generation BCR-ABL TKIs increase the risk of vascular occlusive events compared with imatinib, especially arterial occlusive diseases, and this is in accordance with clinical trial data.^{22–25} However, this cardiovascular risk is controversy for dasatinib because of the low incidence (1.1 per 100 patient-year) of cardiovascular events in clinical trials.^{26,27} Recently, a large retrospective analysis of CP-CML patients treated with BCR-ABL TKIs at the MD Anderson Cancer Center confirmed the increased risk of vascular occlusive events with dasatinib.²⁸ Another controversial point is the effect of imatinib on the cardiovascular system. Indeed, imatinib is associated with low risk of cardiovascular events and it was therefore hypothesized that imatinib prevents their occurrence.^{29,30} Clinical data indicate that most patients developing arterial occlusive events with new-generation BCR-ABL TKIs are high-risk patients, but cardiovascular events also occurred in young and healthy patients. Additional information on clinical safety of BCR-ABL TKIs is described in the Supplementary Material (►Table S1). We assume that the mechanism underlying arterial thrombosis with BCR-ABL TKIs might be multiple. The predominance of arterial events raised concerns about the impact of BCR-ABL TKIs on platelet

functions, atherosclerosis, and metabolism, and precluded prothrombotic states to be responsible of these events.³¹

This review particularly focuses on the contribution of glucose and lipid metabolism, atherosclerosis, and platelets in the occurrence of cardiovascular events with new-generation TKIs. The last section discusses relevant off-targets that might be implicated in the cardiovascular toxicity. The discovery of the mechanism(s) by which arterial occlusive events arose in CML patients would help in the management of patients treated with BCR-ABL TKIs and implement risk minimization measures. Discovery of the pathophysiology of these events in CML patients might also led to the development of predictive biomarkers or to the development of new therapies with no or reduced cardiovascular toxicity profile while keeping an unaltered efficacy.

Impact on Platelet Functions

BCR-ABL TKIs are associated with both bleeding and thrombotic complications. ►Table 1 describes experiments assessing the impact of BCR-ABL TKIs on platelet production and functions. Imatinib and dasatinib induce hemorrhagic events in patients with CML. Interestingly, dasatinib-associated hemorrhages occurred both in patients with and without thrombocytopenia.³² In vitro and in vivo investigations demonstrated that dasatinib affects both platelet functions (i.e., platelet aggregation, secretion, and activation) and platelet formation by impairment of megakaryocyte migration.^{33–36} Furthermore, dasatinib decreases thrombus formation in vitro, in vivo, and ex vivo,³⁴ and decreases the number of procoagulant platelets (i.e., phosphatidylserine-exposing platelets).³⁵ Several dasatinib off-targets are implicated in platelet signaling and functions including members of the SFKs (e.g., Src, Lyn, Fyn, Lck, and Yes) (►Fig. 1).^{37,38} However, SFKs are also inhibited by bosutinib without disturbance of platelet aggregation and adhesion. Dasatinib also inhibits Syk, BTK, and members of the ephrin family^b (e.g., EphA2), all known to be involved in platelet functions.

Experimental assessments of platelet functions with imatinib demonstrate less pronounced effects on platelets. Imatinib inhibits platelet aggregation only at high doses,³⁴ and does not interfere with platelet aggregation in vivo.³⁹ However, in vitro studies also indicate decreased platelet secretion and activation by imatinib.³⁴ The mechanism by which imatinib inhibits platelet functions is unknown. Oppositely to dasatinib, imatinib does not inhibit SFKs, ephrins, BTK, and Syk. A hypothesis also suggests that imatinib induces bleeding disorders because of BCR-ABL rearrangements in megakaryocytic cell lines, leading to clonal expansion of dysfunctional megakaryocytes.⁴⁰

Even if ponatinib induces very few bleeding disorders, assessment of primary hemostasis in CML patients demonstrated that ponatinib induces defect in platelet aggregation. This impairment was found at all ponatinib dosage, in patients with or without low platelet counts.⁴¹ These results

^a T315I: Substitution at position 315 in BCR-ABL from a threonine to an isoleucine. This substitution alters the structure of the ATP-binding pocket and eliminates a crucial hydrogen bond required for binding of first- and second-generation TKIs.

^b Ephrin family: Members of this family are involved in platelet spreading, adhesion to fibrinogen and platelet secretion.

Table 1 In vitro and ex vivo investigations of the effects of BCR-ABL TKIs on platelet production and functions

Endpoints	Methods	Models	TKIs	Findings	Ref.
Platelet production	Platelet count	Murine whole blood	Dasatinib	Thrombocytopenia ↗ platelet production	33
	Flow cytometry (DNA ploidy) Migration assay (Dunn chamber)	Megakaryocyte primary culture	Dasatinib	↗ megakaryocyte differentiation ↗ megakaryocyte migration ↗ proplatelet formation	33
Platelet aggregation	Born aggregometry; Light transmission aggregometry	Washed human platelet	Imatinib	= CRP-, collagen- and thrombin-induced platelet aggregation	38,39,42
	Light transmission aggregometry	Human platelet (PRP)	Imatinib	↗ ADP-induced platelet aggregation ↗ collagen- and CRP-induced platelet aggregation	34
Platelet activation	Light transmission aggregometry, immunostaining (PAC-1)	Human platelet (PRP); patient blood	Dasatinib	↗ ADP-, collagen-, thrombin- and CRP-induced platelet aggregation	34,35,38
	Light transmission aggregometry; Born aggregometry	Human platelet (PRP); Washed human platelet	Nilotinib	= platelet aggregation	34,39,42
	Born aggregometry	Washed human platelet	Ponatinib	↗ CRP-induced platelet aggregation = thrombin-induced platelet aggregation	42
	Immunostaining (PS)	Washed human platelet	Imatinib	= PS exposure	42
	Western blot	Human platelet lysate	Imatinib	= Src, Lyn, LAT, and BTK activation	42
	Immunostaining (PS)	Patient blood	Dasatinib	↗ PS exposure	35
	Immunostaining (PS)	Washed human platelet	Nilotinib	= PS exposure	42
	Immunostaining (PS)	Patient blood	Nilotinib	↗ PS exposure	35
	Western blot	Human platelet lysate	Nilotinib	= Src, Lyn, LAT and BTK activation	42
	Immunostaining (PS)	Patient blood	Bosutinib	↗ PS exposure	35
Granule release	Immunostaining (PS)	Washed human platelet, patient blood	Ponatinib	↗ PS exposure	35,42
	Western blot	Human platelet lysate	Ponatinib	↗ Src, Lyn, LAT and BTK activation	42
	Immunostaining (P-selectin)	Human platelet	Imatinib	↗ thrombin-, PAR-1- and CRP-mediated α-granule release = PAR-4-mediated α-granule release	34
	Immunostaining (P-selectin)	Washed human platelet	Imatinib	= α-granule release	42
	Immunostaining (P-selectin)	Human platelet	Dasatinib	↗ thrombin-, PAR-1-, PAR-4- and CRP-mediated α-granule release	34
	Immunostaining (P-selectin)	Washed human platelet	Nilotinib	= PAR-4-, CRP- and thrombin-mediated α-granule release	34,42

Table 1 (Continued)

Endpoints	Methods	Models	TKIs	Findings	Ref.
	Immunostaining (P-selectin)	Murine platelet	Nilotinib	➔ CRP-, PAR-4- and thrombin-mediated α-granule release	34
	Immunostaining (P-selectin)	Human platelet	Nilotinib	➔ PAR-1-mediated α-granule release	34
	Immunostaining (P-selectin)	Washed human platelet	Ponatinib	➔ α-granule release	42
	Microscopy (platelet spreading)	Washed human platelet	Imatinib	= platelet spreading and lamellipodia formation	42
Platelet spreading	Microscopy (platelet spreading)	Washed human platelet	Nilotinib	= platelet spreading and lamellipodia formation	42
	Microscopy (platelet spreading)	Washed human platelet	Ponatinib	➔ platelet spreading and lamellipodia formation	42
	In vitro flow study, PFA-100	Human blood, murine whole blood	Imatinib	= platelet deposition and thrombus volume = closure time	34,36,44
	Ex vivo and in vitro flow study	Murine whole blood, human whole blood	Imatinib	➔ thrombus volume and aggregate formation	34,42
Thrombus formation	In vitro and ex vivo flow study	Human blood, murine whole blood, patient whole blood	Dasatinib	➔ thrombus volume and platelet deposition	34-36
	PFA-100	Human whole blood	Dasatinib	➔ closure time (collagen/epinephrine activation) = closure time (collagen/ADP activation)	44
	Ex vivo flow study	Murine whole blood, patient whole blood	Nilotinib	➔ thrombus volume (growth and stability)	34
	In vitro flow study	Human whole blood, murine whole blood	Nilotinib	= platelet deposition and thrombus volume	34,36,42
	In vitro flow study	Human blood	Bosutinib	➔ platelet deposition (late)	36
	PFA-100	Patient blood	Ponatinib	➔ closure time	41
	In vitro flow study	Human whole blood	Ponatinib	➔ aggregate formation	42

Abbreviations: ADP, adenosine diphosphate; BTK, Bruton's tyrosine kinase; CRP, C-reactive protein; DNA, deoxyribonucleic acid; LAT, linker for activation of T-cells; PAR, protease-activated receptor; PFA, platelet function assay; PRP, platelet-rich plasma; PS, phosphatidyl serine.

Fig. 1 Signaling pathways supporting platelet adhesion, activation, and aggregation. Tyrosine kinases are involved in several pathways and contribute to platelet adhesion, aggregation, and activation. Important players in platelet signaling are members of the Src family kinases; particularly Lyn, Fyn, and cSRC. These three tyrosine kinases are inhibited by dasatinib which might explain platelet dysfunction encountered with this treatment. Additionally, dasatinib also inhibits BTK, Syk, EphA4, and EphB1—four tyrosine kinases involved in platelet activation and aggregate stabilization. 5HT, 5-hydroxytryptamine; ADP, adenosine diphosphate; Btk, Bruton's tyrosine kinase; Ca, calcium; Eph, ephrin; FcR, Fc receptor; GP, glycoprotein; PAR, protease-activated receptor; PI3K, phosphoinositide 3-kinase; PLC, phospholipase C; TXA2, thromboxane A2; vWF, Von Willebrand factor.

were in accordance with in vitro studies which previously demonstrated similar characteristics than dasatinib (i.e., decrease of platelet spreading, aggregation, P-selectin secretion, and phosphatidylserine exposure).^{35,42} However, in vitro assays tested ponatinib at 1 μ M, a dose far higher than the concentration observed in patients on treatment.⁴³ Nilotinib and bosutinib are not associated with bleeding disorders in CML patients. First in vitro studies demonstrated little or no effect on platelet aggregation and activation with these two TKIs.^{36,39,44} However, recent experiments described prothrombotic phenotype of platelets induced by nilotinib, with increase of PAR-1^c-mediated platelet secretion, adhesion, and activation, without disturbing platelet aggregation.³⁴ Additional studies demonstrated that nilotinib increases secretion of adhesive molecules as well as thrombus formation and stability ex vivo.³⁴

To summarize, dasatinib and imatinib induce hemorrhagic events through alteration of platelet functions, but the

molecular mechanism needs to be better determined. Ponatinib also impairs platelet functions. Therefore, no current data involve platelets in the pathogenesis of arterial thrombosis occurring with dasatinib and ponatinib. Oppositely, nilotinib might induce arterial thrombosis through alteration of platelet secretion, adhesion, and activation.

Metabolic Dysregulation

Glucose Metabolism

BCR-ABL TKIs have contradictory effect on glucose metabolism. Imatinib and dasatinib improve glucose metabolism and type 2 diabetes management in CML patients (i.e., decrease of antidiabetic drug dosage and reversal of type 2 diabetes).^{14,45–49} This clinical profile is in accordance with in vivo studies in which imatinib is effective to prevent the development of type 1 diabetes in prediabetic mice, without impacting the adaptive immune system.⁵⁰ Therefore, imatinib is currently tested in clinical trials for patients suffering from type 1 diabetes mellitus (NCT01781975). The mechanism(s) by which dasatinib and imatinib improve glucose

^c PAR-1: protease-activated receptor 1. PAR receptors mediate cellular effects of thrombin in platelets and endothelial cells.

Fig. 2 Effects of BCR-ABL TKIs on glucose metabolism. Imatinib and dasatinib possess hypoglycemic effects, whereas nilotinib increases blood glucose level and diabetes development. The figure describes glucose metabolism and boxes contain emitted hypotheses for effects of imatinib, dasatinib, and nilotinib on glucose metabolism. Four major hypotheses have been emitted including impact on insulin production by β -cells, β -cell survival, peripheral insulin sensitivity, and hepatic glucose production. ABL, Abelson; FAK, focal adhesion kinase; GLUT, glucose transporter; IRS-1, insulin receptor substrate 1; JNK, c-Jun N-terminal kinases; MEKK1, MAPK/ERK kinase kinase 1; NF- κ B, nuclear factor-kappa B; PDK1, pyruvate dehydrogenase kinase 1; PI3K, phosphoinositide 3-kinase; ROS, reactive oxygen species.

metabolism remains unknown. Global hypotheses suggest that imatinib increases peripheral insulin sensitivity, promotes β -cell survival, or decreases hepatic glucose production (**Fig. 2**).^{51–54} This latter hypothesis (i.e., decreased hepatic glucose production by imatinib) is not currently the preferred theory, whereas it was demonstrated that imatinib weakly affects hepatic glucose production.⁵¹ Several targets might be involved in this metabolic effect. PDGFR has already been linked with type 1 diabetes reversal.⁵⁰ Hägerkvist et al hypothesized that c-Abl inhibition by imatinib promotes β -cell survival through activation of NF- κ B signaling and inhibition of proapoptotic pathways (**Fig. 2**).^{53,54} Inhibition of c-Abl in β -cells might also increase insulin production and contribute to the glucose regulation by imatinib.⁵⁵ It was also speculated that imatinib decreases insulin resistance in peripheral tissues due to c-Abl-dependent JNK inactivation.^{d,51} Similar hypotheses might be translated to dasatinib because of the similar off-target inhibitory profile (i.e., dasatinib also inhibits c-Abl and

PDGFR). It was hypothesized that imatinib and dasatinib impact glucose metabolism through reduced adipose mass.^{51,56} However, clinical data do not demonstrate weight loss in CML patients and do not favor this hypothesis. In both imatinib- and dasatinib-treated patients, increased circulating adiponectin^e level correlates with decreased insulin resistance.^{57,58} This correlation might be explained by the translocation of the glucose transporter GLUT4^f from the cytoplasm to the cell membrane following adiponectin signaling.⁵⁹ Additionally, adiponectin has been related to decreased hepatic glucose production which could be an additional mechanism by which imatinib and dasatinib improve glucose metabolism.⁶⁰ It was speculated that the raise of adiponectin level with imatinib and dasatinib is the consequence of increased adipogenesis subsequent to PDGFR inhibition.⁶¹

^e Adiponectin is a protein regulating glucose metabolism. Adiponectin increases peripheral insulin sensitivity by improving glucose uptake.

^f GLUT4: Glucose transporter type 4. GLUT4 is an insulin-regulated glucose transporter expressed in peripheral tissues.

^d JNK: c-Jun N-terminal kinases. JNK is responsive to stress stimuli and mediates insulin resistance through inhibition of insulin receptor substrate.

Fig. 3 Effects of BCR-ABL TKIs on lipid metabolism. Several hypotheses have been emitted to explain the imatinib-induced hypolipidemic effect. Imatinib regulates expression of genes involved in lipid metabolism: ApoBec1 that regulates ApoB expression through the introduction of a stop codon into ApoB mRNA (ApoB is essential for VLDL production), and LDLR that is implicated in lipid clearance. Imatinib-induced PDGFR inhibition influences LPL synthesis and dysregulates LRP. Dasatinib and nilotinib increase cholesterol plasma level through an unknown mechanism. Global hypotheses can be emitted and include increased hepatic lipid synthesis (possibly related to hyperinsulinemia) and decreased lipid clearance through LDLR functional defect or decreased LPL synthesis. ABC, ATP-binding cassette; C, cholesterol; CETP, cholesteryl ester transfer protein; CM, chylomicron; FA, fatty acid; HMGCoA reductase, hydroxymethylglutaryl-CoA reductase; IDL, intermediate-density lipoprotein; LDL, low-density lipoprotein; LDLR, low-density lipoprotein receptor; LPL, lipoprotein lipase; LRP, lipoprotein receptor-related protein; PDGFR, platelet-derived growth factor receptor; VLDL, very low-density lipoprotein.

Oppositely to imatinib and dasatinib, case reports and clinical trials indicate that nilotinib increases blood glucose level and promotes diabetes mellitus.^{62–65} Indeed, 20% of nilotinib-treated patients developed diabetes after 3 years of treatment,⁶⁵ whereas 29% of patients suffer from increase of fasting glucose after 1 year of therapy.⁶⁴ However, no variations of glycated hemoglobin were reported.^{64,65} Clinical data indicate no direct effect of nilotinib on β -cells, but suggest fasting insulin increase, fasting C-peptide decrease, and an increase of HOMA-IR values (i.e., a model to assess insulin resistance).^{64,66,67} Therefore, the preferred hypothesis to explain the development of hyperglycemia is the manifestation of insulin resistance. Weakened insulin secretion occurred sometimes, but it is likely that this impairment is the consequence of β -cell exhaustion.⁶⁸ However, in vitro experiments demonstrated inhibitory effect of nilotinib on pancreatic cell growth.⁶⁹ Breccia et al proposed an additional hypothesis linking development of hyperglycemia and body mass index. They suggested that the development of hyperglycemia might be the consequence of increase fat level tissue resulting in decrease peripheral insulin sensitivity.⁷⁰ However, dietetic measures to restrict glucose exogenous

uptake in patients who developed hyperglycemia were not successful,⁶³ and nilotinib does not induce changes in patient body weight.⁷¹ Little is known regarding the mechanism by which nilotinib induces insulin resistance. Racil et al suggested that peripheral insulin resistance is mediated by c-Abl inhibition which is involved in insulin receptor signaling ($\rightarrow$ Fig. 2).⁶⁷ This hypothesis is contrary to the hypothesis described with dasatinib and imatinib in which c-Abl enhances insulin sensitivity through c-Abl inhibition. These two hypotheses describe different pathways involving c-Abl but with opposite outcomes. To date, no hypothesis is preferred and additional studies are required to understand the opposite effect on glucose metabolism between TKIs, whereas both have been attributed to c-Abl inhibition. Interestingly, Frasca et al described opposite role of c-Abl in insulin signaling depending on the receptor involved, the signaling pathway, and the cell context.⁷² Similar investigations should be performed in the context of c-Abl inhibition by BCR-ABL TKIs. For bosutinib and ponatinib, little is known regarding their impact on glucose metabolism, but no drastic changes in glucose profile has been reported during clinical trials.

Lipid Metabolism

Similarly with glucose metabolism, effects on lipid metabolism are conflicting between TKIs. Oppositely to *in vivo* study which demonstrated no impact of imatinib on total cholesterol and triglycerides levels in diabetic mice,²⁹ imatinib is associated in CML patients with a rapid and progressive decrease of cholesterol and triglycerides levels.^{66,73–75} First hypothesis relates the inhibition of PDGFR by imatinib (► **Fig. 3**). PDGFR is involved in the synthesis of the lipoprotein lipase (LPL) and in the regulation of the lipoprotein receptor-related protein (LRP).^{73,74} However, all BCR-ABL TKIs possess inhibitory activity against PDGFR but do not share this positive impact on lipid profile. Recently, Ellis et al described that imatinib impairs gene expression of proteins involved in plasma lipid regulation. Indeed, in *in vitro* model of CML cells, imatinib affects gene expression of four genes implicated in lipid synthesis (HMG-CoA reductase^g gene and apobec1^h), lipid clearance (LDLR geneⁱ) and in exchange of lipids from very low-density lipoprotein (VLDL) or low-density lipoprotein (LDL) to high-density lipoprotein (HDL) (CETP^j gene). However, these studies were performed in a model of CML cells and need to be confirmed in more relevant models (e.g., primary cell lines, hepatocytes).⁷⁶ Franceschino et al suggested that imatinib decreases diarrhea-related lipid absorption due to inhibition of c-kit in interstitial Cajal cells (i.e., c-kit signaling is critical for the survival and development of these cells).⁷³ However, this hypothesis is unlikely, few patients (3.3%) developed grade 3/4 diarrhea, and patients treated with interferon- α and cytarabine developed diarrhea at a same rate and do not present lipid level reduction in the phase 3 clinical trial (NCT00333840).

Oppositely, dasatinib and mostly nilotinib are associated with an increase of cholesterol level.^{26,66,77} Nilotinib induces quick rise of total cholesterol, HDL, and LDL (i.e., within 3 months). Nilotinib-induced dyslipidemia are responsive to statin and lipid level normalized after nilotinib discontinuation.⁷⁸ To date, the mechanism by which dasatinib and nilotinib impact lipid metabolism is unknown. Future researches should determine how these treatments induce dyslipidemia. Global hypotheses could be formulated and include an increase of lipid synthesis that might be secondary to insulin resistance and hyperinsulinemia. This hypothesis is particularly relevant with nilotinib and it is also associated with hyperglycemia. Dasatinib and nilotinib might also decrease blood lipid clearance (e.g., disturbance of LDLR and LPL synthesis). The development of dyslipidemia might contribute to the occurrence of arterial occlusive events that occurred with nilotinib and dasatinib.

However, the relationship between impaired lipid metabolism and cardiovascular occlusive events is unknown with BCR-ABL TKIs, and there is no indication that correct management of lipid metabolism can prevent arterial thrombosis (e.g., stenosis occurred in a nilotinib-treated patient despite the management of its hyperlipidemia through statin treatment).⁷⁹ On their side, bosutinib and ponatinib do not disturb lipid metabolism.^{78,80}

Effects on Atherosclerosis

Endothelial Dysfunction

► **Fig. S1** in the Supplementary Material details the role of endothelial cells (ECs) in atherosclerosis. Several *in vitro* and *in vivo* experiments assess the impact of imatinib on EC viability and functions (► **Table 2**). These studies demonstrate that imatinib does not affect EC viability nor induce apoptosis but increases EC proliferation.^{39,81–84} Only one study reports a proapoptotic effect of imatinib on ECs, but their experiments were performed on a cell line (i.e., EA.hy926 cells),⁸⁵ a model less reliable than primary cultures (e.g., HUVEC,^k HCAEC^l). *In vitro* studies also assessed the effect of imatinib on EC functions. In these studies, imatinib does not influence adhesion molecule expressions (i.e., ICAM-1^m and VCAM-1ⁿ), EC migration, reactive oxygen species (ROS) production, nor angiogenesis.^{81,82,85–87} Letsiou et al suggested that imatinib decreases EC inflammation by decreasing the secretion of proinflammatory mediators.⁸⁶ The impact of imatinib on endothelial permeability is not clear. Indeed, *in vitro* studies demonstrate that imatinib increases endothelial permeability by decreasing the level of plasma membrane VE-cadherin,^{o,85,86} whereas *in vivo* experiments indicate decreased vascular leak following imatinib treatment in a murine model of acute lung injury.⁸⁸ Additionally, imatinib has been tested in patients suffering from acute lung injury, a disease characterized by vascular leakage, and demonstrate promising clinical efficacy. Therefore, imatinib might positively affect atherogenesis by decreasing endothelial inflammation and reducing vascular leakage.

Nilotinib and ponatinib reduce EC proliferation and might impair endothelial regeneration.^{39,82,89,90} Additionally, ponatinib induces EC apoptosis, although it is well recognized that high glucose concentration induces EC death,⁹¹ suggesting that nilotinib might, by this intermediary, affect EC viability. Moreover, clinical data indicate that dasatinib induces pulmonary arterial hypertension, whereas imatinib is possibly beneficial in this disease.^{92,93} This pathology is initiated by dysfunction or injury of pulmonary ECs.⁸⁷

^g HMGCoA reductase: 3-hydroxy-3-methyl-glutaryl-coenzyme A reductase. HMGCoA reductase catalysis is the conversion of HMG-CoA to mevalonic acid, an essential step in cholesterol synthesis.

^h Apobec1: Apolipoprotein B mRNA editing enzyme catalytic subunit 1. Apobec1 introduces a stop codon into ApoB mRNA.

ⁱ LDLR: Low-density lipoprotein receptor. This cell surface receptor mediates LDL endocytosis.

^j CETP: Cholesteryl ester transfer protein.

^k HUVEC: Human umbilical vein endothelial cells.

^l HCAEC: Human coronary artery endothelial cells.

^m ICAM-1: Intercellular adhesion molecule 1. ICAM-1 stabilizes leukocyte-endothelial cell adhesion and facilitates leukocyte transmigration.

ⁿ VCAM-1: Vascular cell adhesion molecule 1. VCAM-1 mediates rolling-type and firm adhesion of leukocyte.

^o VE-cadherin: Vascular endothelial cadherin. VE-cadherin is a cell-cell adhesion molecules and implies in endothelial junctions.

Table 2 In vivo and in vitro investigations of the effects of BCR-ABL TKIs on endothelial cell viability and major functions

Endpoints	Methods	Models	TKIs	Findings	Ref.
EC proliferation/ survival	Cell counting; trypan blue staining	EA.hy 926 cell; HCAEC	Imatinib	= EC viability <10µM	84,85
	Caspase assay; Annexin V staining; Hoechst staining; TUNEL assay	HMEG-1; HUVEC; Human pulmonary EC; Mouse EC	Imatinib	= EC apoptosis	81,82,87
	TUNEL assay; Annexin V staining	EA.hy 926 cell	Imatinib	↗ EC apoptosis	85
	MTT cell proliferation assay; ³ H-thymidine incorporation; WST-1 assay; cell counting	HMEG-1; HUVEC; HCAEC	Imatinib	= EC proliferation	39,81,82,84
	Resazurin proliferation assay; PCNA expression	HUVEC; BAEC	Imatinib	↗ EC proliferation (≥1.2 µM)	83
Oxidative stress	Caspase assay; Hoechst staining; Annexin V staining; TUNEL assay	Human pulmonary EC	Dasatinib	↗ EC apoptosis	87
	³ H-thymidine incorporation; WST-1 assay; MTT assay	HUVEC; HCAEC; HMEG-1; HCTAEC	Nilotinib	↗ EC proliferation	39,82,89
	Annexin V staining	HUVEC	Nilotinib	= EC apoptosis	82
	Caspase assay; Annexin V staining	HCAEC; HUVEC	Ponatinib	↗ EC apoptosis	82,90
	³ H-thymidine incorporation; WST-1 assay	HUVEC; HMEG-1; EPC	Ponatinib	↗ EC proliferation	82,90
EC migration	Fluorescent ROS detection; Immunofluorescence (8-oxo-dG)	Human Pulmonary EC; Rat lung	Imatinib	= endothelial ROS	87
	Fluorescent ROS detection; Immunofluorescence (8-oxo-dG)	Human Pulmonary EC; Rat lung	Dasatinib	↗ endothelial ROS	87
	Wound scratch assay; Microchemotaxis assay; Transwell migration assay	HMEG-1; HUVEC; EA.hy 926 cell; HCAEC	Imatinib	= EC migration	81,82,84,85
	Wound scratch assay	HUVEC; HCAEC; HMEG-1	Nilotinib	↗ EC migration	39
	Transwell migration assay	HUVEC	Nilotinib	= EC migration	82
Angiogenesis	Transwell migration assay	HUVEC	Ponatinib	↗ EC migration	82
	Tube-formation assay	HMEG-1; HUVEC	Imatinib	= angiogenesis	81,82
	Tube-formation assay	HUVEC; HCAEC; HMEG-1	Nilotinib	↗ angiogenesis	39
	Tube-formation assay	HUVEC	Nilotinib	= angiogenesis	82
	Tube-formation assay	HUVEC	Ponatinib	↗ angiogenesis	82
Permeability	Permeability to albumin	EA.hy 926 cell	Imatinib	↗ endothelial permeability (10 µM)	85
	Immunofluorescence (VE-cadherin)	EA.hy 926 cell; HPAEC	Imatinib	↗ membrane VE-cadherin (10 µM)	85,86
	BAL protein levels	Mice (2-hit model of ALI)	Imatinib	↗ BAL protein levels	86,88
			Imatinib	= endothelial permeability	94,147

Table 2 (Continued)

Endpoints	Methods	Models	TKIs	Findings	Ref.
CAM expression	Permeability to FITC-Dextran; permeability to HRP	HMEG-1; HUVEC; Human lung micro-vascular EC			
	Immunostaining	HUVEC	Imatinib	↗ intercellular gaps	147
	Evans blue/albumin extravasation	Mice	Imatinib	↗ Evans blue extravasation	147
	Pulmonary microvascular permeability assay; permeability assay (FITC-Dextran)	Mice; HMEG-1; HPAEC	Dasatinib	↗ endothelial permeability	94
	Permeability assay (FITC-Dextran)	HRMEC	Dasatinib	↗ VEGF-induced permeability	148
	Confocal microscopy; ELISA; qRT-PCR; flow cytometry	HMEG-1; Pulmonary EC (rat lung); EA.hy926	Imatinib	= ICAM-1, VCAM-1 and E-selectin expression = soluble ICAM-1, VCAM-1 and E-selectin	81,87,149
	Immunoblotting (VCAM-1)	Human lung EC	Imatinib	↗ VCAM-1 expression	86
	Confocal microscopy	Pulmonary EC (rat lung)	Dasatinib	↗ ICAM-1, VCAM-1 and E-selectin expression	87
	ELISA	Rat	Dasatinib	↗ soluble ICAM-1, VCAM-1 and E-selectin	87
	qRT-PCR; flow cytometry	EA.hy926	Dasatinib	= ICAM-1, VCAM-1 and E-selectin expression	149
Secretory	Unknown	HUVEC	Nilotinib	↗ ICAM-1, VCAM-1 and E-selectin expression ($\geq 1 \mu\text{M}$)	39
	qRT-PCR; flow cytometry	EA.hy926	Nilotinib	↗ ICAM-1, VCAM-1 and E-selectin expression	149
	ELISA (IL-6; IL-8)	Stimulated HPAEC	Imatinib	↗ IL-8 and IL-6 (LPS induced)	86
	qRT-PCR ; ELISA (IL-1 β ; IL-6; TNF- α)	EA.hy926 cell ; HUVEC	Imatinib	= IL-1 β , IL-6 and TNF- α expression and production	149
	qRT-PCR ; ELISA (IL-1 β ; IL-6; TNF- α)	EA.hy926 cell ; HUVEC	Dasatinib	= IL-1 β , IL-6 and TNF- α expression and production	149
	qRT-PCR ; ELISA (IL-1 β ; IL-6; TNF- α)	EA.hy926 cell ; HUVEC	Nilotinib	= IL-6 and TNF- α expression and production ↗ IL-1 β expression and production	149
Adhesion	ELISA (t-PA; PAI-1; ET-1; vWF; total NO)	HCTAEC	Nilotinib	↗ t-PA ↗ PAI-1, ET-1, vWF and total NO	89
	Unknown	HUVEC	Ponatinib	↗ adhesion to plastic surface at 1 μM	90

Abbreviations: 8-oxo-dG, 8-hydroxy-2'-deoxyguanosine; ALI, acute lung injury; BAEC, bovine aortic endothelial cell; BAL, bronchoalveolar level; EC, endothelial cell; ELISA, enzyme-linked immunosorbent assay; EPC, endothelial progenitor cell; ET-1, endothelin 1; FITC, fluorescein isothiocyanate; HCAEC, human coronary artery endothelial cell; HCTAEC, human carotid artery endothelial cell; HMEG-1, human microvascular endothelial cell; HPAEC, human pulmonary artery endothelial cell; HRMEC, human retinal microvascular endothelial cells; HUVEC, human umbilical vein endothelial cell; ICAM-1, intercellular adhesion molecule 1; IL, interleukin; LPS, lipopolysaccharide; NO, nitric oxide; PAI-1, plasminogen activator inhibitor-1; ROS, reactive oxygen species; t-PA, tissue plasminogen activator; TUNEL, terminal deoxynucleotidyl transferase dUTP nick end labeling; VCAM-1, vascular cell adhesion molecule 1; VE-cadherin, vascular endothelial cadherin; vWF, Von Willebrand factor.

Therefore, in vivo and in vitro studies investigated effect of imatinib and dasatinib on pulmonary ECs and demonstrate that dasatinib induces apoptosis on pulmonary ECs mediated by increased mitochondrial ROS production.⁸⁷ Future researches should assess if this effect is also found in arterial ECs and ROS production should also be tested with other new-generation BCR-ABL TKIs.

In addition to their effect on EC viability, nilotinib and ponatinib also influence EC functions, inhibit their migration, and decrease angiogenesis.^{39,82} It was suggested that the antiangiogenic effect of ponatinib is the consequence of VEGFR^p inhibition, but this hypothesis cannot explain the antiangiogenic effect of nilotinib (i.e., nilotinib does not inhibit VEGFR).⁸² Nilotinib also increases adhesion molecule expressions (i.e., ICAM-1, VCAM-1, and E-selectin) in vitro,³⁹ suggesting that nilotinib might increase leukocyte recruitment. However, further experiments are needed to validate this hypothesis (e.g., assessment of endothelium permeability and transendothelial migration). Dasatinib also induces endothelium leakage in vitro, and the RhoA-ROCK^q pathway is involved in this phenomenon.⁹⁴ It was demonstrated that RhoA activation induces the phosphorylation of myosin light chain that increases the actomyosin contractility and disrupt endothelial barrier.⁹⁴ Therefore, increased endothelium permeability is a potential mechanism by which dasatinib and nilotinib promote atherosclerosis development and arterial thrombosis. Likewise, it is plausible that ponatinib affects endothelium integrity because of its inhibitory activity against VEGFR, which is recognized as a permeability-inducing agent. Additional hypotheses suggest that inhibition of Abl kinase (i.e., Arg^r and c-Abl) and PDGFR might also be implicated in vascular leakage.⁸⁵ Finally, Guignabert et al demonstrated that both in rats and in CML patients taking dasatinib, there is an increase of soluble adhesion molecules, which are well-known markers of endothelial dysfunction.⁸⁷

Inflammation

► **Fig. S2** in the Supplementary Material describes the role of immune cells and inflammation process during atherosclerosis. ► **Table 3** summarizes in vitro studies that investigate impacts of BCR-ABL TKIs on survival, proliferation, and major functions of monocytes, macrophages, and T-lymphocytes. Globally, in vitro studies demonstrate that imatinib inhibits the development and maturation of monocytes and alters monocyte functions.^{95,96} Imatinib decreases production of proinflammatory cytokines (i.e., TNF- α ^s and IL-6^t) and diminishes the potential of monocytes to phago-

cytose.^{97,98} These impacts on monocyte functions are possibly related to c-fms^u inhibition.⁹⁹ Imatinib also inhibits macrophage functions in vitro. Imatinib decreases lipid uptake without impacting the lipid efflux and decreases activity and secretion of two matrix metalloproteinases (MMPs; i.e., MMP-2 and MMP-9^v) on a posttranscriptional level.¹⁰⁰ Additionally, imatinib inhibits T-lymphocyte activation and proliferation and decreases proinflammatory cytokines secretion (i.e., IFN- γ ^w).¹⁰¹ The inhibition of monocyte, macrophage, and T-cell functions by imatinib might prevent the development of atherosclerosis or reduce the risk of atherosclerotic plaque rupture.

Effects of new-generation TKIs on inflammatory cells were less studied, but first experiments indicate similarities with imatinib about its impact on monocytes and macrophages. Both dasatinib and nilotinib have similar inhibitory profile on macrophage-colony formation that has been linked to CSFR inhibition.^{96,102} Dasatinib also possesses anti-inflammatory functions by attenuating proinflammatory cytokines production (i.e., TNF- α , IL-6, and IL-12^x) by macrophages and increasing production of anti-inflammatory mediator (i.e., IL-10^y).¹⁰³ These effects are thought to be mediated by SIK^z inhibition, a subfamily of three serine/threonine kinases that regulate macrophage polarization.^{103,104} Finally, dasatinib is associated with decreased T-cell functions and particularly it decreases the production of proinflammatory cytokines (e.g., TNF- α , IFN- γ) and chemotactic mediators.¹⁰⁵ Nilotinib and bosutinib also possess anti-inflammatory activity and decrease cytokine production and T-cell activation.^{103,106} Inhibition of Lck,^{ai} a tyrosine kinase implicated in T-cell receptor signaling, is implicated in the impairment of T-cell functions by dasatinib and nilotinib.^{107,108} It has been hypothesized that nilotinib decreases mast cell activity through c-kit inhibition,^{62,109} which might result in a decrease of the vascular repair system.^{39,62} Clinical profile of nilotinib in patients with CML consolidates this hypothesis and demonstrates a decreased of mast cell level.³⁹ However, similar decreased of mast cell is also reported with imatinib without high rate of arterial thrombosis.¹¹⁰

Globally, BCR-ABL TKIs possess reassuring profile on inflammatory cells. However, impact of new-generation TKIs on several functions of macrophages have not been assessed (e.g., MMP secretion and activity, lipid uptake, and

^p VEGFR: Vascular endothelial growth factor. This protein plays major roles in vasculogenesis and angiogenesis.

^q RhoA-ROCK: Ras homolog gene family, member A—Rho-associated protein kinase. Rho-kinase regulates cytoskeletal reorganization, cell migration, cell proliferation, and survival.

^r Arg: Abelson-related gene (also known as ABL2). Arg possesses cytoskeletal-remodeling functions.

^s TNF- α : Tumor necrosis factor alpha. This cytokine is mainly involved in systemic inflammation and regulates immune cells.

^t IL-6: Interleukin 6. IL-6 is a proinflammatory cytokine secreted by T-cells and macrophages to stimulate immune response.

^u CSFR: Colony-stimulating factor receptor. CSFR drives growth and development of monocytes.

^v MMP-2 and MMP-9 are two proteases capable of degrading extracellular matrix components. These two MMPs are the main proteases involved in atherogenesis.

^w IFN- γ : Interferon gamma. IFN- γ is involved in innate and adaptive immunity and activates macrophages.

^x IL-12: Interleukin 12. IL-12 is involved in T-cell differentiation and functions.

^y IL-10: Interleukin 10. IL-10 exerts immunoregulation and regulates inflammation.

^z SIK: Salt-inducible kinase. SIKs regulate production of anti- and proinflammatory cytokines.

^{ai} Lck: Lymphocyte-specific protein tyrosine kinase. Lck is mostly involved in T-cell maturation.

Table 3 In vitro studies on effects of BCR-ABL TKIs on proliferation, survival, and major functions of monocytes, macrophages, and T-lymphocytes

Endpoints	Methods	Models	TKIs	Findings	Ref.
Monocytes/Macrophages					
Proliferation/ survival	Propidium iodide staining	PBMC	Imatinib	= viability	150
	Cell counting	Ovarian tumor ascites samples	Imatinib	↗ macrophage production	96
	Cell counting	Ovarian tumor ascites samples	Dasatinib	↗ macrophage production	96
	WST-1 assay	Human macrophages	Ponatinib	= macrophage viability	82
Monocyte differentiation	Morphology assessment	Human monocyte	Imatinib	↗ differentiation into macrophages	95
Secretion	ELISA; qPCR	Human monocyte and macrophage; PBMC	Imatinib	↗ TNF- α , IL-6 and IL-8 production	97,150
	ELISA	PBMC; Human monocyte and macrophage	Imatinib	= IL-10 production	150
	ELISA; Bioplex system; nitrite assay	Raw 264.7; bone-marrow derived macrophage	Dasatinib	↗ TNF- α , IL-6, IL-12p40 and NO production	103,151
	qPCR; Bioplex system	Primary macrophage (mice)	Dasatinib	↗ IL-10 production	103
Phagocytosis	Bioplex system	Bone-marrow derived macrophage	Bosutinib	↗ IL-6, IL-12p40 and TNF- α production	103
	qPCR; Bioplex system	Primary macrophage (mice)	Bosutinib	↗ IL-10 production	103
Cholesterol uptake	Antigen-uptake assay	Human monocyte	Imatinib	↗ phagocytosis	97
	Cholesterol uptake assay	THP-1; PBMC	Imatinib	↗ LDL uptake	100
MMP produc- tion/activity	Cholesterol uptake assay	THP-1	Bosutinib	↗ LDL uptake	100
	Zymography	THP-1	Imatinib	↗ MMP-2 and MMP-9 secretion and activity	100
T Lymphocytes					
Proliferation/ survival	³ H-TdR incorporation; CFSE staining; titrated thymidine	Naïve CD4 ⁺ T cell; Human T cell	Imatinib	↗ T-cell proliferation	101,152,153
	Annexin V staining; Caspase assay	Human T cell	Imatinib	= T-cell apoptosis	101,152,153
	Annexin V staining	Human T cell	Imatinib	= T cell apoptosis	
	CFSE dye	Human T cell	Dasatinib	↗ T-cell proliferation	107
	Annexin V staining	PBMC; Human T cell	Dasatinib	= T cell viability	105,107
Secretion	CFSE dye	CD8 ⁺ T cell; PBMC	Nilotinib	↗ T cell proliferation	106,154
	ELISA	Human T cell; CD8 ⁺ and CD4 ⁺ T cell	Imatinib	↗ IFN- γ production	101,107
	ELISA; proteome profile array	Human T cell; PBMC	Dasatinib	↗ TNF- α , IFN- γ , IL-2, IL-6, IL-17 production	105,107
	Proteome profile array	PBMC	Dasatinib	↗ chemotactic factors secretion	105

(Continued)

Table 3 (Continued)

Endpoints	Methods	Models	TKIs	Findings	Ref.
Activation				(SDF-1, MIP-1α, MIP-1β, MCP-1, CXCL-1)	
	ELISPOT assay	CD8 ⁺ T cell	Nilotinib	IFN-γ production	154
	Immunofluorescence	Human T cell	Imatinib	T cell activation	101
	Flow cytometry (CD25, CD69)	Human T cell	Imatinib	= T cell activation	153
	Flow cytometry (CD25, CD69)	Human T cell; PBMC	Dasatinib	T cell activation	105, 107
	Flow cytometry (CD25, CD69)	Human T cell	Nilotinib	T cell activation	154

Abbreviations: CFSE, carboxyfluorescein succinimidyl ester; CXCL1, (C-X-C motif) ligand 1; ELISA, enzyme-linked immunosorbent assay; ELISPOT, enzyme-linked immunospot; IFN, interferon; IL, interleukin; MCP, monocyte chemoattractant protein-1; MIP-1, macrophage inflammatory protein 1; NO, nitric oxide; PBMC, peripheral blood mononuclear cell; qPCR, quantitative polymerase chain reaction; SDF-1, stromal cell-derived factor 1; TNF, tumor necrosis factor.

foam cell formation), whereas effect of ponatinib on inflammatory cells is unknown. The assessment of lipid uptake and foam cell formation is particularly relevant with new-generation TKIs because there are numerous interactions between TKIs and ABC transporters.^{aii,111,112}

Fibrous Cap Thickness

–Fig. S3 in the Supplementary Material describes the mechanism by which atherosclerotic plaque ruptures and induces arterial thrombosis. –Table 4 summarizes in vitro and in vivo experiments performed on VSMCs and fibroblasts. Imatinib decreases VSMC proliferation and growth but results are conflicting about its impact on apoptosis. Some studies demonstrate no impact on SMC apoptosis, whereas others indicate increased SMC death.^{83,113–116} Imatinib also affects VSMC functions and decreases their migration and LDL binding, inducing decreased LDL retention by the sub-endothelium.^{113,117} Imatinib also exerts negative effect on the synthesis of major ECM components (type I collagen and fibronectin A) by fibroblasts, correlating to decreased ECM accumulation in vivo.¹¹⁸ The impact of imatinib on SMCs is thought to be mediated by PDGFR inhibition,¹¹⁴ which is involved in several VSMC functions including VSMC survival and plasticity.¹¹³ Subsequent to the hypothesis that imatinib inhibits PDGFR signaling, prevents abundant SMC and fibroblast proliferation, and inhibits abundant ECM accumulation, imatinib has been tested for the management of several fibrotic diseases (e.g., dermal and liver pulmonary fibrosis, systemic sclerosis).^{30,118,119} Imatinib successfully acts on pulmonary fibrosis and pulmonary arterial hypertension (i.e., a disease involving vascular remodeling mediated by pulmonary SMC proliferation),^{93,114} and has beneficial activity in sclerotic chronic graft-versus-host disease.¹²⁰ Finally, imatinib was tested in vivo for the prevention of cardiovascular diseases and demonstrates efficacy for the treatment of myocardial fibrosis by reducing ECM component synthesis (i.e., procollagen I and III).³⁰ In a rat model, imatinib successfully inhibits stenosis after balloon injury and presents interest in intimal hyperplasia and stenosis after bypass grafts.^{115,116,121–123} Imatinib also successfully prevents arterial thrombosis following microvascular surgery in rabbits.¹²⁴ Imatinib was also encompassed in a stent but do not demonstrate efficacy in restenosis prevention.⁸⁴

Impact of new-generation TKIs on fibrosis was less studied but demonstrate similar inhibitory effect on VSMCs and fibroblasts. Indeed, dasatinib inhibits PDGFR more potently than imatinib,¹¹³ and the hypothesis that dasatinib prevents restenosis similarly with imatinib was emitted. Therefore, a patent has been filed claiming the use of dasatinib for the prevention of stenosis and restenosis.¹²⁵ Compared with imatinib, dasatinib has additional off-targets and is able to

^{aii} ABC transporter: ATP-binding cassette transporters. ABCG1 and ABCA1 are implicated in macrophage reverse cholesterol transport. Their deficiency leads to foam cell formation and atherosclerosis development.

Table 4 In vitro and in vivo studies on effects of BCR-ABL TKIs on proliferation, survival, and major functions of smooth muscle cells and fibroblasts

Endpoints	Methods	Models	TKIs	Findings	Ref.
Proliferation/ survival	Resazurin assay; immunofluorescence; ³ H-thymidine incorporation; BrdU incorporation; MTT assay	HVSMC; BAoSMC; PASMC; ASMC; VSMC; HAoSMC; HCASMC; Rabbit	Imatinib	➔ SMC proliferation	83,84,114–116,123,155
	Caspase assay; PARP (Western blot); JC-1 dye; Annexin V staining	BAoSMC; Dermal fibroblast; PASMC	Imatinib	= SMC/fibroblast apoptosis	83,118,155
	TUNEL; caspase assay	PASMC; HAoSMC; Rabbit	Imatinib	➔ SMC apoptosis (PDGF-stimulated)	114,116,123
	Trypan blue exclusion	HCASMC; A10 cell line	Imatinib	= SMC viability	84
Migration	Cell counting; Propidium iodide staining	A10 cell line, HAoSMC	Dasatinib	➔ SMC proliferation	113,125
	Transwell cell migration assay	HAoSMC; PASMC; HCASMC; A10 cell	Imatinib	➔ SMC migration	84,116,155
	Transwell cell migration assay	HAoSMC; A10 cell	Dasatinib	➔ SMC migration	113,125
	Radiolabel incorporation	Human VSMC	Imatinib	➔ proteoglycan synthesis	117
Secretion/ synthesis	RT-PCR; Western blot; Sircol collagen assay	Dermal fibroblast	Imatinib	➔ COL1A1, COL1A2, fibronectin 1 synthesis ➔ collagen synthesis	118
	RT-PCR	Dermal fibroblast	Imatinib	= MMP-1, MMP-2, TIMP-1, TIMP-2, TIMP-3 and TIMP-4	118
	qRT-PCR	Human fibroblast	Nilotinib	Decreases COL1A1 and COL1A2 synthesis	127
	Sirius red staining	Rat	Imatinib	➔ myocardial fibrosis, liver fibrosis	30,119
Fibrosis	Intima/media ratio	Rat (Balloon injury model)	Imatinib	➔ stenosis	121,122
	Intima/media ratio	Rabbit	Imatinib	➔ intimal thickness	124
	Hydroxyproline, collagen content	Rat liver	Imatinib	➔ hydroxyproline and collagen content	128
	Hydroxyproline, collagen content	Rat liver	Nilotinib	➔ hydroxyproline and collagen content	128
	Sirius red staining	Rat liver	Nilotinib	➔ liver fibrosis	128

Abbreviations: ASMC, arterial smooth muscle cell; BAoSMC, bovine aortic smooth muscle cell; BrdU, bromodeoxyuridine; COL, collagen; HAoSMC, human aortic smooth muscle cell; HCASMC, human coronary artery smooth muscle cell; HVSMC, human vascular smooth muscle cell; MMP, matrix metalloproteinase; PARP, poly(ADP-ribose) polymerase; PASMC, pulmonary smooth muscle cell; PDGF, platelet-derived growth factor; qRT-PCR, quantitative reverse transcription polymerase chain reaction; SMC, smooth muscle cell; TIMP, tissue inhibitor of metalloproteinase; TUNEL, terminal deoxynucleotidyl transferase dUTP nick end labeling; VSMC, vascular smooth muscle cell.

Fig. 4 Specificity of imatinib, dasatinib, nilotinib, and ponatinib toward tyrosine kinases. Green, yellow, red, and blue circles contain tyrosine kinase inhibited by dasatinib, nilotinib, bosutinib, and ponatinib, respectively. Tyrosine kinases in white represent imatinib off-targets. This figure summarizes results from 13 experiments.^{39,43,130,132–137,156–159} In case of conflictual results between studies, a conservative approach has been applied. Additional information is provided in the Supplementary Material.

inhibit Src,^{aiii} a kinase involved in dermal fibrosis in addition to PDGFR.¹²⁶ Therefore, dasatinib was tested in patients with scleroderma-like chronic graft-versus-host disease, a disease resulting from inflammation and progressive fibrosis of the dermis and subcutaneous tissues, and first results are encouraging.¹²⁶ Nilotinib also appears to be clinically efficient in scleroderma-like graft-versus-host disease by reducing collagen expression.¹²⁷ Finally, nilotinib was tested in vivo for the treatment of liver fibrosis and demonstrates decreased fibrotic markers and inflammatory cytokines (IL-1 α , IL-1 β , IFN- γ , IL-6).¹²⁸ However, only low-dose nilotinib was found to be efficient against fibrosis and normalized collagen content.¹²⁸ This lack of antifibrotic effect at higher doses might be explained by inhibition of additional off-targets by nilotinib that affect the benefit of low-dose nilotinib against fibrosis. Arterial thrombosis occurring with dasatinib and nilotinib are probably not the consequence of VSMC impairment, but investigations should be performed on VSMCs rather than on fibroblasts. Additional investigations are warranted to complete impact of BCR-ABL TKIs on VSMC functions (e.g., VSMC apoptosis, proliferation, and migration) and confirm their safety toward VSMCs.

^{aiii}Src is involved in angiogenesis and cell survival and proliferation.

Off-targets

BCR-ABL TKIs bind the highly conserved ATP binding site and are therefore not very specific to BCR-ABL and possess multiple cellular targets (kinases and nonkinase proteins).^{129,130} This allowed the possibility to exploit them in other indications (e.g., PDGFR inhibition by imatinib is used in BCR-ABL-negative chronic myeloid disorders),¹³¹ but this may also induce toxicities and side effects.¹²⁹ The development of arterial thrombotic events with new-generation BCR-ABL TKIs is likely to be related to inhibition of off-targets, as described throughout this review. ► **Fig. 4** describes inhibitory profiles of imatinib, dasatinib, nilotinib, bosutinib, and ponatinib. Globally, imatinib is the most selective BCR-ABL TKIs, whereas dasatinib and ponatinib inhibit numerous off-targets.

However, inhibitory profiles are difficult to determine and several researches published discrepancies. For conflicting results, a conservative approach has been applied in ► **Fig. 4**, but supplementary information (► **Table S2**) describes the tyrosine kinase selectivity profile of the five BCR-ABL TKIs and indicates divergences between studies.^{43,130,132–134} These discrepancies can be explained by the difference in drug concentration and methodologies. To date, several methods have been used to determine inhibitory profile of BCR-ABL TKIs including in vitro kinase assay,^{133–135} kinase expression in bacteriophages,¹³⁶ and

affinity purification methods combined with mass spectrophotometry.^{130,132} However, all these methods suffer from caveats, including the incompatibility to perform live-cell studies. A cell-permeable kinase probe was developed to figure out this problem, but this assay is still limited by the number of off-target tested (i.e., it requires to predefine tested off-targets) and therefore, the missing of targets is possible.¹³⁷ For this reason, the inhibitory activity of each TKI has not been tested toward all tyrosine kinase and ▶ **Fig. 4** includes only off-targets for which at least one of the five BCR-ABL TKI has been tested. Thus, inhibitory profiles need to be carefully considered and it has to keep in mind that BCR-ABL TKI metabolites may possess activity against supplemental off-targets.

As described over this review, PDGF signaling has countless effects on several cells and tissues and is involved in several proatherogenic mechanisms (e.g., adipogenesis, vascular leakage, VSMC viability, and functions) and vascular homeostasis, which led to the suggestion of its implication in the potential beneficial cardiovascular effect of imatinib.^{116,123,138} However, dasatinib, nilotinib, and ponatinib also inhibit PDGFR but increase the risk of arterial occlusive events. This difference of clinical outcome might be explained by the concentration of BCR-ABL TKIs necessary to obtain a same degree of PDGFR inhibition.⁴³ Indeed, Rivera et al reported that when adjusted to the maximum serum concentration, imatinib inhibits more profoundly PDGFR than dasatinib, nilotinib, and ponatinib.⁴³ Therefore, at effective concentration, it is probable that the degree of PDGFR inhibition is too low with dasatinib, nilotinib, and ponatinib to obtain the beneficial effect of PDGFR inhibition on atherosclerosis. Another possible hypothesis concerns the less conclusive specificity of new-generation TKIs which leads to inhibition of additional off-targets that might counterbalance the positive effect of PDGFR inhibition.

Other tyrosine kinases have been incriminated in the occurrence of arterial thrombosis with new-generation TKIs. DDR-1^{aiv} possesses functions in vascular homeostasis, atherogenesis, and is expressed in pancreatic islet cells. However, and similarly with PDGFR, it is inhibited by all BCR-ABL TKIs.^{26,62} Other hypotheses include impairment of VEGF signaling by ponatinib^{43,90} or the inhibition of several ephrin receptors by new-generation TKIs but not by imatinib which might inhibit monocyte recruitment.¹³⁹ Finally, it has been suggested that the inhibition of c-Abl itself is implicated in the increase of the cardiovascular risk. Indeed, imatinib possesses lower inhibitory effect on c-Abl than new-generation TKIs, which might further explain the difference in cardiovascular safety.⁴³ Additionally, c-Abl modulates Tie-2,^{av} a tyrosine kinase that possesses impor-

tant effect on endothelial cell function, angiogenesis, and inflammation.^{140,141}

Perspectives and Conclusions

This review summarizes the data underlying the potential preventive effect of imatinib on the occurrence of arterial thrombosis. Globally, in vitro and in vivo experiments demonstrate that imatinib possesses antiplatelet activity, hypolipidemic and hypoglycemic effects, and inhibits inflammation and atherosclerosis development in several cell types (i.e., decreases of inflammatory cell and VSMC functions and increased vascular permeability). These benefits were largely attributed to PDGFR inhibition. It is currently unknown why new-generation TKIs that also inhibit PDGFR present opposite cardiovascular safety profile and this point needs to be elucidated.

New-generation BCR-ABL TKIs increase the risk of arterial thromboembolism with different clinical features (e.g., time-to-event and absolute rate) and are associated with different safety profiles, suggesting different pathways to explain the pathophysiology. The safety profile of nilotinib is mostly characterized by impaired glucose and lipid metabolism. However, both the molecular mechanism of these alterations and their impact on the occurrence of arterial thrombosis are unknown. Both dasatinib and ponatinib exhibit antiplatelet effect, whereas it was recently suggested that nilotinib potentially induces prothrombotic phenotype of platelets. Based on the clinical characteristics and case reports, atherosclerosis appears the most plausible mechanisms by which new-generation TKIs induce arterial thrombosis. However, in vitro and in vivo studies of viability and functions of SMCs and inflammatory cells demonstrate reassuring impact of dasatinib and nilotinib, even if additional studies are required to complete this evaluation. However, first experiments indicate that dasatinib, nilotinib, and ponatinib influence EC survival and/or endothelium integrity, suggesting a reasonable hypothesis by which new-generation TKIs induce atherosclerosis development and, subsequently, arterial thrombosis. Additional studies on the shedding of functional extracellular vesicles by endothelial cells might be interesting regarding their important role in coronary artery diseases.¹⁴² Finally, the impact of new-generation TKIs on human blood coagulation and fibrinolysis has never been studied and should be addressed.

To conclude, new-generation TKIs increase the risk of arterial thrombosis in patients with CML, whereas imatinib, the first-generation TKI, might prevent the development of cardiovascular events. To date, the cellular events and signaling pathways by which these events occurred are unknown and researches are extremely limited focusing mainly on imatinib and nilotinib. Researches need to be extended to all new-generation BCR-ABL TKIs (i.e., dasatinib, bosutinib, and ponatinib). The understanding of the mechanisms by which new-generation BCR-ABL TKIs induce or promote arterial occlusive events will improve the clinical uses of these therapies. To date, only general risk minimization measures have been proposed (e.g., management of dyslipidemia,

^{aiv}DDR-1: Discoidin domain receptor tyrosine kinase 1. DDR1 is involved in the regulation of cell growth, differentiation, and metabolism.

^{av}Tie-2: Tunica interna endothelial cell kinase. Tie-2 regulates angiogenesis, endothelial cell survival, proliferation, migration, adhesion and cell spreading, cytoskeleton reorganization, and vascular quiescence. Tie-2 also possesses anti-inflammatory functions by preventing the leakage of proinflammatory mediators and leukocytes.

diabetes, arterial hypertension following standard of care).^{14,22,23,143–146} The understanding of the pathophysiology is required to implement the most appropriate risk minimization strategies for thrombotic events and to select patients to whom the prescription of these drugs should be avoided when applicable. Finally, the understanding of the pathophysiology will help in the design of new BCR-ABL inhibitors sparing the toxic targets.

Review Criteria

Relevant articles published from the database inception to July 11, 2017, were identified from an electronic database (PubMed) using the keywords “vascular,” “thrombosis,” “atherosclerosis,” “arteriosclerosis,” “venous,” “arterial,” “hemostasis,” “metabolic,” “metabolism,” “glycemia,” “glycaemia,” “cholesterol,” “triglycerides,” and “platelet” combined with the five approved BCR-ABL TKIs. The search strategy is presented in supplementary files. Articles published in languages other than English were excluded from the analysis. Primary criteria were pathophysiological explanation of arterial thrombotic events. Abstracts and full-text articles were reviewed with a focus on atherogenesis, plaque rupture, platelet functions, and their link with the development of arterial thrombosis with BCR-ABL TKIs. The reference section of identified articles was also examined.

Authors' Contributions

H.H. was responsible for the first draft of the manuscript. F.M., C.C., C.G., J.M.D., and J.D. contributed to the final draft of the manuscript.

Conflicts of Interest

J.D. reports personal fees from Roche Diagnostics, Stago Diagnostica, Bayer Healthcare, and Daiichi-Sankyo; travel grants from Bayer Healthcare, Boehringer Ingelheim, CSL Behring, and Stago Diagnostica outside the submitted work.

F.M. reports personal fees from Boehringer Ingelheim, Bayer Healthcare, and Bristol-Myers Squibb-Pfizer outside the submitted work.

C.G. reports personal fees from Novartis, Celgene, and Amgen outside the submitted work.

The other authors have no conflicts of interest to disclose.

References

- Bhamidipati PK, Kantarjian H, Cortes J, Cornelison AM, Jabbour E. Management of imatinib-resistant patients with chronic myeloid leukemia. *Ther Adv Hematol* 2013;4(02):103–117
- Gorre ME, Mohammed M, Ellwood K, et al. Clinical resistance to STI-571 cancer therapy caused by BCR-ABL gene mutation or amplification. *Science* 2001;293(5531):876–880
- Bixby D, Talpaz M. Mechanisms of resistance to tyrosine kinase inhibitors in chronic myeloid leukemia and recent therapeutic strategies to overcome resistance. *Hematology (Am Soc Hematol Educ Program)* 2009;2009(01):461–476
- Cortes JE, Saglio G, Kantarjian HM, et al. Final 5-year study results of DASISION: the Dasatinib Versus Imatinib Study in Treatment-Naïve Chronic Myeloid Leukemia Patients Trial. *J Clin Oncol* 2016;34(20):2333–2340
- Hochhaus A, Saglio G, Hughes TP, et al. Long-term benefits and risks of frontline nilotinib vs imatinib for chronic myeloid leukemia in chronic phase: 5-year update of the randomized ENESTnd trial. *Leukemia* 2016;30(05):1044–1054
- Brümmendorf TH, Cortes JE, de Souza CA, et al. Bosutinib versus imatinib in newly diagnosed chronic-phase chronic myeloid leukaemia: results from the 24-month follow-up of the BELA trial. *Br J Haematol* 2015;168(01):69–81
- Mealing S, Barcena L, Hawkins N, et al. The relative efficacy of imatinib, dasatinib and nilotinib for newly diagnosed chronic myeloid leukemia: a systematic review and network meta-analysis. *Exp Hematol Oncol* 2013;2(01):5
- Lamontanara AJ, Gencer EB, Kuzyk O, Hantschel O. Mechanisms of resistance to BCR-ABL and other kinase inhibitors. *Biochim Biophys Acta* 2013;1834(07):1449–1459
- Pagnano KB, Bendit I, Boquimpani C, et al; Latin American Leukemia Net (LALNET). BCR-ABL mutations in chronic myeloid leukemia treated with tyrosine kinase inhibitors and impact on survival. *Cancer Invest* 2015;33(09):451–458
- Ursan ID, Jiang R, Pickard EM, Lee TA, Ng D, Pickard AS. Emergence of BCR-ABL kinase domain mutations associated with newly diagnosed chronic myeloid leukemia: a meta-analysis of clinical trials of tyrosine kinase inhibitors. *J Manag Care Spec Pharm* 2015;21(02):114–122
- National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Chronic Myeloid Leukemia. Version 1.2017. 2016
- Thanopoulou E, Judson I. The safety profile of imatinib in CML and GIST: long-term considerations. *Arch Toxicol* 2012;86(01):1–12
- Kalmanti L, Saussele S, Laussek M, et al. Safety and efficacy of imatinib in CML over a period of 10 years: data from the randomized CML-study IV. *Leukemia* 2015;29(05):1123–1132
- Moslehi JJ, Deininger M. Tyrosine kinase inhibitor-associated cardiovascular toxicity in chronic myeloid leukemia. *J Clin Oncol* 2015;33(35):4210–4218
- Breccia M, Alimena G. Pleural/pericardic effusions during dasatinib treatment: incidence, management and risk factors associated to their development. *Expert Opin Drug Saf* 2010;9(05):713–721
- European Medicines Agency. Sprycel - Summary of Product Characteristics 2017. Available at: http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/000709/WC500056998.pdf. Accessed December 22, 2017
- European Medicines Agency. Iclusig - Summary of Product Characteristics 2017. Available at: http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/002695/WC500145646.pdf. Accessed December 22, 2017
- European Medicines Agency. Tasigna - Summary of Product Characteristics 2017. Available at: http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/000798/WC500034394.pdf. Accessed December 22, 2017
- Kantarjian HM, Cortes JE, Kim DW, et al. Bosutinib safety and management of toxicity in leukemia patients with resistance or intolerance to imatinib and other tyrosine kinase inhibitors. *Blood* 2014;123(09):1309–1318
- Saglio G, Kim D-W, Issaragrisil S, et al; ENESTnd Investigators. Nilotinib versus imatinib for newly diagnosed chronic myeloid leukemia. *N Engl J Med* 2010;362(24):2251–2259
- Poch Martell M, Sibai H, Deotare U, Lipton JH. Ponatinib in the therapy of chronic myeloid leukemia. *Expert Rev Hematol* 2016;9(10):923–932
- Douxflis J, Haguët H, Mullier F, Chatelain C, Graux C, Dogné JM. Association between BCR-ABL tyrosine kinase inhibitors for chronic myeloid leukemia and cardiovascular events, major molecular response, and overall survival: a systematic review and meta-analysis. *JAMA Oncol* 2016; doi: 10.1001/jamaoncol.2015.5932

- 23 Haguet H, Douxfils J, Mullier F, Chatelain C, Graux C, Dogné JM. Risk of arterial and venous occlusive events in chronic myeloid leukemia patients treated with new generation BCR-ABL tyrosine kinase inhibitors: a systematic review and meta-analysis. *Expert Opin Drug Saf* 2017;16(01):5–12
- 24 Cortes J, Kim D-W, Pinilla-Ibarz J, et al. Ponatinib efficacy and safety in heavily pretreated leukemia patients: 3-year results of the PACE trial. 20th Congress of EHA; Copenhagen, Denmark, June 12, 2015
- 25 Pasvolosky O, Leader A, Iakobishvili Z, Wasserstrum Y, Kornowski R, Raanani P. Tyrosine kinase inhibitor associated vascular toxicity in chronic myeloid leukemia. *Cardio-Oncology* 2015;1(01):5
- 26 Gora-Tybor J, Medras E, Calbecka M, et al. Real-life comparison of severe vascular events and other non-hematological complications in patients with chronic myeloid leukemia undergoing second-line nilotinib or dasatinib treatment. *Leuk Lymphoma* 2015;56(08):2309–2314
- 27 le Coutre PD, Hughes TP, Mahon FX, et al. Low incidence of peripheral arterial disease in patients receiving dasatinib in clinical trials. *Leukemia* 2016;30(07):1593–1596
- 28 Sam PY, Ahaneku H, Nogueras-Gonzalez GM, et al. Cardiovascular events among patients with chronic myeloid leukemia (CML) treated with tyrosine kinase inhibitors (TKIs). *Blood* 2016;128(22):1919
- 29 Lassila M, Allen TJ, Cao Z, et al. Imatinib attenuates diabetes-associated atherosclerosis. *Arterioscler Thromb Vasc Biol* 2004;24(05):935–942
- 30 Ma LK, Li Q, He LF, et al. Imatinib attenuates myocardial fibrosis in association with inhibition of the PDGFR α activity. *Arq Bras Cardiol* 2012;99(06):1082–1091
- 31 Fossard G, Blond E, Balsat M, et al. Hyperhomocysteinemia and high doses of nilotinib favor cardiovascular events in chronic phase chronic myelogenous leukemia patients. *Haematologica* 2016;101(03):e86–e90
- 32 Kostos L, Burbury K, Srivastava G, Prince HM. Gastrointestinal bleeding in a chronic myeloid leukaemia patient precipitated by dasatinib-induced platelet dysfunction: case report. *Platelets* 2015;26(08):809–811
- 33 Mazharian A, Ghevaert C, Zhang L, Massberg S, Watson SP. Dasatinib enhances megakaryocyte differentiation but inhibits platelet formation. *Blood* 2011;117(19):5198–5206
- 34 Alhawiti N, Burbury KL, Kwa FA, et al. The tyrosine kinase inhibitor, nilotinib potentiates a prothrombotic state. *Thromb Res* 2016;145:54–64
- 35 Lotfi K, Deb S, Sjöström C, Tharmakulanathan A, Boknäs N, Ramström S. Individual variation in hemostatic alterations caused by tyrosine kinase inhibitors: a way to improve personalized cancer therapy? *Blood* 2016;128(22):1908
- 36 Li R, Grosser T, Diamond SL. Microfluidic whole blood testing of platelet response to pharmacological agents. *Platelets* 2017;28(05):457–462
- 37 Senis YA, Mazharian A, Mori J. Src family kinases: at the forefront of platelet activation. *Blood* 2014;124(13):2013–2024
- 38 Gratacap MP, Martin V, Valéra MC, et al. The new tyrosine-kinase inhibitor and anticancer drug dasatinib reversibly affects platelet activation in vitro and in vivo. *Blood* 2009;114(09):1884–1892
- 39 Albrecht-Schgoer K, Huber K, Grebien F, et al. Nilotinib exerts direct pro-atherogenic and anti-angiogenic effects on vascular endothelial cells: a potential explanation for drug-induced vasculopathy in CML. *Blood* 2013;122(21):257
- 40 Shimabukuro-Vornhagen A, Rothe A, Nogova L, Kochanek M, Scheid C, von Bergwelt-Baildon M. Improvement of platelet dysfunction in chronic myelogenous leukemia following treatment with imatinib: a case report. *J Med Case Reports* 2011;5:215
- 41 Neelakantan P, Marin D, Laffan M, Goldman J, Apperley J, Milojkovic D. Platelet dysfunction associated with ponatinib, a new pan BCR-ABL inhibitor with efficacy for chronic myeloid leukemia resistant to multiple tyrosine kinase inhibitor therapy. *Haematologica* 2012;97(09):1444
- 42 Loren CP, Aslan JE, Rigg RA, et al. The BCR-ABL inhibitor ponatinib inhibits platelet immunoreceptor tyrosine-based activation motif (ITAM) signaling, platelet activation and aggregate formation under shear. *Thromb Res* 2015;135(01):155–160
- 43 Rivera VM, Pritchard JR, Gonzalez F, Baker T, Gozgit JM, Hodgson G. Comparative TKI profiling analyses to explore potential mechanisms of Ponatinib-associated arterial thrombotic events. *Blood* 2014;124(21):1783
- 44 Quintás-Cardama A, Han X, Kantarjian H, Cortes J. Tyrosine kinase inhibitor-induced platelet dysfunction in patients with chronic myeloid leukemia. *Blood* 2009;114(02):261–263
- 45 Breccia M, Muscaritoli M, Aversa Z, Mandelli F, Alimena G. Imatinib mesylate may improve fasting blood glucose in diabetic Ph+ chronic myelogenous leukemia patients responsive to treatment. *J Clin Oncol* 2004;22(22):4653–4655
- 46 Breccia M, Alimena G. The metabolic consequences of imatinib mesylate: changes on glucose, lipidic and bone metabolism. *Leuk Res* 2009;33(07):871–875
- 47 Agostino NM, Chinchilli VM, Lynch CJ, et al. Effect of the tyrosine kinase inhibitors (sunitinib, sorafenib, dasatinib, and imatinib) on blood glucose levels in diabetic and nondiabetic patients in general clinical practice. *J Oncol Pharm Pract* 2011;17(03):197–202
- 48 Breccia M, Muscaritoli M, Cannella L, Stefanizzi C, Frustaci A, Alimena G. Fasting glucose improvement under dasatinib treatment in an accelerated phase chronic myeloid leukemia patient unresponsive to imatinib and nilotinib. *Leuk Res* 2008;32(10):1626–1628
- 49 Ono K, Suzushima H, Watanabe Y, et al. Rapid amelioration of hyperglycemia facilitated by dasatinib in a chronic myeloid leukemia patient with type 2 diabetes mellitus. *Intern Med* 2012;51(19):2763–2766
- 50 Louvet C, Szot GL, Lang J, et al. Tyrosine kinase inhibitors reverse type 1 diabetes in nonobese diabetic mice. *Proc Natl Acad Sci U S A* 2008;105(48):18895–18900
- 51 Hägerkvist R, Jansson L, Welsh N. Imatinib mesylate improves insulin sensitivity and glucose disposal rates in rats fed a high-fat diet. *Clin Sci (Lond)* 2008;114(01):65–71
- 52 Han MS, Chung KW, Cheon HG, et al. Imatinib mesylate reduces endoplasmic reticulum stress and induces remission of diabetes in db/db mice. *Diabetes* 2009;58(02):329–336
- 53 Hägerkvist R, Makeeva N, Elliman S, Welsh N. Imatinib mesylate (Gleevec) protects against streptozotocin-induced diabetes and islet cell death in vitro. *Cell Biol Int* 2006;30(12):1013–1017
- 54 Hägerkvist R, Sandler S, Mokhtari D, Welsh N. Amelioration of diabetes by imatinib mesylate (Gleevec): role of beta-cell NF-kappaB activation and anti-apoptotic preconditioning. *FASEB J* 2007;21(02):618–628
- 55 Xia CQ, Zhang P, Li S, et al. C-Abl inhibitor imatinib enhances insulin production by β cells: c-Abl negatively regulates insulin production via interfering with the expression of NKx2.2 and GLUT-2. *PLoS One* 2014;9(05):e97694
- 56 Sadiq S, Owen E, Foster T, et al. Nilotinib-induced metabolic dysfunction: insights from a translational pilot study using in vitro adipocyte models and patient cohorts. 22nd EHA congress; June 22–25, 2017; Madrid, Spain
- 57 Iizuka K, Niwa H, Kato T, Takeda J. Dasatinib improves insulin sensitivity and affects lipid metabolism in a patient with chronic myeloid leukaemia. *BMJ Case Rep* 2016;2016:201
- 58 Fitter S, Vandyke K, Schultz CG, White D, Hughes TP, Zannettino AC. Plasma adiponectin levels are markedly elevated in imatinib-treated chronic myeloid leukemia (CML) patients: a mechanism for improved insulin sensitivity in type 2 diabetic CML patients? *J Clin Endocrinol Metab* 2010;95(08):3763–3767
- 59 Hosch SE, Olefsky JM, Kim JJ. APPLIED mechanics: uncovering how adiponectin modulates insulin action. *Cell Metab* 2006;4(01):5–6

- 60 Lihn AS, Pedersen SB, Richelsen B. Adiponectin: action, regulation and association to insulin sensitivity. *Obes Rev* 2005;6(01):13–21
- 61 Fitter S, Vandyke K, Gronthos S, Zannettino AC. Suppression of PDGF-induced PI3 kinase activity by imatinib promotes adipogenesis and adiponectin secretion. *J Mol Endocrinol* 2012;48(03):229–240
- 62 Aichberger KJ, Herndlhofer S, Schernthaner GH, et al. Progressive peripheral arterial occlusive disease and other vascular events during nilotinib therapy in CML. *Am J Hematol* 2011;86(07):533–539
- 63 Breccia M, Muscaritoli M, Gentilini F, et al. Impaired fasting glucose level as metabolic side effect of nilotinib in non-diabetic chronic myeloid leukemia patients resistant to imatinib. *Leuk Res* 2007;31(12):1770–1772
- 64 Castagnetti F, Breccia M, Gugliotta G, et al. GIMEMA CML Working Party. Nilotinib 300 mg twice daily: an academic single-arm study of newly diagnosed chronic phase chronic myeloid leukemia patients. *Haematologica* 2016;101(10):1200–1207
- 65 Delphine R, Gautier J-f, Breccia M, et al. Incidence of hyperglycemia by 3 years in patients (Pts) with newly diagnosed chronic myeloid leukemia in chronic phase (CML-CP) treated with Nilotinib (NIL) or Imatinib (IM) in ENESTnd. *Blood* 2012;120(21):1686
- 66 Iurlo A, Orsi E, Cattaneo D, et al. Effects of first- and second-generation tyrosine kinase inhibitor therapy on glucose and lipid metabolism in chronic myeloid leukemia patients: a real clinical problem? *Oncotarget* 2015;6(32):33944–33951
- 67 Racil Z, Razga F, Drapalova J, et al. Mechanism of impaired glucose metabolism during nilotinib therapy in patients with chronic myelogenous leukemia. *Haematologica* 2013;98(10):e124–e126
- 68 Ito Y, Miyamoto T, Chong Y, Maki T, Akashi K, Kamimura T. Nilotinib exacerbates diabetes mellitus by decreasing secretion of endogenous insulin. *Int J Hematol* 2013;97(01):135–138
- 69 Hadzijušufovic E, Herndlhofer S, Aichberger KJ, et al. Nilotinib exerts direct effects on vascular endothelial cells and may act as a co-trigger of atherosclerosis in patients with Ph+ CML. *Blood* 2011;118(21):2753
- 70 Breccia M, Loggisci G, Salaroli A, Serrao A, Alimena G. Nilotinib-mediated increase in fasting glucose level is reversible, does not convert to type 2 diabetes and is likely correlated with increased body mass index. *Leuk Res* 2012;36(04):e66–e67
- 71 Saglio G, Larson RA, Hughes TP, et al. Efficacy and safety of Nilotinib in chronic phase (CP) chronic myeloid leukemia (CML) patients (Pts) with type 2 diabetes in the ENESTnd trial. *Blood* 2010;116(21):3430
- 72 Frasca F, Pandini G, Malaguarnera R, et al. Role of c-Abl in directing metabolic versus mitogenic effects in insulin receptor signaling. *J Biol Chem* 2007;282(36):26077–26088
- 73 Franceschino A, Tornaghi L, Benemacher V, Assouline S, Gambacorti-Passerini C. Alterations in creatine kinase, phosphate and lipid values in patients with chronic myeloid leukemia during treatment with imatinib. *Haematologica* 2008;93(02):317–318
- 74 Gottardi M, Manzato E, Gherlinzoni F. Imatinib and hyperlipidemia. *N Engl J Med* 2005;353(25):2722–2723
- 75 Gologan R, Constantinescu G, Georgescu D, et al. Hypolipemiant besides antileukemic effect of imatinib mesylate. *Leuk Res* 2009;33(09):1285–1287
- 76 Ellis M, Krashin E, Hamburger-Avnery O, Gan S, Elis A, Ashur-Fabian O. The anti-leukemic and lipid lowering effects of imatinib are not hindered by statins in CML: a retrospective clinical study and in vitro assessment of lipid-genes transcription. *Leuk Lymphoma* 2017;58(05):1172–1177
- 77 Franklin M, Burns L, Perez S, Yerragolam D, Makenbaeva D. Incidence of type II diabetes mellitus and hyperlipidemia in patients prescribed Dasatinib or Nilotinib as first or second line therapy for chronic myelogenous leukemia (CML). *Blood* 2016;128(22):4766
- 78 Rea D, Mirault T, Cluzeau T, et al. Early onset hypercholesterolemia induced by the 2nd-generation tyrosine kinase inhibitor nilotinib in patients with chronic phase-chronic myeloid leukemia. *Haematologica* 2014;99(07):1197–1203
- 79 Maurizot A, Beressi JP, Manéglier B, et al. Rapid clinical improvement of peripheral artery occlusive disease symptoms after nilotinib discontinuation despite persisting vascular occlusion. *Blood Cancer J* 2014;4:e247
- 80 Breccia M, Molica M, Alimena G. How tyrosine kinase inhibitors impair metabolism and endocrine system function: a systematic updated review. *Leuk Res* 2014;38(12):1392–1398
- 81 Venalis P, Maurer B, Akhmetshina A, et al. Lack of inhibitory effects of the anti-fibrotic drug imatinib on endothelial cell functions in vitro and in vivo. *J Cell Mol Med* 2009;13(10):4185–4191
- 82 Gover-Proaktor A, Granot G, Shapira S, et al. Ponatinib reduces viability, migration, and functionality of human endothelial cells. *Leuk Lymphoma* 2017;58(06):1455–1467
- 83 Vallières K, Petitclerc E, Laroche G. On the ability of imatinib mesylate to inhibit smooth muscle cell proliferation without delaying endothelialization: an in vitro study. *Vascul Pharmacol* 2009;51(01):50–56
- 84 Hacker TA, Griffin MO, Guttormsen B, Stoker S, Wolff MR. Platelet-derived growth factor receptor antagonist STI571 (imatinib mesylate) inhibits human vascular smooth muscle proliferation and migration in vitro but not in vivo. *J Invasive Cardiol* 2007;19(06):269–274
- 85 Vrekoussis T, Stathopoulos EN, De Giorgi U, et al. Modulation of vascular endothelium by imatinib: a study on the EA.hy 926 endothelial cell line. *J Chemother* 2006;18(01):56–65
- 86 Letsiou E, Rizzo AN, Sammani S, et al. Differential and opposing effects of imatinib on LPS- and ventilator-induced lung injury. *Am J Physiol Lung Cell Mol Physiol* 2015;308(03):L259–L269
- 87 Guignabert C, Phan C, Seferian A, et al. Dasatinib induces lung vascular toxicity and predisposes to pulmonary hypertension. *J Clin Invest* 2016;126(09):3207–3218
- 88 Rizzo AN, Sammani S, Esquinca AE, et al. Imatinib attenuates inflammation and vascular leak in a clinically relevant two-hit model of acute lung injury. *Am J Physiol Lung Cell Mol Physiol* 2015;309(11):L1294–L1304
- 89 Katgi A, Sevindik ÖG, Gökbülüt AA, et al. Nilotinib does not alter the secretory functions of carotid artery endothelial cells in a prothrombotic or antithrombotic fashion. *Clin Appl Thromb Hemost* 2015;21(07):678–683
- 90 Hadzijušufovic E, Kirchmair R, Theurl M, et al. Ponatinib exerts multiple effects on vascular endothelial cells: possible mechanisms and explanations for the adverse vascular events seen in CML patients treated with Ponatinib. *Blood* 2016;128(22):1883
- 91 Lorenzi M, Cagliero E, Toledo S. Glucose toxicity for human endothelial cells in culture. Delayed replication, disturbed cell cycle, and accelerated death. *Diabetes* 1985;34(07):621–627
- 92 Shah NP, Wallis N, Farber HW, et al. Clinical features of pulmonary arterial hypertension in patients receiving dasatinib. *Am J Hematol* 2015;90(11):1060–1064
- 93 Frost AE, Barst RJ, Hooper MM, et al. Long-term safety and efficacy of imatinib in pulmonary arterial hypertension. *J Heart Lung Transplant* 2015;34(11):1366–1375
- 94 Dasgupta SK, Le A, Vijayan KV, Thiagarajan P. Dasatinib inhibits actin fiber reorganization and promotes endothelial cell permeability through RhoA-ROCK pathway. *Cancer Med* 2017;6(04):809–818
- 95 Dewar AL, Domaschew RM, Doherty KV, Hughes TP, Lyons AB. Imatinib inhibits the in vitro development of the monocyte/macrophage lineage from normal human bone marrow progenitors. *Leukemia* 2003;17(09):1713–1721

- 96 Brownlow N, Mol C, Hayford C, Ghaem-Maghami S, Dibb NJ. Dasatinib is a potent inhibitor of tumour-associated macrophages, osteoclasts and the FMS receptor. *Leukemia* 2009;23(03):590–594
- 97 Dewar AL, Doherty KV, Hughes TP, Lyons AB. Imatinib inhibits the functional capacity of cultured human monocytes. *Immunol Cell Biol* 2005;83(01):48–56
- 98 Dewar AL, Cambareri AC, Zannettino ACW, et al. Macrophage colony-stimulating factor receptor c-fms is a novel target of imatinib. *Blood* 2005;105(08):3127–3132
- 99 Metharom P, Martin K, Kumar AH, et al. Pleiotropic role for monocyte C-fms protein in response to vascular injury: potential therapeutic target. *Atherosclerosis* 2011;216(01):74–82
- 100 Gacic J, Vorkapic E, Olsen RS, et al. Imatinib reduces cholesterol uptake and matrix metalloproteinase activity in human THP-1 macrophages. *Pharmacol Rep* 2016;68(01):1–6
- 101 Cwynarski K, Laylor R, Macchiarulo E, et al. Imatinib inhibits the activation and proliferation of normal T lymphocytes in vitro. *Leukemia* 2004;18(08):1332–1339
- 102 Brownlow N, Russell AE, Saravanapavan H, et al. Comparison of nilotinib and imatinib inhibition of FMS receptor signaling, macrophage production and osteoclastogenesis. *Leukemia* 2008;22(03):649–652
- 103 Ozanne J, Prescott AR, Clark K. The clinically approved drugs dasatinib and bosutinib induce anti-inflammatory macrophages by inhibiting the salt-inducible kinases. *Biochem J* 2015;465(02):271–279
- 104 Darling NJ, Toth R, Arthur JS, Clark K. Inhibition of SIK2 and SIK3 during differentiation enhances the anti-inflammatory phenotype of macrophages. *Biochem J* 2017;474(04):521–537
- 105 Schade AE, Schieven GL, Townsend R, et al. Dasatinib, a small-molecule protein tyrosine kinase inhibitor, inhibits T-cell activation and proliferation. *Blood* 2008;111(03):1366–1377
- 106 Blake SJ, Lyons AB, Hughes TP. Nilotinib inhibits the Src-family kinase LCK and T-cell function in vitro. *J Cell Mol Med* 2009;13(03):599–601
- 107 Blake S, Hughes TP, Mayrhofer G, Lyons AB. The Src/ABL kinase inhibitor dasatinib (BMS-354825) inhibits function of normal human T-lymphocytes in vitro. *Clin Immunol* 2008;127(03):330–339
- 108 Lee KC, Ouwehand I, Giannini AL, Thomas NS, Dibb NJ, Bijlmakers MJ. Lck is a key target of imatinib and dasatinib in T-cell activation. *Leukemia* 2010;24(04):896–900
- 109 Farha S, Dweik R, Rahaghi F, et al. Imatinib in pulmonary arterial hypertension: c-Kit inhibition. *Pulm Circ* 2014;4(03):452–455
- 110 Juurikivi A, Sandler C, Lindstedt KA, et al. Inhibition of c-kit tyrosine kinase by imatinib mesylate induces apoptosis in mast cells in rheumatoid synovia: a potential approach to the treatment of arthritis. *Ann Rheum Dis* 2005;64(08):1126–1131
- 111 Sun YL, Kumar P, Sodani K, et al. Ponatinib enhances anticancer drug sensitivity in MRP7-overexpressing cells. *Oncol Rep* 2014;31(04):1605–1612
- 112 Yvan-Charvet L, Wang N, Tall AR. Role of HDL, ABCA1, and ABCG1 transporters in cholesterol efflux and immune responses. *Arterioscler Thromb Vasc Biol* 2010;30(02):139–143
- 113 Chen Z, Lee FY, Bhalla KN, Wu J. Potent inhibition of platelet-derived growth factor-induced responses in vascular smooth muscle cells by BMS-354825 (dasatinib). *Mol Pharmacol* 2006;69(05):1527–1533
- 114 Nakamura K, Akagi S, Ogawa A, et al. Pro-apoptotic effects of imatinib on PDGF-stimulated pulmonary artery smooth muscle cells from patients with idiopathic pulmonary arterial hypertension. *Int J Cardiol* 2012;159(02):100–106
- 115 Li L, Blumenthal DK, Masaki T, Terry CM, Cheung AK. Differential effects of imatinib on PDGF-induced proliferation and PDGF receptor signaling in human arterial and venous smooth muscle cells. *J Cell Biochem* 2006;99(06):1553–1563
- 116 Rocha A, Azevedo I, Soares R. Anti-angiogenic effects of imatinib target smooth muscle cells but not endothelial cells. *Angiogenesis* 2007;10(04):279–286
- 117 Ballinger ML, Osman N, Hashimura K, et al. Imatinib inhibits vascular smooth muscle proteoglycan synthesis and reduces LDL binding in vitro and aortic lipid deposition in vivo. *J Cell Mol Med* 2010;14(6B):1408–1418
- 118 Distler JH, Jüngel A, Huber LC, et al. Imatinib mesylate reduces production of extracellular matrix and prevents development of experimental dermal fibrosis. *Arthritis Rheum* 2007;56(01):311–322
- 119 Yoshiji H, Noguchi R, Kuriyama S, et al. Imatinib mesylate (STI-571) attenuates liver fibrosis development in rats. *Am J Physiol Gastrointest Liver Physiol* 2005;288(05):G907–G913
- 120 Salhotra A, Tsai N, Thomas SH, et al. Sclerodermitous chronic GVHD in patients receiving tyrosine kinase inhibitors after allogeneic hematopoietic cell transplantation. *Bone Marrow Transplant* 2015;50(01):139–141
- 121 Makiyama Y, Toba K, Kato K, et al. Imatinib mesilate inhibits neointimal hyperplasia via growth inhibition of vascular smooth muscle cells in a rat model of balloon injury. *Tohoku J Exp Med* 2008;215(04):299–306
- 122 Park D, Kim SM, Min SI, Ha J, Kim IG, Min SK. Inhibition of intimal hyperplasia by local perivascular application of rapamycin and imatinib mesilate after carotid balloon injury. *J Korean Surg Soc* 2013;85(06):296–301
- 123 Leppänen O, Rutanen J, Hiltunen MO, et al. Oral imatinib mesylate (STI571/Gleevec) improves the efficacy of local intra-vascular endothelial growth factor-C gene transfer in reducing neointimal growth in hypercholesterolemic rabbits. *Circulation* 2004;109(09):1140–1146
- 124 Sevim KZ, Silistreli O, Gorgu M, Sevim O, Ergur B. Short-term vasculoprotective effects of imatinib mesylate on intimal hyperplasia of arterial anastomosis: An experimental study using a rabbit model. *Can J Plast Surg* 2012;20(04):223–228
- 125 Wu J, Chen Z, Bhalla KN. Treatment of restenosis and stenosis with dasatinib (2014). USF Patents. 152
- 126 Sánchez-Ortega I, Servitje O, Arnan M, et al. Dasatinib as salvage therapy for steroid refractory and imatinib resistant or intolerant sclerotic chronic graft-versus-host disease. *Biol Blood Marrow Transplant* 2012;18(02):318–323
- 127 Marinelli Busilacchi E, Olivieri J, Viola N, et al. The exaggerated collagen expression in Gvhd-fibroblasts is effectively inhibited by therapeutic concentration of Nilotinib. *Blood* 2016;128(22):4597
- 128 Shiha GE, Abu-Elsaad NM, Zalata KR, Ibrahim TM. Tracking anti-fibrotic pathways of nilotinib and imatinib in experimentally induced liver fibrosis: an insight. *Clin Exp Pharmacol Physiol* 2014;41(10):788–797
- 129 Giles FJ, O'Dwyer M, Swords R. Class effects of tyrosine kinase inhibitors in the treatment of chronic myeloid leukemia. *Leukemia* 2009;23(10):1698–1707
- 130 Rix U, Hantschel O, Dürnberger G, et al. Chemical proteomic profiles of the BCR-ABL inhibitors imatinib, nilotinib, and dasatinib reveal novel kinase and nonkinase targets. *Blood* 2007;110(12):4055–4063
- 131 Pardanani A, Tefferi A. Imatinib targets other than bcr/abl and their clinical relevance in myeloid disorders. *Blood* 2004;104(07):1931–1939
- 132 Bantscheff M, Eberhard D, Abraham Y, et al. Quantitative chemical proteomics reveals mechanisms of action of clinical ABL kinase inhibitors. *Nat Biotechnol* 2007;25(09):1035–1044
- 133 Weisberg E, Manley PW, Breitenstein W, et al. Characterization of AMN107, a selective inhibitor of native and mutant Bcr-Abl. *Cancer Cell* 2005;7(02):129–141
- 134 Buchdunger E, Cioffi CL, Law N, et al. Abl protein-tyrosine kinase inhibitor STI571 inhibits in vitro signal transduction mediated

- by c-kit and platelet-derived growth factor receptors. *J Pharmacol Exp Ther* 2000;295(01):139–145
- 135 Manley PW, Druce P, Fendrich G, et al. Extended kinase profile and properties of the protein kinase inhibitor nilotinib. *Biochim Biophys Acta* 2010;1804(03):445–453
 - 136 Fabian MA, Biggs WH III, Treiber DK, et al. A small molecule-kinase interaction map for clinical kinase inhibitors. *Nat Biotechnol* 2005;23(03):329–336
 - 137 Shi H, Zhang CJ, Chen GY, Yao SQ. Cell-based proteome profiling of potential dasatinib targets by use of affinity-based probes. *J Am Chem Soc* 2012;134(06):3001–3014
 - 138 Liu Y, Wang Z, Kwong SQ, et al. Inhibition of PDGF, TGF- β , and Abl signaling and reduction of liver fibrosis by the small molecule Bcr-Abl tyrosine kinase antagonist Nilotinib. *J Hepatol* 2011;55(03):612–625
 - 139 Sakamoto A, Sugamoto Y, Tokunaga Y, et al. Expression profiling of the ephrin (EFN) and Eph receptor (EPH) family of genes in atherosclerosis-related human cells. *J Int Med Res* 2011;39(02):522–527
 - 140 Khan AA, Sandhya VK, Singh P, et al. Signaling network map of endothelial TEK tyrosine kinase. *J Signal Transduct* 2014;2014:173026
 - 141 Chislock EM, Ring C, Pendergast AM. Abl kinases are required for vascular function, Tie2 expression, and angiopoietin-1-mediated survival. *Proc Natl Acad Sci U S A* 2013;110(30):12432–12437
 - 142 Boulanger CM, Loyer X, Rautou PE, Amabile N. Extracellular vesicles in coronary artery disease. *Nat Rev Cardiol* 2017;14(05):259–272
 - 143 Aghel N, Delgado DH, Lipton JH. Cardiovascular toxicities of BCR-ABL tyrosine kinase inhibitors in chronic myeloid leukemia: preventive strategies and cardiovascular surveillance. *Vasc Health Risk Manag* 2017;13:293–303
 - 144 Damrongwatanasuk R, Fradley MG. Cardiovascular complications of targeted therapies for chronic myeloid leukemia. *Curr Treat Options Cardiovasc Med* 2017;19(04):24
 - 145 Valent P, Hadzijusufovic E, Hoermann G, et al. Risk factors and mechanisms contributing to TKI-induced vascular events in patients with CML. *Leuk Res* 2017;59:47–54
 - 146 García-Gutiérrez V, Jiménez-Velasco A, Gómez-Casares MT, Sánchez-Guijo F, López-Sendón JL, Steegmann Olmedillas JL. [Cardiovascular management of patients with chronic myeloid leukemia from a multidisciplinary perspective, and proposing action protocol by consensus meeting]. *Med Clin (Barc)* 2016;146(12):561.e1–561.e8
 - 147 Aman J, van Bezuij J, Damanafshan A, et al. Effective treatment of edema and endothelial barrier dysfunction with imatinib. *Circulation* 2012;126(23):2728–2738
 - 148 Kim SR, Suh W. Beneficial effects of the Src inhibitor, dasatinib, on breakdown of the blood-retinal barrier. *Arch Pharm Res* 2017;40(02):197–203
 - 149 Sukegawa M, Wang X, Nishioka C, et al. The BCR/ABL tyrosine kinase inhibitor, nilotinib, stimulates expression of IL-1 β in vascular endothelium in association with downregulation of miR-3p. *Leuk Res* 2017;58:83–90
 - 150 Wolf AM, Wolf D, Rumpold H, et al. The kinase inhibitor imatinib mesylate inhibits TNF-alpha production in vitro and prevents TNF-dependent acute hepatic inflammation. *Proc Natl Acad Sci U S A* 2005;102(38):13622–13627
 - 151 Fraser CK, Lousberg EL, Kumar R, Hughes TP, Diener KR, Hayball JD. Dasatinib inhibits the secretion of TNF-alpha following TLR stimulation in vitro and in vivo. *Exp Hematol* 2009;37(12):1435–1444
 - 152 Seggewiss R, Loré K, Greiner E, et al. Imatinib inhibits T-cell receptor-mediated T-cell proliferation and activation in a dose-dependent manner. *Blood* 2005;105(06):2473–2479
 - 153 Dietz AB, Souan L, Knutson GJ, Bulur PA, Litzow MR, Vuk-Pavlovic S. Imatinib mesylate inhibits T-cell proliferation in vitro and delayed-type hypersensitivity in vivo. *Blood* 2004;104(04):1094–1099
 - 154 Chen J, Schmitt A, Chen B, et al. Nilotinib hampers the proliferation and function of CD8+ T lymphocytes through inhibition of T cell receptor signalling. *J Cell Mol Med* 2008;12(5B):2107–2118
 - 155 Perros F, Montani D, Dorfmueller P, et al. Platelet-derived growth factor expression and function in idiopathic pulmonary arterial hypertension. *Am J Respir Crit Care Med* 2008;178(01):81–88
 - 156 Canning P, Tan L, Chu K, Lee SW, Gray NS, Bullock AN. Structural mechanisms determining inhibition of the collagen receptor DDR1 by selective and multi-targeted type II kinase inhibitors. *J Mol Biol* 2014;426(13):2457–2470
 - 157 Hantschel O, Rix U, Superti-Furga G. Target spectrum of the BCR-ABL inhibitors imatinib, nilotinib and dasatinib. *Leuk Lymphoma* 2008;49(04):615–619
 - 158 Remsing Rix LL, Rix U, Colinge J, et al. Global target profile of the kinase inhibitor bosutinib in primary chronic myeloid leukemia cells. *Leukemia* 2009;23(03):477–485
 - 159 Green MR, Newton MD, Fancher KM. Off-Target Effects of BCR-ABL and JAK2 Inhibitors. *Am J Clin Oncol* 2016;39(01):76–84