Orbitopterional Craniotomy Resection of Pediatric Suprasellar Craniopharyngioma

Devon LeFever¹ Chris Storey¹ Bharat Guthikonda¹

¹Department of Neurosurgery, LSU Health Science Center, Shreveport, Louisiana, United States

Address for correspondence Devon LeFever, MD, Department of Neurosurgery, LSU Health Science Center, 1501 Kings Hwy, Shreveport, LA 71103, United States (e-mail: DLeFev@lsuhsc.edu).

J Neurol Surg B 2018;79(suppl S3):S254–S255.

Abstract

The orbitopterional approach provides an excellent combination of basal access and suprasellar access. This approach also allows for less brain retraction when resecting larger suprasellar tumors that are more superiorly projecting due to a more frontal and inferior trajectory. In this operative video, the authors thoroughly detail an orbitopterional craniotomy utilizing a one-piece modified orbitozygomatic technique. This technique involves opening the craniotomy through a standard pterional incision. The craniotomy is performed using the standard three burr holes of a pterional approach; however, the osteotomy is extended anteriorly through the frontal process of the zygomatic bone as well as through the supraorbital rim. In this operative video atlas, the authors illustrate the operative anatomy, as well as surgical strategy and techniques to resect a large suprasellar craniopharyngioma in a 4-year-old male. Other reasonable approach options for a lesion of this size would include a standard pterional approach, a supraorbital approach, or expanded endoscopic transsphenoidal approach. The lesion was quite high and thus, the supraorbital approach may confine access to the superior portion of the tumor. While recognizing that some groups may have chosen the endoscopic expanded transsphenoidal approach for this lesion, the authors describe more confidence in achieving the goal of a safe and maximal resection with the orbitopterional approach.

The link to the video can be found at: <https://youtu.be/eznsK16BzR8>.

Keywords

- craniopharyngioma
- orbitopterional
- pediatric neurosurgery
- suprasellar mass
- orbitozygomatic

Conflict of Interest
None.

Funding
None.

www.thieme.com/skullbasevideos

www.thieme.com/jnlsbvideos

received
October 15, 2017
accepted
December 14, 2017
published online
February 26, 2018

DOI <https://doi.org/10.1055/s-0038-1624589>.
ISSN 2193-6331.

© 2018 Georg Thieme Verlag KG
Stuttgart · New York

License terms

Fig. 1 Preoperative (A) and postoperative (B) sagittal T1 contrast-enhanced images demonstrating the resection of a larger suprasellar and superiorly projecting craniopharyngioma.

Fig. 2 Intraoperative still image demonstrating the left optic nerve (ON), optic tract (OT), optic chiasm, and internal carotid artery (ICA). The tumor wall is noted in the prechiasmatic corridor and a portion of the cyst wall is seen in the retrochiasmatic corridor.