Combined Interhemispheric and Transsylvian Approach for Resection of Craniopharyngioma

Tomohiro Inoue¹ Hideaki Ono² Akira Tamura² Isamu Saito²

¹Department of Neurosurgery, NTT Medical Center Tokyo, Tokyo, Japan

²Department of Neurosurgery, Fuji Brain Institute and Hospital, Fujinomiya Shi, Japan

Address for correspondence Tomohiro Inoue, MD, PhD, Department of Neurosurgery, NTT Medical Center Tokyo, 5-9-22 Higashi Gotanda Shinakawa-ku, Tokyo 141-0022, Japan (e-mail: t.inouenttsu@gmail.com).

J Neurol Surg B 2018;79(suppl S3):S251.

Abstract

Keywords

- interhemispheric approach
- lamina terminalis
- pterional approach
- craniopharyngioma

We present a 37-year-old male case of cystic suprasellar huge craniopharyngioma, who presented with significant memory disturbance due to obstructive hydrocephalus. Combined interhemispheric and pterional approach was chosen to resect huge suprasellar tumor. Interhemispheric trans-lamina terminalis approach was quite effective to resect third ventricular tumor, while pterional approach was useful to dissect tumor out of basilar perforators and stalk.

The link to the video can be found at: <https://youtu.be/BoYIPa96kdo>.

Conflict of Interest
None.

www.thieme.com/skullbasevideos

www.thieme.com/jnlsbvideos

received
September 18, 2017
accepted
December 18, 2017
published online
February 20, 2018

DOI <https://doi.org/10.1055/s-0038-1625968>.
ISSN 2193-6331.

© 2018 Georg Thieme Verlag KG
Stuttgart · New York

License terms