

Inclusion Criteria: Age between 18 and 65 years. Patients undergoing clipping, coiling for anterior circulation aneurysm with WFNS grade III, IV, and V. ICU admission within 48 to 72 hours of surgery.

Exclusion Criteria: Patient not giving consent. Contraindication for application of NIRS sensors (pneumocephalus). Data of rSO_2 , $SjvO_2$, and CBFV were collected at the following time points—arrival in the ICU (baseline), second hourly daily during the ICU stay (up to 5 days) prior to and after intervention. The significant changes were recorded.

Results: Injured side NIRS values were less compared with normal side. Injured side TCD values were always higher by > 25 cm/sec. The side-side differences existed for NIRS and TCD. The trend values were similar in all patients and consistent.

Conclusion: Interhemispheric differences in rSO_2 and TCD could be considered as an early sign of VS-induced DCI rather than absolute threshold values. However, large studies are needed to test our observations.

Keywords: rSO_2 , TCD, $SjvO_2$, vasospasm

References

1. Mutoh T, Ishikawa T, Suzuki A, Yasui N. Continuous cardiac output and near-infrared spectroscopy monitoring to assist in management of symptomatic cerebral vasospasm after subarachnoid hemorrhage. *Neurocrit Care* 2010;13(3):331–338
2. Poon WS, Wong GK, Ng SC. The quantitative time-resolved near infrared spectroscopy (TR-NIRS) for bedside cerebrohemodynamic monitoring after aneurysmal subarachnoid hemorrhage: can we predict delayed neurological deficits? *World Neurosurg* 2010;73(5):465–466

A003 Anesthetic Consideration in a Case of Von Hippel–Lindau Disease for Cervical Intramedullary Cyst Excision

Hitesh G. Nathani,¹ Bhoomika Thakore,¹ Joseph Monteiro,¹ Hetal Rathod¹

¹Department of Neuroanaesthesia, P. D. Hinduja Hospital, Mumbai, Maharashtra, India

Introduction: Von Hippel–Lindau (VHL) disease is an inherited autosomal dominant disorder with predilection to develop characteristic tumors in the cerebellar hemispheres, retina, brain stem, and spinal cord with a prevalence of approximately 1/50,000.

Methodology/Description: A 29-year-old, 55-kg woman presented with tingling and numbness in all four limbs, difficulty in walking, constipation, and urge for micturition. MRI revealed multiple cystic enhancing lesions in cervical and dorsal cord with significant cord edema. Diagnosed with VHL disease, pancreatic cyst, and diabetes mellitus for past 5 years on treatment. Multiple hemangiomas in retina. Operated for right partial nephrectomy 5 years back. Patient had a Glasgow Coma Scale (GCS) of 15/15 preop. Power of 4/5 on right side, 5/5 on left side. Routine hematological, biochemical values, thyroxine, cortisol, chest X-ray, serum electrolytes, calcium and phosphorus levels, 12-lead electrocardiogram, and echocardiogram were normal. Patient was posted for

cervical laminectomy and intramedullary cyst excision from C2 to C7 level. Routine monitors were attached, left radial artery was cannulated. Patient induced with inj. midazolam, fentanyl, propofol, and atracurium. Anesthesia maintained with oxygen–air mixture with sevoflurane (<0.5 MAC), dexmedetomidine, and propofol infusion. Intraop neuromonitoring—somatosensory evoked potential (SSEP), motor evoked potential (MEP), bispectral index, neuromuscular monitoring done. MEP—amplitude decreased < 50 intraoperatively with baseline values, while SSEP were maintained. Postoperatively, patient had right upper and lower limb power grade 0/5, which improved significantly by POD-3 to preop level 4/5. Patient was discharged on POD-8.

Conclusion: There is very little literature available for anesthetic management in a VHL disease. Anesthetic management of VHL is by itself a challenge; this case was more challenging due to the presence of cervicodorsal and pancreatic cyst with multiple hemangiomas. The case was successfully managed by vigilant clinical and invasive hemodynamic and neuromonitoring with successful outcome.

Keywords: von Hippel–Lindau, anesthetic consideration, neuromonitoring

References

1. Murthy T, Pratyush G, Prabhakar T, Singh P, Mohan C. Anaesthetic implication of von Hippel Lindau disease. *Med J Armed Forces India* 2006;62(2):181–183
2. Mugawar M, Rajender Y, Purohit AK, Sastry RA, Sundaram C, Rammurti S. Anesthetic management of von Hippel–Lindau syndrome for excision of cerebellar hemangioblastoma and pheochromocytoma surgery. *Anesth Analg* 1998;86(3):673–674

A004 Novel Use of Transcranial MEP and SSEP in a Case of Sciatic Nerve Tumor

Nitin Manohar,¹ Anand Balasubramaniam¹

¹Department of Neuroanesthesia, Yashoda Hospitals Secunderabad, Hyderabad, Telangana, India

Introduction: Sciatic nerve monitoring is commonly done in pelvic surgeries and total hip replacements to preserve sciatic nerve function and prevent foot drop.

Methodology/Description: We present a case of a 24-year-old woman with severe pain in left posterior thigh extending into the leg and foot. MRI revealed sciatic nerve tumor (neurofibroma) extending into the sciatic foramen. We planned for intraoperative neurophysiological monitoring to preserve her sciatic nerve function. As the tumor was extending into the sciatic foramen, we could not stimulate the proximal nerve or do nerve action potentials. So, we planned for transcranial (motor evoked potentials [MEPs]). MEP monitoring in the muscles supplied by sciatic nerve (biceps femoris, gastrocnemius, tibialis anterior, extensor hallucis longus, and abductor hallucis longus). Posterior tibial somatosensory evoked potentials (SSEPs) were also used to monitor sensory component. TIVA with entropy guidance using propofol and dexmedetomidine was used to maintain anesthesia. For control, opposite limb MEP and SSEP were also monitored. During